

Saunders' Question Compend

ESSENTIALS OF
MATERIA MEDICA AND THERAPEUTICS
HENRY MORRIS M.D.


RM

126

M88







SEP 8 1898

ESSENTIALS
OF
MATERIA MEDICA, THERAPEUTICS, AND
PRESCRIPTION-WRITING.

SINCE the issue of the first volume of the
Saunders Question-Compends,

OVER 160,000 COPIES

of these unrivalled publications have been sold.
This enormous sale is indisputable evidence
of the value of these self-helps to students
and physicians.

SAUNDERS' QUESTION-COMPENDS, No. 7.

ESSENTIALS OF MATERIA MEDICA, THERAPEUTICS,
AND PRESCRIPTION WRITING

ARRANGED IN THE FORM OF

QUESTIONS AND ANSWERS

PREPARED ESPECIALLY FOR

STUDENTS OF MEDICINE.

BY

HENRY MORRIS, M.D.,

Fellow of the College of Physicians of Philadelphia; Honorary Member of the Altoona Academy of Medicine; Associate Member of the Association of Military Surgeons of the United States; Member of the American Medical Association, of the Pennsylvania State Medical Society, of the Philadelphia County Medical Society, of the Pathological Society of Philadelphia; Physician to St. Joseph's Hospital.

FIFTH EDITION, REVISED AND ENLARGED.

PHILADELPHIA:

W. B. SAUNDERS,

925 WALNUT STREET.

1898.

RM. 26
.M88

12653

Entered according to Act of Congress, in the year 1898, by

W. B. SAUNDERS.

In the Office of the Librarian of Congress, at Washington.

SEP 8 1898



TWO COPIES RECEIVED

PRESS OF
W. B. SAUNDERS, PHILADA.

43142
July 19,
1898.

Recd., M. W. J., M.D., 20-22

To the Memory of
DR. THEODORE HORWITZ,
WHOSE EARLY DEATH
DEPRIVED THE MEDICAL PROFESSION OF AN EARNEST
AND TALENTED WORKER,
THIS BOOK IS AFFECTIONATELY INSCRIBED
BY HIS FRIEND,
THE AUTHOR.

PREFACE TO THE FIFTH EDITION.

IN the preparation of this edition it has been the object of the author not only to follow closely the last edition of the U. S. Pharmacopœia, but also to bring the book, as nearly as possible, thoroughly abreast of the times. Much that was obsolete has therefore been omitted, and many new drugs, especially among the groups of antipyretics and hypnotics, have been added, while the older subject-matter has been carefully revised and corrected.

In deference to the criticisms of English and Canadian medical journals, the dosage of each preparation has been expressed in the metrical weights and measures as well as in apothecaries' weight and wine measure. This change has been rendered necessary also by the adoption of the metrical system by the Medical Corps of the United States Army, many of whose officers have in the past used, and I hope in the future will continue to use, this little book for reference.

One word in regard to former criticisms, which on the whole have been most kind and considerate : The author is criticised for his classification of medicines. He thinks, with Lauder Brunton, that the time has come in scientific medicine when an attempt at classification, however imperfect and tentative, should be made, instead of giving up the entire subject as hopeless, and arranging the remedies in alphabetical order ; in which case, probably, as in a book before him, remedies as diverse in their effects and uses as *croton oil*, *cubebs*, *cyanide of potassium*, *digitalis*, *elaterium*, and *ergot* would follow each other, to the endless confusion of the student, who knows that the only way he can

master a subject thoroughly is by systematizing his knowledge and his method of study, which is hardly possible if he learns in succession, and without any connection with each other, the effects and uses of *oxide of zinc*, *oxygen*, *pancreatin*, *paraldehyd*, and *pareira*, to take a few examples of such classification at haphazard. While admitting that his classification is not free from faults, the author regards it as much preferable to the examples quoted.

By adopting a classification based on a therapeutical rather than a physiological method, the entire book is a therapeutical index; and while the names of diseases do not appear in the table of contents, it is in reality, from its arrangement, an index to the remedies adapted to the treatment of various diseased conditions.

The author acknowledges his indebtedness to numerous standard text-books, as the National Dispensatory; the U. S. Pharmacopœia; Wood's, Hare's, Lauder Brunton's, Foster's, and Wilson's Therapeutics; Maisch's Organic Materia Medica; Bloxam's Chemistry; Blythe on Poisons; as well as various medical annuals, as those of Sajous, Gould, the International Medical Annual; and to writers in the various medical journals, whose papers, though too numerous to mention, form the great bulk of new matter in this edition.

The author takes pleasure in thanking Dr. Henry C. Haden for his valuable work in gathering references and abstracts; and his acknowledgments are also due to Miss Caroline M. Irwin for the exhaustive and thorough index, which he is sure will add much to the value of this edition.

In conclusion, he renews the dedication to the memory of his friend, the late Dr. Theodore Horwitz, and while thanking the profession and critics for their kind reception of former editions, expresses the hope that they will find the present edition equally worthy of their confidence.

HENRY MORRIS.

313 S. 16th St., Philadelphia.

PREFACE.

The object of this little book is, not to replace or attempt to replace with the student the larger text-books on *Materia Medica*, but simply to give him, in condensed form, the principles of this branch of medicine, without a knowledge of which he can neither pass his final examinations, nor practice his profession intelligently.

The author hopes that, if properly used, this book will be of service to the student and young practitioner, but he is sure, from his experience as a teacher, that neither this nor any other "compend" will suffice to form the groundwork of what is really the study of a lifetime—the Science of Medicine.

In regard to the classification of drugs, he would say that this work was commenced before the excellent arrangement contained in the last edition of Prof. Wood's *Therapeutics* was published, and not being satisfied with any of the groupings which he had seen, he was induced to adopt a classification based on therapeutic rather than physiological grounds, which, however, he is pleased to think agrees very nearly with that of the latest edition of Wood. In conclusion he must acknowledge his indebtedness not only to the standard text-books, such as the admirable works of Wood, Bartholow, Ringer, Lauder Brunton and many others, but also to many writers in the medical journals, both of this and other countries, to whom want

of space has prevented him from giving due credit in the appropriate place. He must also acknowledge the kindness of his friend, Dr. Henry D. Moore, to whom he is indebted for the preparation of the index.

Believing that the critic is the best friend of the medical author, he will thankfully receive any criticisms or hints from either public or private sources.

HENRY MORRIS.

313 South 16th St., Philadelphia.

TABLE OF CONTENTS.

	PAGES
DEFINITIONS. —(Materia Medica, Pharmacy, Pharmacology, Therapeutics, Remedies, Pharmacopœia, Dispensatory, Official preparations),	17-23
WEIGHTS AND MEASURES. —(Troy or Apothecaries' weights, Wine or Apothecaries' measure, Metrical or decimal system of weights, Tables showing the value of Apothecaries' weights in the metrical system, Table showing the value of Apothecaries' measure in the metrical system, Approximate measures),	23-26
PRESCRIPTION WRITING. —(Purposes for which medicines are combined, Precautions to be observed in prescription writing, Component parts of a prescription, Method of writing a prescription, Calculation of dose, General rules for dosage), . . .	26-30
INFLUENCES THAT MODIFY EFFECTS OF MEDICINES. —(Age, Sex, Idiosyncrasy, Constitution, Habitual use, Condition of digestive and absorbent systems, Diseases, Mode of administration),	30-32
ADMINISTRATION OF MEDICINES. —(Enepidermic, Epidermic, Endermic, and Hypodermic methods, By the mucous membranes, By intravenous injection),	32
CLASSIFICATION OF MEDICINES,	33, 34
GROUP I.—SYSTEMIC MEDICINES,	35-244
CLASS I.—MEDICINES ACTING ON THE BODY AT LARGE, BY THEIR INFLUENCE ON DIGESTION, NUTRITION (tissue waste and repair), AND CALORIFICATION, . .	35-90
ORDER 1.— <i>Digestants</i> —(Pepsin, Ingluvin, Pancreatin, Papain),	35-37
ORDER 2.— <i>Tonics</i> —(Quassia, Gentian, Calumba, Chirata, Serpentaria. Eucalyptus, Prunus Virginiana, Anthemis,	

	PAGES
Matricaria, Eupatorium, Cascarilla, Absinthium, Cinchona and its alkaloids, Hydrastis, Preparations of Iron and of Manganese, Phosphorus, Mineral acids, Lactic acid), . . .	37-59
ORDER 3.— <i>Alteratives</i> —(Preparations of Mercurv and of Gold, Iodine, the Iodides, Arsenical preparations, Cod-liver oil, the Phosphates and Hypophosphites, Calcium chloride, Ammonium chloride, Colchicum, Sarsaparilla, Guaiac, Mezereum, Menispermum, Calendula, Sassafras, Stillingia, Dulcamara, Lappa),	59-77
ORDER 4.— <i>Antipyretics</i> —(Antipyrine, Acetanilid, Phenacetin, Phenocoll, Quinine, Salicin, the Salicylates, Salol, Salophen, Guaiacol, Resorcin, Hydroquinone, Pyrocatechin, Chinoline, Thalline, Kairine),	77-90
CLASS II.—MEDICINES ACTING ON THE BODY, BY THEIR	
EFFECTS ON THE VARIOUS ORGANS,	90-202
(a) ON THE NERVOUS SYSTEM,	90-146
ORDER 1.— <i>Hypnotics</i> —(Opium and its alkaloids, Chloral, Chloralamide, Chloralose, the Bromides, Hyoscyne, Sulphonal, Trional, Tetronal, Paraldehyd, Urethan, Cannabis Indica, Humulus, Lactucarium),	90-107
ORDER 2.— <i>Mydriatic Anodynes</i> —(Belladonna, Stramonium, Hyoscyamus, Duboisia, Coca, and their alkaloids, Antipyrine),	107-116
ORDER 3.— <i>Anæsthetics</i> —(Ether, Chloroform, Nitrous oxide, Methylene bichloride, Ethyl bromide, Bromoform). <i>Local Anæsthetics</i> —(Cold, Rhigolene, Ether, Ethyl chloride, Cocaine, Eucaïne, Menthol, Guaiacol, Tincture of aconite),	116-126
ORDER 4.— <i>Antispasmodics</i> —(Musk, Valerian, the Valerianates, Asafetida, Camphor, Monobromated camphor, Ethereal oil, Hoffmann's anodyne, Cypripedium, Scutellaria), .	126-130
ORDER 5.— <i>Ercito-motors</i> —(Nux vomica and its alkaloids, Rhus toxicodendron, Picrotoxin),	130-134
ORDER 6.— <i>Depresso-motors</i> —(Physostigma, Chloral, the Bromides, the Nitrites, Gelsemium, Prussic acid, the Cyanides, Oil of Bitter Almond, Lobelia, Conium, Woorara, Tobacco),	134-146

PAGES

(b) ON THE ORGANS OF CIRCULATION, 146-169

ORDER 1.—*Stimulants*—(Alcohol, Ether, Ammonium, Amyl nitrate, Nitroglycerin, Atropine, Digitalis, Caffeine, Adonidine, Strophanthus, Sparteine, Strychnine, Cimicifuga, Convallaria), 146-159

ORDER 2.—*Sedatives*—(Antimonial preparations, Aconite and its alkaloids, Veratrum Viride, Veratrine, Gelsemium, Pulsatilla, Arnica, the Vegetable acids), 159-169

(c) ON THE SEXUAL ORGANS, 169-177

ORDER 1.—*Aphrodisiacs*—(The Tonics generally, especially Phosphorus, Zinc phosphide, and Iron. The Phosphates and Hypophosphites, Gold, Arsenic, Cod liver oil, Opium, Alcohol, Cannabis Indica, Musk, Camphor, Nux vomica, Stimulants, Hamamelis, Ergot, Cantharides, Myrrh), 169

ORDER 2.—*Anaphrodisiacs*—(Nauseants, Purgatives, Opium (full doses), the Bromides, Monobromated camphor, Iodine and the Iodides, Potassium nitrate, Ammonium chloride, Belladonna, Tobacco, Conium, Cardiac sedatives, Digitalis, Cocaine), 170

ORDER 3.—*Oxytocics*—(Ergot, Hydrastis, Hydrastinine hydrochlorate, Cotton-root bark, Quinine), 170-174

ORDER 4.—*Uterine Sedatives*—(Opium, Cannabis Indica, the Bromides, Chloral, Tartar emetic, Ether, Viburnum prunifolium), 174-175

ORDER 5.—*Emmenagogues*—(Preparations of Iron, of Arsenic, and of Manganese, Myrrh, Aloes, the Salines, Potassium permanganate, Savine, Tansy, Apiol, Cantharides, Guaiac), 175-177

(d) ON THE ALIMENTARY CANAL, 177-202

ORDER 1.—*Emetics*—(Ipecac, Apomorphine, Mustard, Zinc sulphate, Copper sulphate, Alum, Turpeth mineral, Tartar emetic), 177-181

ORDER 2.—*Gastric Sedatives*—(Calomel, Ice, Lime-water, Brandy, Morphine, Diluted hydrocyanic acid, Creasote, Bismuth, Carbolic acid, Thymol, Sulphurous acid, the Sulphites, Charcoal, the Bromides, Nux vomica, Cereum

	PAGES
oxalate, Ipecac (minute doses), Fowler's solution, Counter-irritation),	181-182
ORDER 3.— <i>Carminatives</i> —(Asafetida, Eucalyptus, Ether, Capsicum, Pepper, Cinnamon, Nutmeg, Mace, Cloves, Allspice, Ginger, Cardamoms, Aromatic powder, Aromatic fluid extract, Oil of cajuput, Calamus, Oil of gaultheria, Oil of lavender flowers, Peppermint, Spearmint, Oil of rosemary, Pennyroyal, Horehound, Sage, Oil of thyme, Fennel, Caraway, Anise, Coriander, Star anise, Orange-peel, Orange-flowers, Vanilla, Oil of turpentine),	182-188
ORDER 4.— <i>Cathartics</i> —(Tamarinds, Manna, Cassia fistula, Frangula, Cascara sagrada, Castor oil, Sulphur, Sulphurated potassa, Rhubarb, Juglans, Aloes, Senna, Lepandra, Euonymus, Mercurials, Magnesia, Magnesium carbonate, Epsom salt, Effervescent magnesium citrate and its solution, Glauber's salt, Sodium phosphate and pyrophosphate, Potassium sulphate, Potassium bitartrate, Rochelle salt, Seidlitz powder, Manganese sulphate, Jalap, Bryony, Podophyllum, Iris, Tetterwort, Scammony, Colocynth, Gamboge, Compound cathartic pills, Vegetable cathartic pills, Elaterin, Croton oil, Enemata),	188-199
ORDER 5.— <i>Anthelmintics</i> —(Spigelia, Chenopodium, Santonica and Santonin, Aspidium, Pomegranate and Pelletierine, Cusso, Pumpkin-seed, Kamala, Oil of turpentine, Calomel, Sulphur, Infusion of quassia, Sodium chloride, Naphtol, Tannin, Alum-water, Lime-water, Injections of sweet oil),	199-202
CLASS III.—MEDICINES ACTING ON THE BODY BY THEIR EFFECT ON THE VARIOUS SECRETIONS AND EXCRETIONS,	202-244
ORDER 1.— <i>Diuretics</i> —(Solution of potassa, Potassium carbonate, Potassium bicarbonate, Potassium citrate, Potassium acetate, Cream of tartar, Potassium nitrate, Potassium chlorate, Lithium carbonate, Lithium citrate, Lithium benzoate, Sodium acetate, Strontium lactate, Squill, Digitalis, Caffeine, Sodio-theobromine salicylate, Apocynum, Broom, Dandelion, Sweet spirit of nitre, Mercurials, Buchu, Pareira, Uva ursi, Chimaphila, Triticum, Juniper, Oil of erigeron, Oil of santal, Corn-silk, Turpentine, Copaiba, Cubeb, Matico, Cantharides),	202-217

ORDER 2.—*Diaphoretics*—(Ipecac, Dover's powder, Antimonials, Aconite, Veratrum viride, Salicylic acid, Potassium citrate, Spirit of mindererus, Sweet spirit of nitre, Pilocarpus), 217-221

ORDER 3.—*Expectorants*—(Ipecac, Apomorphine, Tartar emetic, Pilocarpine, Lobelia, Grindelia, Ammonium chloride, Senega, Quillaja, Quebracho, Asclepias, Eriodictyon, Garlic, Sanguinaria, Ammoniac, Benzoin, Peruvian balsam, Balsam of tolu, Tar, Terpin hydrate, Terebene, Squill, Turpentine, Eucalyptus, Serpentaria, Prunus Virginiana), 221-228

ORDER 4.—*Astringents*—(Tannic acid, Gallic acid, Nutgalls, Catechu, Kino, Krameria, Logwood, White oak bark, Geranium, Witch-hazel, Red and pale rose, Sumach, Chestnut leaves, the Lead salts, Alum, Aluminum sulphate, the Salts of copper, of zinc, of silver, and of bismuth, Cerium oxalate), 228-241

ORDER 5.—*Antacids*—(Solution of potassa, Spirit and aromatic spirit of ammonia, Solution of soda, Sodium carbonate, Sodium bicarbonate, Lime-water, Syrup of lime, Precipitated calcium carbonate, Prepared chalk and its preparations), 241-244

GROUP II.—TOPICAL MEDICINES, 244-269

CLASS I.—*ANTISEPTICS*.—(Solution of hydrogen dioxide, Potassium permanganate, Chlorine-water, Chlorinated lime, Labarraque's solution, Mercuric chloride, Bromine, Iodine, Mercuric oxide, Iodoform, Iodol, Sulphurous acid, the Sulphites and sulphides, Boric acid and the borates, Iron sulphate, Lead nitrate, Carbolic acid and the carbolates, Creasote, Benzoic acid and the benzoates, Naphtalin, Naphtol, Formaldehyd, the Antipyretics, Antiseptic oils, Thymol), . 244-256

CLASS II.—*IRRITANTS*, 256-262

ORDER 1.—*Rubefacients*—(Mustard, Capsicum, Oil of turpentine, Ammonia, Burgundy pitch, Canada pitch), . . 256-258

ORDER 2.—*Vesicants*—(Cantharides, Potáto-bug, Ammonia), 258-259

ORDER 3.—*Pustulants*—(Croton oil, Antimonial ointment), 259-260

	PAGES
ORDER 4.— <i>Escharotics</i> —(Caustic potash, Caustic soda, Arsenious acid, Bromine, Zinc chloride, Acid nitrate of mercury, Corrosive sublimate, Potassium bichromate, the Mineral acids, Silver nitrates, Copper sulphate, Alum), . .	260-262
CLASS III.— <i>DEMULCENTS</i> —(Water, Acacia, Tragacanth, Slippery elm-bark, Marshmallow, Flax-seed, Sassafras pith, Quince-seed, Liquorice-root, Iceland moss, Irish moss, Starch, Isinglass),	263-266
CLASS IV.— <i>EMOLLIENTS</i> —(Lard, Lanolin, Suet, Spermaceti, Wax, Oleic acid, Glycerine, Cacao-butter, Cotton-seed oil, Oil of benne, Petrolatum, the Demulcents), . . .	266-267
CLASS V.— <i>PROTECTIVES</i> —(Plasters, Ointments, Plaster of Paris, Collodion, India-rubber, Solution of sodium silicate),	268-269
CLASS VI.— <i>COLORING AGENTS</i> —(Cochineal, Saffron, Red saunders),	269
INDEX,	271

QUESTIONS AND ANSWERS

ON THE

ESSENTIALS OF MATERIA MEDICA AND THERAPEUTICS.

What is meant by the term “materia medica”?

Materia medica is that division of medical science which treats of all that relates to the medicines employed in the treatment of disease. It embraces the study of the source of remedies, their physical and chemical properties, their action in medicinal doses on the human body both in health and disease, their poisonous effects and the means by which they may be antagonized, and the methods of preparing and administering them; and is closely allied to botany, mineralogy, zoölogy, chemistry, physiology and pathology.

How may this knowledge be subdivided?

Into *Materia Medica Proper*, *Pharmacy*, *Pharmacology* including *Toxicology*, and *Therapeutics*.

What is materia medica proper?

By materia medica proper is understood a description of the source, and the physical and chemical characteristics of remedies.

What is pharmacy?

Pharmacy (φάρμακον, a medicine) relates to the preparation and combination of remedies.

What is pharmacology?

Pharmacology (φάρμακον, a medicine, and λόγος, a discourse), though originally meaning a “treatise on drugs” or “on pharmacy,” is now usually applied only to the study of the action of drugs on the body, and embraces toxicology (τοξικόν, poison with

which arrows are smeared, and λόγος, a discourse), which deals with poisons and their antidotes.

What are therapeutics?

Therapeutics (Θεράπεια, I take care of, or, cure) deals with the application of remedies to disease, and may be either *empirical* or *rational*.

How does empirical differ from rational therapeutics?

When a remedy is given in any disease because it has succeeded in similar cases before, although the manner in which it acts is often not understood, it is given empirically. When a remedy is prescribed because, from a knowledge of its action on the body, it is likely that it will antagonize or overcome a particular pathological condition, it is administered on the rational plan. Rational therapeutics is the more scientific, and is the outgrowth of pharmacology.

What are remedies?

Remedies are the agents employed in the treatment of disease.

How may remedies be classified?

Remedies are subdivided into *mechanical*, *physical*, *dietetic*, *hygienic* and *pharmaceutical* agents. The first four classes are usually treated of in works especially devoted to such subjects.

What are mechanical remedies?

Venesection; local bloodletting by means of leeches, cups, scarifications, incisions or punctures; bandages; splints; sutures; ligatures; acupressure needle and forceps; acupuncture; aspiration; frictions; massage, etc., are mechanical remedies and are described in works on surgery.

What are physical remedies?

When light, heat or cold, electricity, galvanism or other of the physical forces are used in the treatment of disease, they are spoken of as *physical remedies*. They are treated of at length in works on physics and in books devoted to them exclusively.

What is meant by dietetic remedies?

All relating to the food of the sick* or to special plans of diet, are included under this head.

* In many diseases the diet is of equal or of more importance than medication, e. g., in diabetes mellitus, and in various forms of dyspepsia.

What are hygienic remedies?

The clothing, bathing and cleanliness of the sick, the ventilation of the sick room, etc., belong to this subdivision, and will be found in works on practice of medicine, but more particularly in those relating to hygiene.

What are pharmaceutical remedies?

Pharmaceutical remedies or "medicines, are substances not essentially alimentary, which, when applied to the body, so alter or modify its vital functions as to be rendered applicable to the treatment of diseases." * It is preëminently with this subdivision that *materia medica* deals.

What is meant by "official" preparations?

This term is applied to such remedies as are authorized (*officialis*) by the Pharmacopœia.

How does a pharmacopœia differ from a dispensatory?

A *pharmacopœia* (φάρμακον, a medicine, and ποίεω, to make) is a book containing directions for the preparation of medicines with the object of uniformity of nomenclature, preparation and strength. The U. S. Pharmacopœia has been decennially revised since 1820, when it was first authorized by a representative convention of druggists and physicians, held at Washington. It is an authoritative list of remedies, with formulæ for such preparations as may be in general use. "A *dispensatory* differs from a pharmacopœia in containing the physical and medical history of the various substances." It "contains the whole of the pharmacopœia" "with additions by the author, under whose authority alone it usually appears" (Dunglison). It treats, also, of the therapeutics, modes of administration and doses of the various drugs, "while the pharmacopœia is restricted to the mode of preparing them."

How may the official preparations be classified?

They may be classified, according to their physical state, into *solids*, *semi-solids*, and *liquids*.

Name the official solid preparations.

Lozenges, papers, pills, powders, and triturations.

* Biddle's *Materia Medica*, 11th Ed., 1889, p. 46.

What are the official semi-solid preparations?

Confections, extracts, cerates, liniments, oleates, ointments, plasters, and suppositories.

Mention the official liquid preparations.

Solutions, mixtures, waters, infusions, decoctions, tinctures, spirits, wines, glycerites, fluid extracts, honeys, syrups, vinegars, and oleo-resins.

What are lozenges?

Lozenges or troches (*trochisci*) are dry, solid medicines made of powders incorporated usually by means of mucilage and sugar, intended to be slowly dissolved in the mouth.

What are papers?

Papers (*chartæ*) are made by soaking bibulous paper in medicinal solutions (*e.g.*, charta potassii nitratis), or by spreading medicinal mixtures on paper (*e.g.*, charta sinapis).

What are pills?

Pills (*pilule*) are preparations of such size that they can be conveniently swallowed whole. They usually contain some excipient to give the proper degree of tenacity to their particles, and are rolled into a spherical form.

What are powders?

Powders (*pulveres*) are dry solid medicines in a minute state of subdivision.

What are triturations?

Triturations (*triturationes*) consist of 1 part of a medicinal substance and 9 parts of sugar of milk thoroughly mingled and reduced to a fine powder in a mortar.

What is a confection?

Confections or conserves (*confectiones*) are remedies made into a soft mass with sugar, honey, etc., which render them more pleasant to the taste.

What are extracts?

Extracts (*extracta*) are made by evaporating solutions of vegetable substances (or in some cases the juices expressed from plants)

to a pilular consistence. They are called aqueous or watery, alcoholic and acetous extracts, according to the menstruum used as a solvent

What are cerates?

Cerates (*cerata*) are unctuous preparations, intended for external application, mixed with wax, spermaceti or resin, which gives them a firm consistence, and prevents them from melting at the temperature of the body.

What are liniments?

Liniments (*linimenta*) are oleaginous preparations, intended for external application with friction.

What are oleates?

Oleates (*oleata*) are solutions in oleic acid of metallic salts or alkaloids, and are applied externally.

What are ointments?

Ointments or unguents (*unguenta*) are medicaments incorporated with some fatty substance, and intended for external use. They are of the consistence of lard.

What are plasters?

Plasters (*emplastra*) are preparations spread on linen, muslin, silk or sheepskin, and are intended to adhere closely to the surface to which they are applied.

What are suppositories?

Suppositories (*suppositoria*) are made by incorporating medicinal substances with oil of theobroma, and are intended for introduction into the rectum or vagina. They are of conical form, to facilitate their insertion.

What are solutions?

Solutions (*liquores*) are preparations in which a non-volatile substance is entirely dissolved in the menstruum (usually water).

What are mixtures?

Mixtures (*mixture*) are preparations in which, by means of some viscid substance, an insoluble medicine is suspended in water.

What are waters?

Waters (*aque*) are aqueous solutions of gaseous or volatile substances.

What are infusions?

Infusions (*infusa*) are aqueous preparations obtained by *pouring* hot or cold water on vegetable substances and allowing it to remain sufficiently long to extract the virtues of the drug.

What are decoctions?

Decoctions (*decocta*) are aqueous preparations obtained by *boiling* vegetable substances in water for a few minutes and allowing them to cool in closely-covered vessels.

What are tinctures?

Tinctures (*tincturae*) are alcoholic solutions, usually of non-volatile substances. Alcohol or diluted alcohol is generally used as a solvent, but occasionally the aromatic spirits of ammonia is employed, the product being called an *ammoniated tincture*.

What are spirits?

Spirits (*spiritus*) are alcoholic solutions of gaseous or volatile substances.

What are wines?

Wines (*vina*) are tinctures made with stronger white wine instead of alcohol.

What are glycerites?

Glycerites (*glycerita*) are medicinal substances mixed with glycerin.

What are fluid extracts?

Fluid extracts (*extracta fluida*) are liquid preparations of uniform strength, 1 c.c. of which represent 1 Gm. (about 1 minim to the grain) of the crude drug, prepared chiefly with alcohol and glycerin.

What are honeys?

Honeys (*mellita*) are medicines dissolved in honey.

What are syrups?

Syrups (*syrupi*) are composed of *medicinal* substances, or of *flavoring* ingredients in sugar and water. *Simple syrup* is a concentrated solution of sugar in water.

What are vinegars?

Vinegars (*aceta*) are medicinal solutions in diluted acetic acid.

What are oleoresins?

Oleoresins (*oleoresinæ*) are concentrated extracts consisting chiefly of the resin dissolved in the oil of the drug from which they are obtained.

May medicines be prescribed in other forms than those referred to above?

Medicines are often administered in *capsules* (small, elongated gelatine boxes with rounded ends, which dissolve readily in the stomach), *granules* or *parvules* (very minute pills), *emulsions* (oils or resins suspended in a mixture by means of some viscid material), *enemata* (liquids injected into the rectum for their local or systemic effect), or *injections* (liquids forced by a syringe into the mucous passages or cavities, under the skin or into a blood vessel). *Inhalations* of vapors may be employed for their local effect on the broncho-pulmonary tract or for their influence on the system at large, and *gargles* may be used as sedative, astringent or disinfecting and deodorizing applications to the buccal and pharyngeal mucous membranes. Externally, *poultices* (simple or medicated), *lotions* (washes), and *fomentations* (hot lotions applied on flannel or cloth and allowed to remain in contact with the surface), are also often used.

What weights and measures are used in prescribing and dispensing medicines?

Troy or apothecaries' weight is used in dispensing solids, while *wine or apothecaries' measure* is employed for liquids; occasionally the *metrical system* is used in prescribing.

What is apothecaries' weight?

In apothecaries' weight the pound is divided into ounces, drachms, scruples and grains, as follows:—

20 grains (gr.) = 1 scruple (℥).

60 grains or 3 scruples = 1 drachm (ʒ).

480 grains or 24 scruples or 8 drachms = 1 ounce (℥).

5760 grains or 288 scruples or 96 drachms or 12 ounces = 1 pound (lb).

The grain (gr.), drachm (ʒ) and ounce (℥) should alone be used in prescription writing.

What is wine measure ?

In wine measure the gallon is divided into pints, fluid-ounces, fluid-drachms and minims, thus :—

60 minims (℥) = 1 fluid-drachm (ʒ).

480 minims or 8 fluid-drachms = 1 fluid-ounce (ʒ).

7680 minims or 128 fluid-drachms or 16 fluid-ounces = 1 pint (O).

61440 minims or 1024 fluid-drachms or 128 fluid-ounces or 8 pints = 1 gallon (C).

The minim (℥), fluid-drachm (ʒ), fluid-ounce (ʒ) and, very seldom, the pint (O) are used in prescriptions.

Describe the metrical system of weights.

In the *metrical or decimal system* the gramme is multiplied or divided by ten, Greek prefixes denoting the multiplication, and Latin the subdivisions of the unit ; thus :—

10 milligrammes = 1 centigramme.

100 milligrammes, or 10 centigrammes = 1 decigramme.

1000 milligrammes, or 100 centigrammes, or 10 decigrammes = 1 gramme.

10,000 milligrammes, or 1000 centigrammes, or 100 decigrammes, or 10 grammes = 1 decagramme.

100,000 milligrammes, or 10,000 centigrammes, or 1000 decigrammes, or 100 grammes, or 10 decagrammes = 1 hectogramme.

1,000,000 milligrammes, or 100,000 centigrammes, or 10,000 decigrammes, or 1000 grammes, or 100 decagrammes, or 10 hectogrammes = 1 kilogramme.

In writing prescriptions in the metrical system, the gramme (Gm.) alone is used for both solids and fluids, the abbreviation being *preceded* by *Arabic figures* or decimals denoting the amount of the drug (*e. g.*, 10.0 Gm.—0.02 Gm., etc.).

What is a gramme ?

The gramme is the weight of a cubic centimeter (c. c.) of distilled water at 4° C. (39.5° F.).

What is a centimeter ?

A centimeter ($\frac{1}{100}$ of an inch) is the $\frac{1}{100}$ of a metre (39.37 inches), which is the ten-millionth part of a quarter of a meridian of the earth (or of the distance from the pole to the equator).

What relation does the metrical system of weights bear to apothecaries' weight and measure?

1 gramme = 15.5 grains ; 1 decigramme (0.1 Gm.) = 1.5 grains ;
 1 centigramme (0.01 Gm.) = .15 (or $\frac{1}{6}$) grain ; 1 milligramme
 (0.001 Gm.) = .015 (or $\frac{1}{64}$) grain ; 1 grain (gr.j) = .06 gramme ; 1
 drachm (ʒj) = 4.0 grammes (approximately) ; 1 ounce (ʒj) =
 32.0 grammes (approximately).

The following table gives the value of grains in the metrical system. (The fractions have not been carried out beyond the fourth decimal point.)

Grains	Grammes	Grains	Grammes
$\frac{1}{1000}$	= 0.0006	3	= 0.194
$\frac{1}{500}$	= 0.0013	4	= 0.259
$\frac{1}{250}$	= 0.0027	5	= 0.324
$\frac{1}{80}$	= 0.008	10	= 0.648
$\frac{1}{40}$	= 0.016	30 or ʒss	= 1.944
$\frac{1}{20}$	= 0.032	60 or ʒj	= 3.888
$\frac{3}{40}$	= 0.049	120 or ʒij	= 7.775
1	= 0.065	240 or ʒss	= 15.551
2	= 0.130	480 or ʒj	= 31.103

As the weight of a c. c. of fluid depends upon its density, 1.0 Gm. of a light liquid (*e. g.*, alcohol) will measure more than the same weight of a heavy liquid (*e. g.*, glycerin). Approximately, however, m℥ = 0.06 Gm. ; fʒj = 4.0 Gm. (a little less), and fʒj = 32.0 Gm. (a little less).

The following table gives the value of minims (of water at 4° C. in vacuo) in the metrical system. For ordinary purposes these values may be used without correction.

Minims	Grammes	Minims	Grammes
1	= 0.06161	20	= 1.232
2	= 0.123	30 or fʒss	= 1.848
3	= 0.185	60 or fʒj	= 3.696
4	= 0.246	120 or fʒij	= 7.393
5	= 0.308	240 or fʒss	= 14.786
10	= 0.616	480 or fʒj	= 29.573
15	= 0.924	960 or fʒij	= 59.147

What approximate measurements are used in medicine?

Medicines are sometimes ordered in *drops* instead of *minims*. As the size of the drop depends on a variety of circumstances, this is a bad practice. A *teaspoonful* is often used for $f\overline{3}j$, or 5.0 Gm. ; a *tablespoonful* for $f\overline{3}ss$, or 20.0 Gm. ; a *wineglassful* for $f\overline{3}ij$; but these are uncertain measures, and should only be employed when an excess of the quantity intended would be harmless. Graduated medicine glasses are easily obtained, and are much more reliable.

May a physician prescribe other formulæ than those contained in the Pharmacopœia?

The physician often extemporizes formulæ to meet the requirements of the individual case.

For what purposes are different remedies often combined in a single prescription?

Remedies are combined to increase or to moderate the effects of some single medicine, or to overcome some action which it is undesirable to produce ; to procure the simultaneous action of drugs having different effects upon the body, or to form a new chemical compound, often differing in its action from those of the substances forming it ; for convenience in administering, or for rendering nauseous drugs more palatable.

What precautions should be used in prescription writing?

The physician should be careful, while writing the prescription,—

1. To bear constantly in mind the object for which the medicine is wanted, in order that he may better combine his remedies to accomplish his purpose.
2. The writing should be clear and distinct, and the words not so abbreviated as to be unintelligible or susceptible of more than one meaning. Attention to these points will prevent the occurrence of mistakes in compounding the medicine.
3. The doses should neither be large enough nor close enough together (in the case of powerful medicines) to cause injurious effects. It is better to give too little than too much.
4. Medicines which form poisonous compounds (as calomel and sodium chloride) should not be combined in the same prescription.
5. Remedies should not be combined (except for particular pur-

poses) which are physiological antagonists, *i. e.*, which by their actions on the body render each other inert.

6. Remedies should not be combined for internal administration which are chemically incompatible, *i. e.*, which will decompose or precipitate each other.

7. The prescription should be as simple as possible, and contain as few ingredients as the nature of the case will allow.

8. It is better not to order a larger quantity of medicine than the patient will probably require.

9. Never send out a prescription without having carefully reread it, to be certain no mistake has been made.

10. Never give a patient a prescription which has been corrected. If a mistake has been made, rewrite the prescription entirely.

Of what parts does a prescription consist?

A prescription consists of four parts :—

1st. The *superscription* (*super*, “over,” and *scribo*, “I write”), which consists of the sign *R* (which is possibly an abbreviation for *Recipe*, “take”), and in some cases the name and address of the patient, with the date.

2d. The *inscription*, the body of the prescription, containing the names of the drugs which are to be used, with the amount of each.

3d. The *subscription* (*sub*, “under,” and *scribo*, “I write”), the directions to the apothecary regarding the preparation of the medicine.

4th. The *signature*, commencing with *S.* or *Sig.* (*Signa*, “write”), and including directions for taking the medicine, the physician’s name, and frequently the date, which may be placed here instead of at the beginning of a prescription.

It is customary to write the superscription and signature in English and the inscription and subscription in much abbreviated *medical Latin*.

The *inscription* formerly consisted almost universally of a *basis* (the principal medicine), an *adjuvant* (to aid its action), a *corrective* (to prevent unpleasant effect), and a *vehicle*. Although inscriptions thus formed are still sometimes seen, particularly in purgative prescriptions, yet the tendency to simplicity in prescribing is increasing

daily, and prescriptions are often found containing only the basis and the vehicle.

Is a knowledge of Latin necessary in writing a prescription?

A very limited knowledge of Latin is sufficient for ordinary prescription writing, because, owing to the common practice of abbreviating, the last syllable of the Latin words, which varies with the case, is generally omitted.

Describe the method of writing a prescription.

Having written the superscription,* the physician places the names of the drugs which he wishes to administer, in a vertical column under the sign R., writing them in Latin and in the genitive case. He next determines how many doses he wishes to prescribe, and multiplying the single dose of each drug by the total number of doses in the prescription, he sets the amount opposite the corresponding medicine, expressing the quantity by the proper signs and numerals if it is written in apothecaries' weight and measure, or by Arabic figures and abbreviations if written in grammes. Thus, having determined to prescribe an iron mixture containing 24 doses, he might write—

R.	Tincturæ Ferri Chloridi, . . .	f℥ij	}	or	{	8		Gm.
	Acidi Acetici Diluti,	f℥j				4		"
	Liquoris Ammonii Acetatis, . . .	f℥j				32		"
	Elixir Aromatici,	f℥xiiij				52		"

Next comes the directions to the druggist, which in this case is simply to mix the medicines together and which is expressed by the Latin word *misce*, usually abbreviated to M.

Then follows Sig. or S. (signa), which means the druggist must write what follows on the label, as a direction for the patient, "Take one teaspoonful in water 3 times a day after eating," ending the prescription with the physician's name.

May the above prescription be written in any other way?

Instead of writing "Elixir Aromatici f℥xiiij," he may write "Elixir Aromatici *quantum sufficit* f℥ij," which means "as much of the aromatic elixir as will suffice (to make the whole mixture measure) f℥ij."

* The patient's name and address and the date are often omitted.

Why are the titles of the drugs written in the genitive case?

The verb "*recipe*" (R., "take thou") governs not the drug, but the quantity to be taken, which is placed consequently in the accusative case ("take thou two drachms"); the name of the drug *limits* the meaning of the quantity expressed, and according to rule is placed in the genitive case and governed by the amount ("take thou two drachms" of what?—"of the tincture of the chloride of iron").

Given a prescription containing a number of doses of different ingredients, how may a single dose of any one drug be ascertained?

Calculate the number of doses in the mixture and divide the quantity of any ingredient by the number, the result is a single dose.

Thus, the above prescription is a $\text{f}\overline{\text{3}}\text{ij}$ mixture, of which the dose is a teaspoonful or $\text{f}\overline{\text{3}}\text{j}$. As $\text{f}\overline{\text{3}}\text{viij} = \text{f}\overline{\text{3}}\text{j}$, there are 3 times $\text{f}\overline{\text{3}}\text{viij}$ or $\text{f}\overline{\text{3}}\text{xxiv}$ in the mixture, consequently 24 doses. To ascertain the amount of tincture of chloride of iron in each dose, convert the quantity of the tincture to minims and divide by 24: $\text{m}60 = \text{f}\overline{\text{3}}\text{j}$, therefore $\text{m}120 = \text{f}\overline{\text{3}}\text{ij}$; $\text{m}120 \div 24 = \text{m}v$, the quantity of the tincture in a single dose of the mixture.

How many doses should a mixture contain?

A physician should avoid prescribing more of a medicine than his patient will probably want. As a general rule, 24 doses is a sufficient amount, and forms a prescription in which the single doses of the ingredients are easily calculated.

Thus, if the dose be a teaspoonful ($\text{f}\overline{\text{3}}\text{j}$), 24 doses would make a $\text{f}\overline{\text{3}}\text{ij}$ mixture (see above); if, however, the dose is a dessertspoonful ($\overline{\text{3}}\text{ij}$), 24 doses would be twice as much, $\text{f}\overline{\text{3}}\text{vj}$. As $\text{m}iiss \times 24 = \text{m}60$ or $\text{f}\overline{\text{3}}\text{j}$, the latter amount represents 24 doses of $\text{m}iiss$ each, and consequently, if an entire mixture contains 24 doses, $\text{f}\overline{\text{3}}\text{j}$ of any of its components will contain 24 doses of $\text{m}iiss$.

As solids, when in a thorough state of solution, are equally diffused throughout the menstruum, if $\overline{\text{3}}\text{j}$ or gr. 60 of any solid is dissolved in 24 doses of a solution, each dose will contain gr. $iiss$ of the solid. Hence, in prescribing—

1st, write for a $\text{f}\overline{\text{3}}\text{ij}$ mixture if the dose is to be a teaspoonful ($\text{f}\overline{\text{3}}\text{j}$), or
 " $\text{f}\overline{\text{3}}\text{vj}$ " " " " " dessertspoonful ($\overline{\text{3}}\text{ij}$),

in either case the mixture will contain 24 doses.

2d, recollect that if each dose of such a mixture is to contain gr. iiss or ℥iiss of any substance, there should be ℥j or f℥j of the substance in the mixture;

if each dose should contain gr. v or ℥v, there should be ℥ij or f℥ij;
 " " " gr. x or ℥x, " " ℥ss or f℥ss;
 " " " gr. xx or ℥xx, " " ℥j or f℥j.

These rules simplify prescription-writing for beginners:

— Can any general rule be formulated for dosage of official preparations?

As a general rule the *fluid extracts* may be prescribed in ℥x doses. (*Exceptions* (the poisons), the fluid extracts of aconite (℥ss-ij), belladonna (℥j-ij), colchicum root (℥ij-v), colchicum seed (℥ij-x), digitalis (℥j-ij), nux vomica (℥j-v), sanguinaria (℥j-v), squill (℥j-ij), stramonium (℥j), veratrum viride (℥j-iv).) *Infusions* and *decoctions* in f℥ss-f℥ij doses. (*Exception*, infusion of digitalis, f℥j-iv.) *Syrups*, f℥j-f℥ij doses. (*Exceptions*, syrup of the iodide of iron, ℥v-xxx; compound syrup of squill, ℥x-xxx.) *Tinctures*, ℥v-f℥j doses. (*Exceptions* (the poisons), tinctures of aconite, ℥ij-ij; belladonna, ℥vij-xxx; colchicum, ℥v-xxx; digitalis, ℥v-x; iodine, ℥j-x; ipacac and opium, ℥v-xv; nux vomica, ℥j-x; opium, ℥ij-x; deodorized opium, ℥ij-x; physostigma, ℥v-xv; squill, ℥v-xxx; stramonium, ℥v-xv; veratrum viride, ℥ij-x.) *Powders*, gr. ij-x. *Volatile oils*, gtt. j-ij.

Many preparations can be administered in larger doses than indicated by this general rule.

Mention some circumstances which, by modifying the effects, will influence the dose of medicines.

As the activity of a drug is increased or lessened by the age, sex, idiosyncrasy, constitution and habits of the patient, as well as by the condition of the digestive and absorbent systems, disease, and manner of administration, these circumstances must be considered in relation to dosage.

How does age influence the activity of a drug?

Children and old persons are more readily influenced by most drugs, particularly narcotics, than adults in the prime of life; but,

as a rule, they bear larger *proportionate* doses of purgatives. No scheme for calculating the doses from the age of the patient alone can be relied on in practice.

What is Dr. Young's rule for calculating the dose for children?

Convert the child's age into a fraction by using the age itself as a numerator and the age + 12 as a denominator, the result being the proportion of the adult dose which should be administered to the child. Thus, at three years $\frac{3}{3+12} = \frac{3}{15} = \frac{1}{5}$ of that of the adult. At 21 years of age the full dose is given.

What is Dr. Cowling's rule?

Dr. Cowling proposed to divide the patient's *next birthday* by 24; thus, at three years the dose would be $\frac{4}{24} = \frac{1}{6}$ of that of the adult.

How may the dose for a child be more easily calculated in the metric system?

Dr. Lauder Brunton proposes to multiply the next birthday by 4 and divide the result by 100; or, what is the same thing, to multiply the full dose by the child's next birthday, then by 4, and remove the decimal point two places to the left. Thus, if the dose for an adult be 1. Gm., that for a child of 3 years will be $\frac{1 \times 4 \times 4}{100} = .160$ Gms.

How do sex, idiosyncrasy, constitution and habits influence medicinal effects?

Men generally bear larger doses of medicines than women. Many persons manifest idiosyncrasies with respect to certain drugs; thus, some are very susceptible to the poisonous effects of even minute doses of arsenic; others cannot take quinine or opium or belladonna or the iodides, and in others very dilute solutions of cocaine applied to the mucous membranes may cause alarming symptoms. As a rule, the larger and more robust the individual, the less easily is he influenced by drugs. When a medicine is habitually taken, the system becomes tolerant to it, and larger doses are required to produce any effect. This is especially the case with opium and purgatives.

In what way does the condition of the digestive and absorbent symptoms modify the effects of medicine?

During digestion, larger amounts of medicine are borne than can be taken fasting. In certain conditions of the alimentary canal, as in the third stages of yellow fever and of cholera, during a congestive chill of the gastro-enteric variety, or during profound opium narcosis, absorption from the stomach is slow, if it take place at all.

What effect does disease have upon the influence of medicines on the system?

In certain diseases great tolerance is manifested to medicine; thus, in peritonitis and cerebro-spinal fever enormous doses of opium are taken with advantage; in typhoid fever the tolerance to alcohol is marked.

Does the method of administration modify the effect of medicines?

It does: diuretics act more powerfully when much diluted; saline purgatives when concentrated. Usually medicines act more rapidly and powerfully when given by the mouth than when administered per rectum; still more so when injected under the skin, and most powerfully when injected into the blood.

How may medicines be administered?

Medicines may be administered by the *skin*, being either applied without friction (*euepidermic method*), or with friction (*epidermic method*), or else directly to the derma, which has been denuded by means of a blister (*endermic method*). They may also be given *hypodermically*, by means of a small syringe armed with a long, hollow needle. When thus administered, the dose should be from one-half to two-thirds of that which is required by the mouth, and the medicine should be in a thorough state of solution. Medicines are also applied to all the *mucous tracts* in the form of solutions, vapors (by atomization), powders (by insufflation), etc. The gastro-intestinal tract is most generally used when a systemic effect is desired, and the remedy is given either by the mouth or by the rectum. Suppositories and ointments are occasionally used per vaginam for their influence on the system. The broncho-pulmonary mucous membrane is used for the inhalation of vapors, as ether. *Intravenous injection* is rarely practiced except as a last resort.

How are medicines classified ?

Medicines are usually classified either according to their physiological action or to their therapeutic uses. The latter is the better arrangement.

The difficulty of grouping medicines is increased by the fact that even as yet the study of pharmacology is in its infancy, so that the exact physiological group to which many of them belong is not definitely determined. The action of most medicines, also, is complex, affecting different organs at the same time, and in many instances varying much according to the dose and the manner of administration.

They may be divided accordingly into—

GROUP 1.—SYSTEMIC MEDICINES.

Class 1.—Medicines acting on the body at large by their influence on digestion, nutrition (tissue waste and repair), and calorification.

Order I. Digestants. *Order II.* Tonics. *Order III.* Alteratives. *Order IV.* Antipyretics.

Class 2.—Medicines acting on the body by their effect on its various organs.

A. On the Nervous System:—

Order I. Hypnotics. *Order II.* Mydriatic Anodynes. *Order III.* Anæsthetics. *Order IV.* Antispasmodics. *Order V.* Excito-motors. *Order VI.* Depresso-motors.

B. On the Organs of Circulation:—

Order I. Stimulants. *Order II.* Sedatives.

C. On the Sexual Organs:—

Order I. Aphrodisiacs. *Order II.* Anaphrodisiacs. *Order III.* Oxytocics. *Order IV.* Uterine Sedatives. *Order V.* Emmenagogues.

D. On the Alimentary Canal:—

Order I. Emetics. *Order II.* Gastric Sedatives. *Order III.* Carminatives. *Order IV.* Cathartics. *Order V.* Anthelmintics.

Class 3.—Medicines Acting on the Various Secretions:—

Order I. Diuretics. *Order II.* Diaphoretics. *Order III.* Expectorants. *Order IV.* Astringents. *Order V.* Antacids.

GROUP 2.—TOPICAL MEDICINES.

Class 1. Antiseptics.

Class 2. Irritants.

Order I. Rubefacients. *Order II.* Vesicants. *Order III.*
Suppurants. *Order IV.* Escharotics.

Class 3. Demulcents.

Class 4. Emollients.

Class 5. Protectives.

Class 6. Coloring Agents.

GROUP I.—SYSTEMIC MEDICINES.

Class I.—Medicines Acting on the Body at Large by Their Influence on Digestion, Nutrition (Tissue Waste and Repair) or Calorification.

ORDER I.—DIGESTANTS.

What are digestants?

Digestants are animal or vegetable substances given to aid digestion when the normal alimentary secretions are insufficient for this function.

Name the digestants.

1. *Pepsinum* (*pepsin*), a proteolytic ferment obtained from the glandular layer of fresh stomachs from healthy pigs.
2. *Ingluvin*, a preparation from the gizzard of the domestic chicken.
3. *Pancreatinum* (*pancreatin*), a mixture of the enzymes naturally existing in the pancreas of warm-blooded animals, usually obtained from the fresh pancreas of the hog (*sus scrofa*).
4. *Papain*, a ferment obtained from the fruit of *Carica papaya* (*Nat. Ord.* Papayaceæ), a tree of South America.

What are the physiological actions of these substances?

1. *Pepsin* (when combined with an acid) digests albumen, which it converts into albuminose.
2. *Ingluvin* also aids the digestion of albumen, this action being probably due to a bitter principle which it contains and which stimulates the gastric glands.
3. *Pancreatin* (in neutral or alkaline solutions) converts albumen into albuminose and starches into glucose and aids in emulsifying fats.
4. *Papain* digests albumen and dissolves false membranes, ascariides and tænia.

What are the therapeutic uses of the digestants?

1. *Pepsin* is useful when the secretion of the peptic glands is not sufficient for the purposes of digestion, as in *atonic dyspepsia*, *apep-*

sia of infants, anæmia, chlorosis, gastric ulcer and cancer and infantile diarrhœa. It has been used for the relief of *reflex vomiting*, as that of *pregnancy*. It is an invaluable addition to nutritive enemata, and has been *injected into tumors* to arrest their growth and aid in their absorption.

2. *Ingluvin* is used chiefly for the relief of *vomiting of pregnancy*, also in *indigestion accompanied with flatulency*, and in *acute dyspepsia*.

3. *Pancreatin* is used in *intestinal indigestion*, and to aid in emulsifying fats. It is also added to nutritive enemata.

4. *Papain* is sometimes used in *dyspepsia*, to aid the digestion of albuminous foods, but more frequently for its solvent action on *false membranes and tænia*.

Pepsin, pancreatin (under the name of *trypsin*) and *papain* have been successfully used to *dissolve the membranes of croup and diphtheria*. They are used in solution, either in the form of spray or applied by a brush to the part.

What are the preparations and doses of the digestants ?

Pepsinum saccharatum (*saccharated pepsin*) should digest 300 times its own weight of freshly coagulated and disintegrated egg-albumen, dose gr. v-xxx (0.3-2.0).

Ingluvin (not official) may be given in doses of gr. v-xv (0.3-1.0).

Pancreatinum (*pancreatin*) in doses of gr. v-xv (0.3-1.0).

Papain (not official) in doses of gr. j-v (0.065-0.49).

Name some of the substances which are incompatible with the digestants.

Alcohol and many of the *mineral salts* destroy their properties.

Alkalies (when sufficiently powerful) interfere with the action of pepsin and ingluvin, while materially aiding that of pancreatin.

Acids retard and finally destroy the action of pancreatin while aiding those of pepsin and ingluvin.

How and at what time should the digestants be given ?

Pepsin and ingluvin should be given with meals, either in solution with an acid or in capsules, powder or compressed pill.

Often they are combined with sodium bicarbonate, to which bismuth is sometimes added, and administered before eating.

Pancreatin is usually given from 2 to 4 hours after meals, combined with an alkali.

Papain is taken after meals, in powder, capsule or compressed pill.

ORDER II.—TONICS.

What are tonics ?

Tonics are remedies which, in conditions of debility, impart a gradual and permanent increase of vigor to the body or its various parts, restoring the energy and strength which were wanting.

How may tonics be divided ?

They may be divided into *vegetable* and *mineral* tonics.

VEGETABLE TONICS.

What subdivisions of vegetable tonics are made ?

VEGETABLE TONICS are divisible into *simple or pure bitters*, *aromatic bitters* and *peculiar bitters*.

Simple Bitters.

Mention the simple bitters.

1. QUASSIA, the *wood* of *Picræna excelsa* (*Nat. Ord.* Simarubææ, a lofty tree of the West Indies.
2. GENTIANA (*gentian*), the *root* of the *Gentiana lutea* (*Nat. Ord.* Gentianeæ), a plant of the mountainous portions of Central and Southern Europe.
3. CALUMBA, the *root* of *Jateorrhiza palmata* (*Nat. Ord.* Menispermaceæ), a plant of the southeastern coast of Africa.
4. CHIRATA, *Swertia chirata* (*Nat. Ord.* Gentianeæ), a *plant* of India.

What are their chief chemical constituents ?

1. *Quassia-wood* contains *quassin*, a neutral bitter principle ;
2. *Gentian*, an acid, *gentisic acid* (gentisin) ; a fixed oil, and a bitter principle, *gentianin* or *gentiopicroin* ;
3. *Calumba*, an alkaloid, *berberine* (which is found in many other plants, and has been used as a tonic and febrifuge) ; a bitter principle, *columbin*, and an acid, *columbic acid* ; and

4. *Chirata*, a neutral bitter principle, *chiratin*; and an amorphous acid, *ophelic*.

What are the physiological actions of the simple bitters?

The simple bitters increase the secretions, and impart tone to the mucous membranes; increase the appetite and digestion, and thus promote nutrition and improve the condition of the blood, but when taken continuously for too long a time they disorder digestion and cause gastric catarrh. Gentian, in full doses, by increasing the intestinal secretion, acts as a mild laxative.

Calumba is least likely to disorder the stomach.

What are the therapeutic uses of the simple bitters?

They are given (1) to *increase secretion and promote appetite in atonic dyspepsia*, in *chronic gastric catarrh* and during *convalescence* from acute diseases. Calumba is the mildest, and, being best adapted to weak stomachs, may be used in *vomiting of pregnancy* and in *sea-sickness*; quassia, gentian and chirata are well adapted to cases of dyspepsia with relaxation or torpor of the mucous membrane and resulting constipation. (2) *To give tone to the mucous membranes*, in *diarrhoea* and in *accumulations of flatus* in the bowel from a relaxed condition, calumba may be used. (3) *As vehicles* for the administration of other medicines, as acids or alkalies. (4) *To destroy ascarides*, an infusion of quassia may be used as an enema.

What are the preparations and doses of these substances?

1. EXTRACTUM QUASSIÆ (*extract of quassia*), dose gr. j-v (0.065-0.5), principally used as an excipient.

EXTRACTUM QUASSIÆ FLUIDUM (*fluid extract of quassia*), dose $\text{m}_{\text{V-XXX}}$ (0.308-1.848).

TINCTURA QUASSIÆ (*tincture of quassia*), f 3j-ij (4.0-8.0).

Quassin (not official), dose gr. $\frac{1}{4}$ - $\frac{1}{2}$ (0.016-0.032).

2. EXTRACTUM GENTIANÆ (*extract of gentian*), dose gr. j-xxx (0.065-2.0).

EXTRACTUM GENTIANÆ FLUIDUM (*fluid extract of gentian*), dose f 3ss-j (2.0-4.0).

TINCTURA GENTIANÆ COMPOSITA (*compound tincture of gentian*), dose f 3ss-ij (2.0-8.0).

3. EXTRACTUM CALUMBÆ FLUIDUM (*fluid extract of calumba*), dose f 3ss-j (2.0-4.0).

TINCTURA CALUMBÆ (*tincture of calumba*), dose fʒss-ij (2.0-8.0).

Berberine (not official), dose gr. j-x (0.065-0.648) as a sulphate or hydrochlorate.

4. EXTRACTUM CHIRATÆ FLUIDUM (*fluid extract of chirata*), dose ʒx-xx (0.616-1.3).

TINCTURA CHIRATÆ (*tincture of chirata*), dose ʒv-fʒj (0.308-4.0).

What medicines should not be prescribed with gentian?

Iron sulphate and the salts of silver and of lead are incompatible with gentian.

How and at what time should the simple bitters be given?

The simple bitters are given before eating, either alone or combined with other medicines in pills or in mixtures or solutions.

Aromatic Bitters.

How do the aromatic differ from the simple bitters?

The aromatic bitters are more *stimulating* to the mucous tract than the simple bitters, because of the aromatic volatile oils which they contain; they also contain tannic or gallic acid, which renders them more or less *astringent*.

Mention the aromatic bitters.

1. SERPENTARIA (*Virginia snakeroot*), the *rhizome* and *rootlets* of *Aristolochia serpentaria* (*Nat. Ord.* Aristolochiaceæ) and other varieties of *Aristolochia*, indigenous plants;

2. EUCALYPTUS, the *leaves* of *E. globulus* or blue gum-tree (*Nat. Ord.* Myrtaceæ), a tree of Australia;

3. PRUNUS VIRGINIANA, the *bark* of *P. serotina* or wild cherry (*Nat. Ord.* Rosaceæ), an indigenous tree;

4. ANTHEMIS, the *flower-heads* of *A. nobilis* or chamomile (*Nat. Ord.* Compositæ), a native of Europe;

5. MATRICARIA, the *flower-heads* of *M. Chamomilla* or German chamomile (*Nat. Ord.* Compositæ);

6. EUPATORIUM, the *leaves* and *flowering tops* of *E. perfoliatum*, boneset or thoroughwort (*Nat. Ord.* Compositæ), a common indigenous plant;

7. CASCARILLA, the *bark* of *Croton Eluteria* (*Nat. Ord.* Euphorbiaceæ), a small tree of the West India Islands; and

8. **ABSINTHIUM**, the *leaves* and *tops* of *Artemisia Absinthium* or wormwood (*Nat. Ord.* Compositæ), a European plant.

What are the chief chemical constituents of the aromatic bitters?

They all contain a *volatile oil*, a *bitter principle*, *resin* and *tannic acid*.

The *volatile oil of eucalyptus* is composed of two terpenes, *cymol* and *eucalyptol*; that of *Prunus Virginiana* contains *hydrocyanic acid*, which is formed by the reaction in water of a peculiar ferment (not emulsin) existing in the bark, with the bitter principle; that of *absinthium* contains *absinthol*.

The *bitter principle of serpentaria* is called *aristolochin*; that of *anthesis*, *anthesis acid*; that of *eupatorium*, *eupaturin*; that of *cascarilla*, *cascarillin*; and that of *absinthium*, *absinthin*.

What are the physiological actions of the aromatic bitters?

Owing to the bitter principles which they contain, their action on digestion, nutrition and secretion is similar to that produced by the simple bitters. They have, in addition, a stimulating influence on the mucous membranes, due to the volatile oils (which are eliminated by the bronchial mucous membrane, skin and kidneys, and increasing their secretions are, somewhat, expectorants, diaphoretics and diuretics), and an astringent effect, owing to their tannic acid. In large doses they are emetic and laxative. *The oil of eucalyptus* destroys germs, and is an irritant when applied locally. It quickens the pulse and respiration and lowers the blood pressure. In large doses it causes diarrhoea. *Prunus Virginiana* has a sedative influence over the pneumogastric nerve, from the hydrocyanic acid which is developed in it, and thus quiets cough. *Absinthium* in large doses, or when habitually used in the form of the *liqueur (absinthe)*, produces epileptiform convulsions.

What are the therapeutic uses of the aromatic bitters?

They are used for the same purposes as the simple bitters.

Serpentaria is used as a *cardiac stimulant* and *tonic* in low fevers, and occasionally as a *stimulating expectorant* in capillary bronchitis and low pneumonias. *Eucalyptus* is used in *chronic catarrhal conditions* of the gastric, broncho-pulmonary, and genito-urinary, mucous

membranes, and from its *destructive influence on germs* it is of benefit in vomiting due to *sarcina ventriculi*. As an *antimalarial* agent it is far inferior to cinchona, but sometimes proves useful where the latter has failed. As a *local* application, it is employed in stomatitis. *Prunus Virginiana* is also used as a *stomachic tonic*, but is especially employed as a *cough-relieving agent* in bronchial catarrh and in phthisis.

What are the preparations and doses of these substances?

1. EXTRACTUM SERPENTARIÆ FLUIDUM (*fluid extract of serpentaria*), dose f3ss-j (2.0-4.0).

TINCTURA SERPENTARIÆ (*tincture of serpentaria*), dose f3j-ij (3.0-7.0).

2. EXTRACTUM EUCALYPTI FLUIDUM (*fluid extract of eucalyptus*), dose f3ss-j (2.0-4.0).

OLEUM EUCALYPTI (*oil of eucalyptus*), dose m_v-xv (0.3-1.0) (in capsules or in emulsion).

EUCALYPTOL, a neutral body obtained from the volatile oil of *Eucalyptus globulus* and of some other species of *Eucalyptus*, dose m_v-xv (0.3-1.0).

3. EXTRACTUM PRUNI VIRGINIANÆ FLUIDUM (*fluid extract of prunus virginiana*), dose f3j-ij (3.0-7.0).

INFUSUM PRUNI VIRGINIANÆ (*infusion of prunus virginiana*), dose f3ss-ij (15.0-60.0).

SYRUPUS PRUNI VIRGINIANÆ (*syrup of prunus virginiana*), dose f3j-f3ss (3.5-15.0).

4 and 5. There are no official preparations of *anthemis* and *matriaria*, the latter of which is not much used in this country. They are usually given in infusion (chamomile tea) in doses of f3ij (60.0) t. i. d., as a stomachic tonic.

6. EXTRACTUM EUPATORII FLUIDUM (*fluid extract of eupatorium*), dose f3ss-j (2.0-4.0).

7 and 8. *Cascarilla* and *absinthium* may be given in infusion. There are no official preparations.

At what time should the aromatic bitters be administered?

They should be given shortly before meals when intended to promote appetite and digestion, but when given for their effects on the bronchial or vesical mucous membranes, or for their action on the heart, they should be given between meals.

*Peculiar Bitters.***What are the peculiar bitters?**

They are such bitters as, owing to peculiarity of action, cannot be classed with either of the preceding groups.

Name the peculiar bitters.

Cinchona and hydrastis.

CINCHONA.

What is cinchona?

Cinchona is the *barks* of all species of *Cinchona* (*Nat. Ord. Rubiaceæ*) which contain not less than 5 per cent. of total alkaloids, and at least 2.5 per cent. of quinine.

What is the habitat and what are the principal varieties of cinchona?

The cinchona trees are natives of the mountainous regions of the western parts of S. America, from 22° south latitude to 10° north latitude, but have been successfully cultivated in India, Ceylon, Java, the West Indies and other parts of the world. The two principal varieties are *cinchona calisaya* from which is derived the *cinchona flava* (yellow or calisaya bark) and the *cinchona succirubra*, whence comes the *cinchona rubra* (red bark). Other varieties are *C. micrantha* and *C. condaminea*, from which is obtained *cinchona pallida* or pale bark, no longer official.

How is the bark obtained?

The bark was formerly obtained by stripping the trees, but as this killed them and threatened to extinguish the genus, the present method has been adopted. One-fifth the circumference of the tree is denuded of bark in longitudinal strips each year, in the dry season, for five years, at the end of which time the bark has again grown over the portion first denuded. The process is then repeated.

What are the chemical constituents of cinchona?

The most important constituents are the alkaloids *quinine* and *cinchonine*, of which the former is found to preponderate in the yellow bark, the latter in pale barks, and both exist in about equal proportions in red bark. Other important alkaloids are *quinidine* and *cinchonidine*. Combined with these alkaloids are *kinic*, *kinovic*, *cincho-tannic* and *kinovo-tannic acids*. Other less important alkaloids are

also found, as *aricine*, *paricine*, *quinamine* and *paytine*. *Quinicine* and *cinchonine* are amorphous modifications of quinine and cinchonine.

Describe a chemical test for the cinchona alkaloids.

If *quinine* and *quinidine*, or *their salts*, are treated with fresh chlorine-water, an emerald-green precipitate will form when ammonia-water is added. When the same test is applied to *cinchonine*, *cinchonidine*, or *their salts*, a white precipitate results.

What are the physiological effects of cinchona and its alkaloids?

Locally cinchona and its alkaloids are irritant and antiseptic, destroying minute organisms or inhibiting their movements.

Internally it acts on the *alimentary canal* as a simple bitter, in small doses increasing the appetite and digestion; if long continued, producing indigestion and gastric catarrh. In ordinary doses cinchona constipates. In large doses cinchona or its alkaloids irritate the mucous membrane and cause vomiting and diarrhoea.

Nutrition is stimulated and the excretion of waste products increased by small doses; large doses, however, diminish the amount of urea and uric and phosphoric acids in the urine. In malarial fever the products of waste tissue are much increased.

In the *blood* it interferes with the oxygen-carrying function of the red corpuscles, and diminishes the number, inhibits the movements, and prevents the migration of the white corpuscles, both in health and disease.

Circulation: in small doses the cardiac action is increased; large doses, by acting on the cardiac motor ganglia, depress the heart, sometimes causing it to intermit, and finally arrest it in diastole; the blood pressure is lowered.

The *temperature* in health is very slightly, if at all, influenced; in fevers a rapid decline takes place, due to the action on the blood and circulation.

Nervous system; small doses stimulate the cerebral functions; large doses cause *cinchonism* (*i. e.*, a constricted feeling in the forehead; giddiness and tinnitus aurium, with impairment of hearing and sometimes of vision); after toxic doses these symptoms are aggravated and delirium, weak pulse, coma (sometimes convulsions), and in rare

cases death, supervene. It probably reduces the reflex excitability of the spinal cord, and although it never causes, may energize uterine contractions when present. *Elimination* takes place chiefly by the urine, the greater portion escaping from the system within eight hours.

Cutaneous eruptions, as erythema, urticaria or herpes are produced in some subjects by even small doses of cinchona or its alkaloids.

What are the therapeutic uses of cinchona and its alkaloids?

1. *As a tonic.* The cinchona preparations, as well as the salts of its alkaloids, are largely used to improve the appetite and digestion in *debility*, the *latter stages of inflammations* and in *convalescence* from acute diseases generally, also in *atonic dyspepsia* (combined with a mineral acid), in *relaxed conditions* of the alimentary mucous membrane and in *summer complaint* of children. The alkaloids are also used as tonics to sustain the vital forces in *diphtheria*, *erysipelas*, *pycemia*, *septicæmia* in all *typhoid conditions*, in the *eruptive fevers*, and in *surgical shock*. In the various *neuralgiæ* combined with other remedies (as morphine or belladonna, iron or arsenic), it is a valuable agent. From its influence on the migration of leucocytes and on cell proliferation it is used (in larger doses) in the *early stages of inflammation*.

2. *As an antipyretic* in all conditions of hyperpyrexia quinine may be used in large doses, repeated in about six or eight hours to maintain its effects, but the newer and more powerful antipyretics have largely superseded this practice.

3. *As an antiperiodic*, cinchona has no rival in the materia medica. The salts of the alkaloids are preferred for this purpose. Quinine may be used as a *prophylactic* against malaria in small doses, given early in the morning (gr. iij-x, 0.2-0.65). In the treatment of *intermittent* (gr. x (0.65) during the sweating stage, repeated five hours before the next paroxysm is expected), *remittent* (gr. xxx (2.0) per day in two doses, given at any time), *hemorrhagic* and *pernicious malarial fevers* (gr. xx-3j (1.3-4.0) administered promptly by the mouth or rectum or even subcutaneously), it acts as a specific. It is useful also in malarial cachexia, but must often be combined or alternated with arsenic and other treatment.

4. *Locally*, quinine may be applied in solution as an astringent and antiseptic to the nares in *summer catarrh*, etc., or to the mouth in *muguet*.

What are the preparations and doses of cinchona and its alkaloids?

EXTRACTUM CINCHONÆ (*extract of (yellow bark of) cinchona*), dose as a *tonic* gr. x-xxx (0.65-2.0).

EXTRACTUM CINCHONÆ FLUIDUM (*fluid extract of (yellow bark of) cinchona*), dose as a *tonic* fʒj (4.0).

TINCTURA CINCHONÆ (*tincture of (yellow bark of) cinchona*), dose fʒj-iv (4.0-15.0).

TINCTURA CINCHONÆ COMPOSITA (*compound tincture (Huxham's) of cinchona*), containing red bark, serpentaria, bitter orange peel, glycerin, alcohol, and water, dose fʒj-iv (4.0-15.0).

INFUSUM CINCHONÆ (*infusion of cinchona*), containing aromatic sulphuric acid, dose fʒss-ij (15.0-60.0).

QUININÆ SULPHAS (*quinine sulphate*), the basic sulphate, not very soluble in water, but rendered more soluble by the addition of diluted sulphuric acid, dose as a *tonic*, gr. j-vj (0.065-0.4); as an *antipyretic*, gr. x-xx (0.65-1.3), repeated as indicated; as an *anti-periodic*, gr. xvj-xx (1.0-1.3) or more.

QUININÆ BISULPHAS (*quinine bisulphate*), the normal sulphate, more soluble, but containing less quinine than the basic sulphate, and given in slightly larger doses.

QUININÆ HYDROBROMAS (*quinine hydrobromate*), better adapted for hypodermic injection than the sulphates; dose, by the mouth, the same as of quinine sulphate, a little less when used hypodermically.

QUININÆ HYDROCHLORAS (*quinine hydrochlorate*), also adapted for subcutaneous use; dose, as of the preceding preparations.

QUININÆ VALERIANAS (*quinine valerianate*), dose gr. j-vj (0.065-0.389).

QUINIDINÆ SULPHAS (*quinidine sulphate*), dose slightly larger than those of quinine.

CINCHONINÆ SULPHAS (*cinchonine sulphate*), dose $\frac{1}{2}$ larger than those of quinine.

CINCHONIDINÆ SULPHAS (*cinchonidine sulphate*), dose gr. j-xv (0.065-1.0) according to the effects desired.

What medicines are incompatible with cinchona?

The *alkalies* and *alkaline earths* precipitate the alkaloids of cinchona; *tannic, gallic* and *tartaric acid*, the *tincture* and the *compound solution of iodine* form insoluble compounds with them; and the *ferric salts* precipitate cincho-tannic acid from the preparations of cinchona.

How are cinchona and its alkaloids administered?

The preparations of *cinchona* are frequently combined with the mineral acids or administered alone as tonics; the alkaloids are given in solution, freshly made pills or capsules, or may be made into suppositories, or used subcutaneously. They are often combined with iron, arsenic and other medicines. Saccharin is said to disguise their unpleasant taste.

HYDRASTIS.

What is hydrastis?

HYDRASTIS is the *rhizome* and *roots* of *H. canadensis*, yellow root or golden seal (*Nat. Ord. Ranunculacæ*), an indigenous plant.

What alkaloids does it contain?

It contains *hydrastine* (not hydrastin, which is an eclectic preparation) and *berberine*.

What are its physiological actions?

It acts on the appetite, digestion, nutrition and secretions like the simple bitters; it increases the secretions, especially of the glandular appendages of the intestine and probably that of the liver; when given to animals in sufficient doses, it is said to act on the nervous system like strychnine, but less powerfully, stimulating the reflex centres of the spinal cord, heightening the irritability of the motor nerves and causing incoördination, tremor, convulsions, and, finally, death from tetanic spasm of the respiratory muscles. It is also claimed that it always provokes uterine contractions.

What are the therapeutical applications of hydrastis?

It is used as a tonic, to promote appetite, digestion, and nutrition in *atonic dyspepsia* and *chronic gastric catarrh*. From its effects on secretions, it is used in *catarrhal jaundice*, and in *constipation* from deficient secretion, and as a *diuretic* and *blennorrhætic* in *chronic interstitial nephritis* and *chronic cystitis*. It has also been recom-

mended internally to check *uterine hemorrhage*. As an anti-periodic in *intermittent fever* and *chronic malaria* it is inferior to quinine.

Locally, it has been used as an application to sore mouth, fissured nipples, sores and ulcers generally, and as an injection in gleet and vaginal leucorrhœa.

What are the preparations and doses of hydrastis?

EXTRACTUM HYDRASTIS FLUIDUM (*fluid extract of hydrastis*), dose $\mathfrak{m}_{\text{v-f3j}}$ (0.3-4.0); as a stomachic tonic, $\mathfrak{m}_{\text{v-xv}}$ (0.3-1.0) before meals.

TINCTURA HYDRASTIS (*tincture of hydrastis*), dose $\mathfrak{m}_{\text{x-f3j}}$ (0.6-4.0).

GLYCERITUM HYDRASTIS (*glycerite of hydrastis*) is chiefly used as a local application.

HYDRASTININÆ HYDROCHLORAS (*hydrastinine hydrochlorate*) has been used to check *uterine hemorrhage*, and also as a *heart tonic* (*vid. oxytocics*); dose gr. $\frac{1}{4}$ -1 (0.016-0.06).

MINERAL TONICS.

Name the mineral tonics.

The MINERAL TONICS are the *preparations of iron* and of *manganese* (which are sometimes classed separately as *hæmatinics* or blood medicines), *phosphorus*, *zinc phosphide*, the *mineral acids*, and *lactic acid*, although of animal origin, is usually considered in connection with the latter.

FERRUM—IRON.

What is the source of iron?

Iron is the most common of all the metals, and is found widely distributed both in the metallic state and as oxide, sulphide, carbonate, phosphate, etc.

What are the physiological effects of iron?

Metallic iron is inert, but when swallowed it is dissolved by the acid of the stomach, and forms salts. It probably enters the blood as an albuminate. Any iron which is not absorbed passing into the intestines, is converted into a sulphide, and colors the stools black. Iron exists normally in the red corpuscles of the blood in the pro-

portion of 1 part of iron to 230 parts of corpuscles. When deficient in the blood, under a course of iron the number of corpuscles increase, the blood is improved in quality, and consequently the tissues are better nourished and all the functions of the body are better performed. In appropriate cases it promotes the appetite and digestion by stimulating the gastric glands, but if given for too long a time it disorders digestion.

The *more soluble preparations* of iron, if given in large doses, cause nausea and vomiting; its *salts, with the mineral acids and with iodine*, possess more or less toxic properties.

The *sulphates, nitrates and chlorides* of iron are astringent, diminishing the secretions of the mucous membranes, thus causing constipation, and coagulating albumen, thus proving hæmostatic. Iron is eliminated by the fæces, bile and urine.

What are the therapeutic uses of iron?

1. *As a general tonic* to improve the appetite and digestion, when it should be combined with a purgative, as aloes.

2. *To improve the condition of the blood* and aid nutrition, in *anæmia* and *chlorosis*; in *malarial anæmia* (with quinine); in *various cachexiæ*, as the *scrofulous, strumous or syphilitic (iodides)*; in *amenorrhœa, dysmenorrhœa* or *menorrhagia* associated with *anæmia*; in *fatty degeneration* and in *dilatation of the heart*; in many forms of *neuralgia*, and in many *nervous affections* associated with *anæmia*, as *epilepsy* or *chorea*; in *convalescence* from acute diseases, and after *exhausting hemorrhages*. In *erysipelas* iron is almost a specific (tincture of the chloride $\text{m}_{\text{xx-xxx}}$ (1.2-2.0) every 2 or 3 hours), and in diphtheria it is also useful.

3. *As an astringent* it is sometimes used internally in *chronic diarrhœa* and in dysentery. In *chronic albuminuria* it is useful both as a tonic and astringent.

4. *As a hæmostatic*, in all *passive hemorrhages* in the *anæmic* (tincture of the chloride), and in hemorrhages from the lungs, stomach, bowels or kidneys (subsulphate, tersulphate or nitrate in drop doses largely diluted and frequently repeated).

5. *Locally* it is employed as a *styptic* (subsulphate), to check the *bleeding from hemorrhoids* and *leech bites*, or to arrest *capillary oozing* from a large surface; in the form of spray in *epistaxis*, and

hæmoptysis, and it is sometimes injected into the cavity of the uterus (more or less dangerous) to check bleeding from that organ. As an *astringent*, it is applied to syphilitic vegetations (Monsel's solution) and as an injection (largely diluted) in gonorrhœa and leucorrhœa.

What are the preparations of iron and their doses?

FERRUM REDUCTUM (*reduced iron—Quevenne's iron*), an impalpable powder of metallic iron, well borne by the stomach, and not astringent; dose gr. j-v (0.065-0.324).

FERRI OXIDUM HYDRATUM (*hydrated ferric oxide*) and FERRI OXIDUM HYDRATUM CUM MAGNESIA (*hydrated ferric oxide with magnesia*), used almost exclusively as antidotes in arsenical poisoning; dose fʒss (16.0), every 5 or 10 minutes.

TROCHISCI FERRI (*troches of iron*), each lozenge contains of ferric hydrate nearly gr. v (0.30).

EMPLASTRUM FERRI (*plaster of iron*) contains ferric hydrate, lead plaster, Burgundy pitch, and olive oil.

FERRI CARBONAS SACCHARATUS (*saccharated ferrous carbonate*), dose gr. ij-x (0.130-0.650).

MASSA FERRI CARBONATIS (*mass of ferrous carbonate—Vallet's mass*), dose gr. iij-v (0.20-0.30).

PILULÆ FERRI CARBONATIS (*pills of iron carbonate—Blaud's pills*). Each pill is intended to contain about gr. j (0.0648) of ferrous carbonate, an excellent preparation in anæmia and chlorosis; dose 2-5 pills.

MISTURA FERRI COMPOSITÆ (*compound iron mixture—Griffith's anti-hectic mixture*), dose, in anæmia or chlorosis, fʒss-ij (15.0-60.0).

FERRI SULPHAS (*ferrous sulphate—green vitriol or copperas*). One of the most active and astringent preparations of iron. It is irritant to the stomach, and in overdoses a poison. It is used locally as a styptic and astringent, and in solution to disinfect privies and drains; dose (in pill) gr. ss-ij (0.03-0.13).

FERRI SULPHAS EXSICCATUS (*dried ferrous sulphate*), ferrous sulphate deprived of its water of crystallization; dose gr. ¼-2 (0.016-0.13).

FERRI SULPHAS GRANULATUS (*granulated ferrous sulphate*), dose gr. j-v (0.065-0.30).

LIQUOR FERRI TERSULPHATIS (*solution of ferric sulphate*), a solution of normal ferric sulphate; rarely used except in making hydrated ferric oxide.

LIQUOR FERRI SUBSULPHATIS (*solution of ferric subsulphate—Monsel's solution*), a solution of basic ferric sulphate. It may be given internally, largely diluted, as a hæmostatic; dose $\mathfrak{m}\text{j-x}$ (0.06–0.6). It is chiefly used locally as a styptic.

FERRI CHLORIDUM (*ferric chloride*), rarely used internally; dose gr. j–ij (0.065–0.13).

LIQUOR FERRI CHLORIDI (*solution of ferric chloride*), dose $\mathfrak{m}\text{ij-v}$ (0.123–0.30), largely diluted; externally it is used as a styptic.

TINCTURA FERRI CHLORIDI (*tincture of ferric chloride*), one of the best preparations of iron for internal use. It possesses diuretic properties; dose $\mathfrak{m}\text{v-xxx}$ (0.30–1.85).

LIQUOR FERRI ET AMMONII ACETATIS (*solution of iron and ammonium acetate—Basham's mixture*), an excellent preparation, combining tonic and diuretic properties, very valuable in chronic desquamative nephritis and chronic dropsies generally; dose $\mathfrak{f}\overline{3}\text{ss-j}$ (15.0–30.0).

FERRI IODIDUM SACCHARATUM (*saccharated ferrous iodide*), dose gr. iij–x (0.20–0.65).

SYRUPUS FERRI IODIDI (*syrup of ferrous iodide*), dose $\mathfrak{m}\text{v-xxx}$ (0.30–1.85).

PILULÆ FERRI IODIDI (*pills of ferrous iodide*), dose 1 to 3 pills.

FERRI PHOSPHAS SOLUBILIS (*soluble ferric phosphate*), dose gr. ij–v (0.13–0.32).

FERRI PYROPHOSPHAS SOLUBILIS (*soluble ferric pyrophosphate*), dose gr. ij–v (0.13–0.32).

FERRI HYPOPHOSPHIS (*ferric hypophosphite*), dose gr. v–x (0.32–0.65).

LIQUOR FERRI NITRATIS (*solution of ferric nitrate*), used internally as an astringent and hæmostatic; dose $\mathfrak{m}\text{ij-v}$ (0.123–0.30), freely diluted.

FERRI CITRAS (*ferric citrate*), dose gr. ij–x (0.13–0.65).

LIQUOR FERRI CITRATIS (*solution of ferric citrate*), dose $\mathfrak{m}\text{ij-x}$ (0.183–0.61).

LIQUOR FERRI ACETATIS (*solution of ferric acetate*), rarely used internally; dose $\mathfrak{m}\text{x-xxx}$ (0.61–1.85).

FERRI LACTAS (*ferrous lactate*), a mild chalybeate, agreeing well with the stomach, and useful in anæmia and chlorosis, with weak digestion; dose gr. ij-x (0.13-0.65).

FERRI VALERIANAS (*ferric valerianate*), dose gr. j-ij (0.065-0.20).

FERRI ET POTASSII TARTRAS (*iron and potassium tartrate—potassio-tartrate of iron*) may be prescribed with alkalies; dose gr. ij-x (0.13-0.65).

FERRI ET AMMONII TARTRAS (*iron and ammonium tartrate*), a mild preparation; dose gr. ij-x (0.13-0.65).

FERRI ET AMMONII SULPHAS (*ferric ammonium sulphate—ammonio-ferric alum*) is given internally as an astringent in chronic diarrhœa and dysentery; dose gr. j-v (0.065-0.32).

FERRI ET AMMONII CITRAS (*iron and ammonium citrate*), dose gr. ij-v (0.13-0.32).

VINUM FERRI CITRATIS (*wine of iron citrate*), dose fʒj (4.0).

FERRI ET QUININÆ CITRAS (*iron and quinine citrate*),

FERRI ET QUININÆ CITRAS SOLUBILIS (*soluble iron and quinine citrate*), each contains 85 per cent. of ferric citrate, and 12 per cent. of the alkaloid, and may be given in doses of gr. j-v or x (0.065-0.32 or 0.65).

VINUM FERRI AMARUM (*bitter wine of iron*), dose fʒj-ij (4.0-8.0).

FERRI ET STRYCHNINÆ CITRAS (*iron and strychnine citrate*) contains 98 per cent. of iron and ammonium citrate and 1 per cent. of the alkaloid; dose gr. ij-v (0.13-0.32).

SYRUPUS FERRI QUININÆ ET STRYCHNINÆ PHOSPHATUM (*syrup of iron, quinine, and strychnine phosphates*), dose fʒj (4.0).

What medicines are incompatible with the iron preparations?

I. *The alkaline preparations of iron* are incompatible with the acids and acidulous salts.

II. *The salts formed by mineral acids with iron*, with the alkalies and alkaline earths and their carbonates.

III. *The iodides, bromides and salts containing vegetable acids*, with the alkalies and alkaline earths and their carbonates, and with acids and (other) acidulous salts.

IV. *All the iron preparations*, with substances containing tannic or

gallic acids, hence they cannot be prescribed with any of the vegetable tonics except the simple bitters.

When and how should iron be administered?

The fluid preparations should all be given freely diluted. The solid preparations may be given combined with other remedies in pilular form or in solution. Iron causes less gastric irritation and enters the blood more readily if given with or immediately after meals. If gastric irritability is present in anæmia, a course of bitters with the acids had better precede the administration of iron, which, as these symptoms subside, may be given in the form of the lactate, followed by the bitter wine, and finally by the dried sulphate with arsenic in pill, or by the tincture of the chloride. In anæmia iron should be given in rapidly increasing doses until headache, flushing of the face, and some nausea supervene, when the dose should be diminished or the medicine withheld for a few days. It is important to overcome the resulting constipation by occasional laxatives.

MANGANUM—MANGANESE.

What are the sources of manganese?

Manganese is obtained from the native (impure) dioxide, and also from the carbonate.

What are its physiological actions?

Manganese exists normally in the blood, hair, bile, etc., associated with iron; the proportion to iron in the red corpuscles being as 1 to 20. Given internally in small doses, it improves the appetite, digestion and nutrition; in larger doses, it depresses the heart's action and lowers the blood pressure; if large doses are given for some time, the effects resemble those of zinc (wasting and feebleness, staggering gait, and finally paraplegia). In toxic doses it is a gastro-intestinal irritant. The *sulphate* is an emeto-cathartic and cholagogue. *Potassium permanganate* is a powerful oxidizing agent, and hence an antiseptic. It increases the menstrual flow.

For what medicinal purposes is manganese used?

It has been used (1) *to improve the condition of the blood in anæmia and chlorosis and in various cachexia*, either combined with iron

or as a substitute for it; (2) *to improve the appetite and digestion in gastrodynia and pyrosis*; (3) *as a cholagogue in jaundice* when of malarial origin (sulphate); (4) *as an antidote to snake bites* and other animal poisons, and in poisoning by morphine (potassium permanganate internally and locally); (5) in *amenorrhœa* (potassium permanganate); and (6) *locally as a disinfectant and to correct fetor in scarlatina and diphtheria* (permanganate as a gargle), or applied to *cancerous or other ulcers, abscesses, otorrhœa, etc.*

What are the preparations and doses of manganese?

MANGANI DIOXIDUM (*manganese dioxide*), dose gr. j-x (0.065-0.65), in pill, powder, or capsule.

MANGANI SULPHAS (*manganese sulphate*), dose gr. j-v (0.065-0.32).

POTASSII PERMANGANAS (*potassium permanganate*), dose gr. ss-ij (0.032-0.13) in distilled water, in capsule, or made into pill with kaolin.

What medicines are incompatible with manganese?

The salts of lead, silver, and mercury, and the caustic alkalies are chemically incompatible with the manganese preparations.

PHOSPHORUS.

What is the source from which phosphorus is obtained?

Phosphorus is obtained from the calcium phosphate of bones.

What are the physiological effects of phosphorus?

Phosphorus exists normally in nervous tissue and in the bones. *Locally* applied, phosphorus causes inflammation, ulceration, and even gangrene of the skin. *The vapor*, when inhaled, irritates the mucous membrane, and if caries of the teeth exist, may cause caries or necrosis of the maxillæ.

In medicinal doses, phosphorus increases the *appetite and digestion* (if too long continued, it disorders digestion), and causes eructations of hydrogen phosphide; increases *nutrition*, the *cardiac action* and cutaneous circulation, *mental activity*, *sexual activity*, the formation of the *compact tissue of bone*, and the excretion of phosphates by the *urine*. It is *eliminated* by the liver, kidneys, mucous membrane and skin.

What are the toxic effects of phosphorus?

From 1 to 3 hours after a poisonous dose has been swallowed, violent *epigastric pains* occur, followed by nausea and *vomiting* (the vomited matter *smelling of phosphorus* and being *luminous in the dark*); *vomiting of blood* occurs at a later period; violent purging takes place if the phosphorus passes into the bowel. *Jaundice* may supervene after several days if the patient lives. The pulse is at first strong, soon becoming feeble and rapid. There is great thirst, restlessness, anxiety, noisy delirium or coma and death, preceded by convulsions. The *urine* is diminished, contains albumen, blood, and (after jaundice occurs), biliary pigment. After death, acute fatty degeneration of various viscera, as the liver, kidneys, heart and the walls of the arterioles and capillaries is found: the red corpuscles are altered in form and the blood contains much fat.

What is the treatment of acute poisoning by phosphorus?

1. Give *cupric sulphate* gr. v (0.32) at once as an emetic and as an antidote (forming a less soluble phosphide).
2. Give *hydrated magnesia* as a purgative.
3. Continue the use of *cupric sulphate* in small doses (gr. $\frac{1}{12}$ (0.005) every fifteen or twenty minutes) as an antidote or emetic, or, better, give *turpentine* (French acid turpentine or old common, crude turpentine, which forms with phosphorus an almost insoluble spermaceti-like mass) as soon as possible, to neutralize the poison.
4. Give *opium* to counteract the general depression.
5. If the blood is much damaged, transfusion should be resorted to.
6. Never, under any circumstances, give any oils (except turpentine) or fats, as they increase the solubility of phosphorus.

What are the medicinal uses of phosphorus?

Phosphorus is used as a *tonic to promote nutrition*: (1) In various nervous affections associated with anæmia, as, *cerebral exhaustion*, *locomotor ataxia*, *paralysis agitans*, and the various *neuralgiæ*; (2) in general diseases, as, *osteomalacia*, *rickets* and *pernicious anæmia*; (3) in *functional impotence*, and (4) as a substitute for arsenic in certain skin diseases, as *lupus* and *psoriasis*.

What are the preparations of phosphorus and their doses?

PILULÆ PHOSPHORI (*pills of phosphorus*); each pill contains gr. $\frac{1}{100}$ (0.006) of phosphorus; dose j-v pills, t. i. d.

SPIRITUS PHOSPHORI (*spirit of phosphorus—tincture of phosphorus*) contains 1.2 gm. of phosphorus to 1000 c.c. of the preparation, and is used in making

ELIXIR PHOSPHORI (*elixir of phosphorus*); each c.c. contains 0.00025 gm. of phosphorus; dose ℥x-f3j (0.61–4.0).

OLEUM PHOSPHORATUM (*phosphorated oil*) contains phosphorus (1 per cent.) dissolved in ether and almond oil; dose ℥j-v (0.06–0.31).

ZINCI PHOSPHIDUM (*zinc phosphide*) is identical in its actions and uses with phosphorus, but not so active; dose gr. $\frac{1}{20}$ – $\frac{1}{12}$, or even $\frac{1}{2}$ (0.003–0.03) given in pill or granule.

What medicines are incompatible with phosphorus?

Hydrated magnesia, lime-water, cupric sulphate, powdered charcoal and oil of turpentine are chemically incompatible with it.

What substances aid its absorption?

The oils and fats.

What caution should be observed in prescribing it?

Great care should be taken, especially when large doses are prescribed, that the remedy be not administered for a long period continuously, for fear of causing fatty degenerations of the viscera.

MINERAL ACIDS.

What are the general effects of the acids and alkalies upon secretions?

LAW I. *Acids* increase *alkaline* secretions.

LAW II. *Acids* decrease *acid* secretions.

LAW III. *Alkalies* increase *acid* secretions.

LAW IV. *Alkalies* decrease *alkaline* secretions.

How may these laws be applied to the treatment of digestive disturbances?

1. *To decrease excessive acidity of the gastric juice—*

(a) Give an acid before meals to diminish its secretion, or

(b) An alkali after meals to neutralize the acid present.

2. *To increase the acidity of the gastric juice when it is deficient in that quality—*

- (a) Give an alkali before meals to increase the secretion of acid, or
 (b) An acid after meals to supply the deficiency.

Name the mineral acids.

The mineral acids are *sulphuric*, *nitric*, *hydrochloric*, *nitro-hydrochloric* and *phosphoric*.

What are their physiological effects?

Locally the strong acids are *escharotics*, abstracting water from the tissues and combining with the bases. *Sulphuric* and *phosphoric* acids penetrate the tissues more deeply than the others. *Sulphuric* produces a *black*, while *nitric* acid causes a *yellow* eschar.

Internally, in a diluted state and in medicinal doses, they increase the *appetite*, *digestion* and *nutrition* (hydrochloric and phosphoric being more pronounced in their effects). If taken for some time continuously, they diminish the acid of the gastric juice and disorder digestion (sulphuric acid more quickly than the others). Sulphuric acid possesses an astringent influence over the secretions of the skin and bowels; nitric acid, on the contrary, appears to increase the alimentary secretions and that of the liver.

They are *eliminated* by the secretions of the kidneys, bowels and skin. *In large doses* they act as irritant corrosive poisons.

How should poisoning by the mineral acids be treated?

1. Give *alkalies* or *alkaline earths* (tooth powder or soap may be used) to neutralize the acid.
2. Give *eggs*, *milk*, or other *demulcents* to protect the surface.
3. Give *opium* and *nutritive* and *stimulating enemata*, or *intravenous injection* of ammonia, to counteract the resulting depression.
4. As stricture of the oesophagus often follows in these cases if they survive, the passage of an oesophageal bougie should be practiced within 3 or 4 months after the poisoning.

What are the therapeutic uses of the mineral acids?

1. *For their effects on digestion and nutrition.* To prevent excessive formation of acid (both where too much is secreted and where it is formed by fermentation of the food), diluted hydrochloric or phosphoric acid is given before meals; or in *atonic dyspepsia* and *indigestion* due to deficient acid, they may be taken after meals (often

combined with pepsin). *In fevers*, they are used to aid digestion and secretions generally and allay thirst (by stimulating the secretions of the mouth and fauces, thus keeping these parts moist).

2. *For their effects on secretion.* Diluted nitric or nitro-hydrochloric acid is used internally to increase the secretions, and as a bath (fʒiij to water Cj), in *chronic hepatic congestions* and *inflammations*, in *malarial* and *catarrhal jaundice* and in *duodenal catarrh*. As an *astringent* in *diarrhœas*, *dysentery* and *cholera* diluted sulphuric acid is given combined with opium; and to arrest profuse sweating diluted or aromatic sulphuric acid may be temporarily employed.

3. *As a hæmostatic* the same agent is often used in uterine or intestinal hemorrhage and occasionally in purpura.

4. *In the treatment of, or as a preventive to, lead poisoning* diluted sulphuric acid is administered (forming an insoluble lead sulphate).

5. *Locally, as an escharotic*, fuming nitric acid may be applied to *chancre*s, *phagedenic ulcers*, *hospital gangrene* or *gangrene in wounds*. It is also used to remove *warts*, and as an intra-uterine application in various forms of *endometritis*. As a *stimulating lotion* to *unhealthy granulations*, and to check bleeding from hemorrhoids, weak solutions of nitric acid are used.

What are the preparations of the mineral acids, and what are their doses?

1. ACIDUM SULPHURICUM (*sulphuric acid—oil of vitriol*) contains not less than 92.5 per cent. of *absolute sulphuric acid*, and is too strong for internal use.

ACIDUM SULPHURICUM DILUTUM (*diluted sulphuric acid*) contains 10 per cent., by weight, of the *official sulphuric acid*, and is used internally as a *tonic*, *refrigerant*, *astringent*, and *hæmostatic*; dose mʒj-xx (0.123-1.23) freely diluted.

ACIDUM SULPHURICUM AROMATICUM (*aromatic sulphuric acid—elixir of vitriol*) contains 10 per cent. of the *official acid* with alcohol, tincture of ginger, and oil of cinnamon. It is used as a pleasant substitute for the preceding preparation and is given in the same doses.

2. ACIDUM NITRICUM (*nitric acid*) contains 68 per cent. of the *absolute acid*, and is not employed internally.

ACIDUM NITRICUM DILUTUM (*diluted nitric acid*) contains 10 per cent., by weight, of *absolute nitric acid*; dose mʒj-xx (0.123-1.23) freely diluted.

3. **ACIDUM HYDROCHLORICUM** (*hydrochloric acid—muriatic acid*) contains nearly 32 per cent., by weight, of *absolute acid*, and is not used internally.

ACIDUM HYDROCHLORICUM DILUTUM (*diluted hydrochloric acid—diluted muriatic acid*) contains 10 per cent. of the absolute acid. It is (probably) a normal constituent of the gastric juice, and is used as an aid to digestion; dose $\mathfrak{m}_{\text{v-xx}}$ (0.31–1.23) freely diluted.

4. **ACIDUM NITROHYDROCHLORICUM** (*nitrohydrochloric acid—nitromuriatic acid—aqua regia*) is used as a stomachic tonic and to increase the secretions of the liver and intestines; dose $\mathfrak{m}_{\text{ij-x}}$ (0.123–1.23) freely diluted.

ACIDUM NITROHYDROCHLORICUM DILUTUM (*diluted nitrohydrochloric acid—diluted nitromuriatic acid*); dose $\mathfrak{m}_{\text{ij-xx}}$ (0.123–1.23) freely diluted.

5. **ACIDUM PHOSPHORICUM** (*phosphoric acid*) contains 85 per cent., by weight, of absolute ortho-phosphoric acid, and is not used internally.

ACIDUM PHOSPHORICUM DILUTUM (*diluted phosphoric acid*) contains 10 per cent., by weight, of absolute ortho-phosphoric acid, and is chiefly used as an aid to digestion; dose $\mathfrak{m}_{\text{v-xxx}}$ (0.31–1.85) freely diluted.

What medicines are incompatible with the mineral acids?

1. Alkalies and their carbonates and the salts of lime and lead are incompatible with the mineral acids, which also decompose salts of the vegetable acids, uniting with their bases.

2. The mercurial salts should not be given during the administration of hydrochloric or nitro-hydrochloric acids or their preparations, as they may react with the mercury and form corrosive sublimate.

How and when are these acids administered?

They are administered before or after meals, according to the indications present, and may be given freely diluted with water either alone or with glycerin and syrup, or with some of the bitters.

They should be taken through a glass tube to prevent injury to the teeth, or the mouth may be washed out immediately after their use with some weak alkaline solution.

ACIDUM LACTICUM—LACTIC ACID.

What is lactic acid?

Lactic acid is a syrupy liquid, formed by the souring of milk.

What are its effects and uses?

It is a normal ingredient of the gastric juice. In *medicinal doses* it increases the appetite, aids digestion and stimulates nutrition. In *large doses* (f3j) or *long continued* it disorders digestion, causing flatulency and epigastric pain. *Injected into the peritoneal cavity* (dog's) it produces endocarditis, and when given for a very long time (in diabetes) it has caused acute rheumatism. *Lactic acid is used to diminish the acidity* of the gastric juice, or to *replace the deficiency of the acid*; to aid digestion and nutrition in *oxalic, phosphatic or uric acid diathesis*; in *diabetes*, and *locally* as a solvent for the false membranes of croup and diphtheria. It has been claimed that it is a hypnotic in *insomnia* due to nervous excitement, but this is still *sub judice*.

What are its preparations and doses?

ACIDUM LACTICUM (*lactic acid*) contains 75 per cent. of absolute lactic acid. Dose $\mathfrak{m}_{\text{V-xxx}}$ (0.30–1.85) freely diluted and often prescribed in a mixture with pepsin.

ORDER III.—ALTERATIVES.

What are alteratives?

Alteratives are medicines which, without exerting any very perceptible action on any particular organ, so modify the nutritive processes as to enable nature to restore healthy action in many diseased conditions.

Mention the principal alteratives.

The preparations of mercury and of gold; iodine and the iodides; the arsenical preparations; cod-liver oil; the phosphates and hypophosphites; the chlorides of ammonium and calcium; colchicum; sarsaparilla; guaiac; and perhaps mezereon; menispermum; calendula; sassafras, stillingia, dulcamara, and lappa.

HYDRARGYRUM—MERCURY.

What is mercury?

Mercury is a liquid metal principally obtained from the native sulphide or cinnabar.

What are the physiological effects of mercury?

Metallic mercury (quicksilver) is inert either when swallowed or applied to the skin, but if retained in the alimentary canal or after prolonged contact with the skin, it often becomes converted into an oxide and produces constitutional effects. *When inhaled* in a state of vapor it frequently causes salivation, ulceration of the mouth, necrosis and wasting; or in other cases loss of memory, vertigo, shaking palsy and other nervous symptoms.

Minute doses of mercurials, not too frequently repeated, increase the number of red corpuscles in the blood and improve nutrition. In *small and repeated doses* they stimulate the secretions of the skin and mucous membranes, the liver and kidneys and the salivary glands and pancreas, and promote absorption. *Corrosive sublimate* is an hepatic stimulant of considerable power, but the other preparations probably increase the flow of bile by irritating the duodenal mucous membrane, thus causing reflex contractions of the gall bladder. By stimulating the liver, *corrosive sublimate* increases the formation of urea, and thus proves diuretic. It also feebly stimulates the intestinal glands. *In moderate doses* the mercurials are cathartic (especially calomel and blue mass). *In large doses*, or continued for some time, the secretions are much increased and become pathological; the gums are swollen and tender, the mouth sore, the teeth feel elongated, salivation occurs, which becomes profuse if the medicine is continued; the breath is offensive, a metallic taste is experienced; the proportion of albumen, fibrin and red corpuscles to the fluid of the blood is diminished; the body wastes, and, if the abuse of mercury be continued, mercurial fever, diarrhoea, ulceration and sloughing of the gums, loosening of the teeth and sometimes necrosis of the alveolar processes, albuminuria, marasmus and grave nervous symptoms occur. Salivation (ptyalism) is more easily caused by blue mass and calomel, less so by corrosive sublimate and mercury with chalk.

Corrosive sublimate, beside its cholagogue and diuretic properties,

retards hyperplasia of connective tissue, is a powerful gastro-intestinal irritant, and locally the best antiseptic, proving rapidly destructive to germs.

What are the symptoms and treatment of poisoning by corrosive sublimate?

The symptoms are violent gastro-enteritis with vomiting and purging, bloody stools, abdominal pain and tenderness, albuminous urine and death from collapse. The treatment consists in (1) giving an antidote, *albumen* is the best (*white of eggs*, milk or wheat flour); (2) giving an emetic; and (3) the free use of opiates and demulcents.

What are the therapeutic uses of mercury?

Mercury is used (1) as an *antisyphilitic*, either by the mouth (yellow iodide, calomel, blue mass, corrosive sublimate), fumigation (calomel), inunction (mercurial ointment or oleate), or hypodermic injection (albuminate, peptonate, or even calomel or corrosive sublimate). It is of most use in secondary syphilis, the iodides (alone or combined with the red iodide or corrosive chloride) giving better results in tertiary syphilis.

(2) As an *antiplastic and sorbefacient* in *acute glandular affections*, *iritis* and the *latter stages of inflammations* after plastic effusion has occurred. In *diphtheria* and *membranous croup* calomel is advantageously given in repeated doses guarded by minute doses of opium. In the latter disease the subsulphate is often used as an emetic. Mercurials are useful in ileo-colitis of infants (gray powder) and in acute dysentery (corrosive sublimate). (3) As an *indirect tonic* and to *stimulate secretions*, calomel, gray powder or blue mass are used in dyspepsia attended with constipation and torpid liver, and in the early stages of acute diseases. (4) As a *purgative* calomel and blue mass are often used, frequently followed by some saline. (5) *Locally* as *antiseptics* mercuric chloride or iodide are almost indispensable in surgical practice. Various preparations are used locally in various *skin diseases*, especially of parasitic origin, to *promote absorption* in *goitre*, *enlarged spleen*, *chronic epididymitis* and *orchitis* and other glandular swellings. The ointments are also used in conjunctivitis.

What are the preparations of mercury and their doses?

MASSA HYDRARGYRI (*mass of mercury—blue mass*), dose as a

laxative, gr. v–xv (0.33–1.0) ; as an *alterative*, *antisyphilitic*, or *sorbefacient*, gr. ss–ijj (0.03–0.2), t. i. d., guarded by opium, and often combined in the treatment of syphilis with iron or quinine.

UNGUENTUM HYDRARGYRI (*mercurial ointment—blue ointment*). Not used internally. Used externally for *inunction* in syphilis ; as a *sorbefacient*, to *destroy pediculi* ; and as a dressing to *syphilitic ulcers*.

EMPLASTRUM HYDRARGYRI (*mercurial plaster*), used locally as a *sorbefacient*.

EMPLASTRUM AMMONIACI CUM HYDRARGYRO (*plaster of ammoniac with mercury*), is more stimulating than the former.

HYDRARGYRUM CUM CRETÂ (*mercury with chalk—gray powder*), contains mercury, 38 parts, with prepared chalk and sugar of milk, and is used as an *alterative* and *antisyphilitic* ; dose for an adult gr. j–v (0.06–0.32), for a child gr. $\frac{1}{4}$ (0.02) t. i. d. ; as a mild laxative gr. v–xv (0.33–1.0) (for a child gr. ij–v (0.2–0.33)). It should not be prescribed in pill form.

HYDRARGYRI OXIDUM RUBRUM (*red mercuric oxide*) and

HYDRARGYRI OXIDUM FLAVUM (*yellow mercuric oxide*) are rarely used except in the form of

UNGUENTUM HYDRARGYRI OXIDI RUBRI (*ointment of red mercuric oxide—red precipitate ointment*) and

UNGUENTUM HYDRARGYRI OXIDI FLAVI (*ointment of yellow mercuric oxide*), each containing 10 per cent. of mercuric oxide, and which are applied in *granular* and other forms of *conjunctivitis* and in *corneal ulcers*.

OLEATUM HYDRARGYRI (*mercurial oleate*), contains 10 per cent. of the yellow oxide dissolved in oleic acid. Used chiefly for *inunction* in syphilis.

HYDRARGYRI CHLORIDUM MITE (*mercurous chloride—calomel*), dose, as a *purgative* or *anthelmintic*, gr. ss–j–x (0.03–0.06–0.6) at bedtime ; as an *antisyphilitic* gr. $\frac{1}{10}$ –j (0.006–0.06), t. i. d., with opium ; as an *alterative* and *absorbent* gr. ss–j (0.03–0.06) every 2 or 3 hours ; for its effect on *secretions* gr. ss–j (0.03–0.06) t. i. d. ; to *arrest vomiting* gr. $\frac{1}{10}$ –j (0.006–0.06) combined with sodium bicarbonate and frequently repeated. It is used externally as a dusting powder to the conjunctiva, and as an ointment in various skin affections. It is also used by *fumigation* in syphilis.

HYDRARGYRI CHLORIDUM CORROSIVUM (*mercuric chloride—cor-*

rosive sublimate), used as an *alterative* and *antisyphilitic*; dose gr. $\frac{1}{16}$ – $\frac{1}{8}$ (0.004–0.008) by the mouth, gr. $\frac{1}{48}$ – $\frac{1}{16}$ (0.001–0.004) by hypodermic injection. Locally, in weak solutions as an injection in *gonorrhœa* after the more acute stage has passed (gr. $\frac{1}{2}$ – $\frac{1}{8}$ –f $\overline{3}$ viii (0.005–0.008–236.5)) or as a *tenuicidal* in ringworm, etc., or for the destruction of pediculi. As an *antiseptic* gr. vijss to hot water Oij = 1 part to 2000.

HYDRARGYRI IODIDUM FLAVUM (*mercurous iodide*), dose as an *antisyphilitic* gr. $\frac{1}{8}$ –j (0.008–0.065) combined with opium in pill. Used also locally, in the form of ointment, to syphilitic ulcers.

HYDRARGYRI IODIDUM RUBRUM (*mercuric iodide*), dose as an *alterative* or *antisyphilitic* gr. $\frac{1}{16}$ – $\frac{1}{4}$ (0.004–0.016). It is often combined with potassium iodide in the treatment of tertiary syphilis, and is used locally in the form of ointment to promote the absorption of glandular swellings, goitre, enlarged spleen, etc. As an *antiseptic* it is used in solution of gr. iv–vijss (0.259–0.454) to hot water Oij (950.3). Potassium iodide increases its solubility.

HYDRARGYRI CYANIDUM (*mercuric cyanide*) is used in syphilis as a substitute for corrosive sublimate, dose gr. $\frac{1}{16}$ – $\frac{1}{12}$ (0.004–0.005).

HYDRARGYRUM AMMONIATUM (*ammoniated mercury—white precipitate*) is only used externally as

UNGUENTUM HYDRARGYRI AMMONIATI (*ointment of ammoniated mercury*), which contains 10 per cent. of ammoniated mercury, and is used especially in parasitic skin diseases and to destroy pediculi.

HYDRARGYRI SUBSULPHAS FLAVUS (*yellow mercurial subsulphate—turpeth mineral*), dose as an *alterative* gr. $\frac{1}{4}$ –ss (0.016–0.032); as an *emetic* in croup, gr. j–v (0.065–0.324), in syrup, repeated every 10 or 15 minutes until free vomiting occurs.

UNGUENTUM HYDRARGYRI NITRATIS (*ointment of mercurial nitrate—citrine ointment*), used externally (usually diluted with other ointment) in various skin affections, especially those of parasitic origin, to destroy pediculi and as a dressing for venereal ulcers.

LIQUOR HYDRARGYRI NITRATIS (*solution of mercuric nitrate—acid nitrate of mercury*), used as an *escharotic*. Diluted with water, it forms a useful stimulating application to indolent ulcers.

What substances are incompatible with the mercurial preparations?

The *metallic preparations* and *calomel* are incompatible with the

chlorides, hydrochloric and nitro-hydrochloric acids, which are apt to form with them corrosive sublimate. *Calomel* is also incompatible with the alkalies, their carbonates, alkaline earths, soap and the hydrosulphates. *Corrosive sublimate* is decomposed by nearly everything, and *green mercurous iodide* is converted into the more active red mercuric iodide by combination with iodine and other iodides.

AURUM—GOLD.

What are the physiological effects of the salts of gold?

Locally they are caustic. *Internally* in *small doses* they increase the appetite, digestion, glandular secretions, and excretions of urine and perspiration; they stimulate the nervous system, and particularly the spinal cord; they stimulate the sexual organs, and in women (probably) increase the menstrual flow. In *larger doses*, or after prolonged use, they disorder digestion, cause salivation without ulceration of the gums, diminish the oxygen-carrying function of the red globules, and stimulate the lymphatic system, thus causing rapid wasting. In *overdoses* they are gastro-intestinal irritants, the *symptoms* and *antidote* being the same as those of poisoning by corrosive sublimate.

What are the medicinal uses of the gold salts?

They are used as substitutes for corrosive sublimate in the treatment of *tertiary syphilis*; as *alteratives* in scrofula, chronic Bright's disease, myelitis, chronic ovarian inflammations and neuralgia; and as *aphrodisiacs* in functional impotence.

The so-called "*bichloride of gold*" or "*Keely cure*" does not consist in the administration of gold bichloride, as no such salt is known. The cures are probably largely effected by suggestion. It frequently fails to effect a cure, and many relapses occur.

What are the preparations and doses of the gold salts?

AURI ET SODII CHLORIDUM (*gold and sodium chloride*) is the only official preparation; dose gr. $\frac{1}{30}$ – $\frac{1}{10}$ (0.002–0.006), in pill or capsule.

IODUM—IODINE.

What is iodine?

Iodine is a non-metallic element, obtained from the ashes of seaweed.

What are the physiological effects of iodine and the iodides?

Iodine is a *disinfectant* and *antiseptic*. *Locally* applied, it is an *irritant caustic* to the skin and mucous membranes, staining the former yellow and causing superficial inflammation and desquamation, and sometimes even vesication. The *vapor*, when inhaled, is irritant to the air passages. In sufficient amounts it is a gastrointestinal irritant. It enters the blood readily, and, combining with sodium or potassium, corresponds, in its constitutional effects, to the *iodides*. They *stimulate* the *lymphatic system* and cause the *absorption* especially of newly-formed tissues and diseased cells, thus removing indolent swellings, inflammatory and syphilitic deposits and various metallic poisons, as lead, mercury, etc., which may have been deposited in the tissues. When *too long administered*, they cause great emaciation and depression of the vital powers. They are rapidly *eliminated* by the mucous membranes (irritating them in their passage), salivary glands and kidneys.

Given in *large quantities*, they cause *iodism*, *i. e.*, malaise, some fever, frontal headache, coryza, lachrymation, sore throat, difficult swallowing, various eruptions on the face and shoulders and antaphrodisiac effects, sometimes permanent. The susceptibility varies greatly in different persons, but these symptoms may in great measure be prevented by copious draughts of water frequently taken during the course of treatment.

What is the treatment of acute poisoning by free iodine?

A decoction of starch is the *antidote*, followed by emetics, and opium and demulcents.

What are the therapeutic uses of iodine and the iodides?

They are employed as *alteratives* and *absorbents* in the active manifestations of *tertiary syphilis* (iodides chiefly), in the treatment of *scrofulous* and *glandular enlargements*, in the early stages of

cirrhosis of the liver or kidneys, in the various forms of *sclerosis* of the nervous centres, and in *chronic poisoning* by mercury, lead and other metals; to aid the absorption of *lingering pneumonic consolidation* (ammonium iodide) and of *plastic* and *serous* inflammatory *exudations* (potassium iodide), in *aneurisms* (large doses of potassium iodide), in *chronic rheumatism* and in *malaria* (iodine preparations).

For their effects on the *mucous secretions* the iodides are given in *chronic bronchitis*, *spasmodic asthma*, *catarrhal pneumonia*, *acute catarrh*, *hay fever* and *duodenal catarrh*. It is highly recommended internally in the treatment of *typhoid fever*, and has been used to check the *vomiting of pregnancy*. *Locally*, iodine is used by inhalation in *catarrh*, *coryza* and *hay fever*, and is applied to aid *absorption* and as a *counter-irritant* to *chronic inflammations* and *chronic glandular* and other *swellings*. The *tincture* is *injected* undiluted into the serous cavity after the removal of the fluid in *empyema*, *hydrocele*, etc., and is applied to the uterine cavity in various forms of *endometritis*.

What are the preparations and doses of iodine and the iodides?

LIQUOR IODI COMPOSITUS (*compound solution of iodine—Lugol's solution*), containing iodine (5 parts), potassium iodide (10 parts), and water (85 parts); dose ℥v–xv (0.308–0.924) t. i. d.

TINCTURA IODI (*tincture of iodine*), containing 8 per cent. of iodine; dose ℥ij–xv (0.123–0.924) t. i. d. It is chiefly used externally.

UNGUENTUM IODI (*iodine ointment*) contains iodine (4 parts), potassium iodide (1 part), water (2 parts), and benzoinated lard (93 parts).

POTASSII IODIDUM (*potassium iodide*), dose gr. ij–xv–ʒj (0.130–0.972–3.888) t. i. d.

UNGUENTUM POTASSII IODIDI (*ointment of potassium iodide*) is much less powerful than iodine ointment.

AMMONII IODIDUM (*ammonium iodide*), dose gr. ij–xv (0.132–0.972).

SODII IODIDUM (*sodium iodide*), dose gr. v–xxx (0.324–1.944).

SULPHURIS IODIDUM (*sulphur iodide*) is but little used; dose gr. ½–j (0.032–0.065).

STRONTII IODIDUM (*strontium iodide*) is supposed to be as efficacious as the other iodides and to be less likely to cause iodism; dose gr. v-x (0.324-0.648).

The *mercurial iodides* and *iron iodide* have already been mentioned. *Arsenic, zinc and lead* iodides will be described in their appropriate places.

What are the incompatibilities of these preparations?

The *iodides* are incompatible with the acids and acidulous salts, with the soluble metallic salts generally; with most of the alkaloids and with sweet spirits of nitre. Iodine is precipitated from the *tincture* on the addition of water.

How are iodine and the iodides administered?

Lugol's solution may be given in syrup and water. The iodides may be administered in compressed pill or capsule, or they may be given dissolved in water flavored with orange syrup, or in one of the bitter tinctures, or in compound syrup of sarsaparilla. Nothing effectually disguises their unpleasant taste.

ARSENUM—ARSENIC.

What is arsenic?

Arsenic is a metallic element, found in various ores combined with other metals as an arsenide.

Mention some of the tests for arsenic.

When *in powder*, if heated with charcoal in a small test tube, it sublimes and condenses in the form of a brilliant steel gray ring of minute octahedral crystals, soluble in sodium or calcium hypochlorite. When *in solution*, the addition of sulphuretted hydrogen or ammonium sulphide produces a lemon-yellow arsenic trisulphide, insoluble in diluted hydrochloric acid, but soluble in a solution of ammonium carbonate: also the addition of ammonia water followed by a solution of silver nitrate, gives a canary-yellow silver arsenite. Marsh's and Reinsch's tests should also be used.

What are the physiological effects of arsenical preparations?

Metallic arsenic is inert, but when swallowed may be converted into arsenous acid and cause poisoning. No matter how administered, it has a selective action on the gastro-intestinal and broncho-pulmonary mucous membranes. *Locally*, arsenous acid is a painful *caustic*, and when in sufficient amount produces so much inflammation that absorption cannot take place. The *inhalation* of arsenical fumes (as from wall papers) have caused poisoning. Given *internally in small doses* it improves the appetite, digestion, nutrition and condition of the blood; stimulates the secretions of the alimentary canal and increases peristalsis; slightly stimulates the respiratory and circulatory centres and the cerebral functions; and lessens the excretion of carbonic acid and urea. *In larger doses* (not toxic), or when *taken for some time continuously*, it irritates the mucous membranes, disorders digestion, gives rise to *epigastric pain*, *nausea* and vomiting, diarrhoea, tenesmus and sometimes dysentery, palpitation, oppressed breathing, *œdema of the eyelids*, albuminuria, cutaneous eruptions, tremor and disordered sensation. It is *eliminated* by the mucous membrane, liver, kidneys and skin. *Tolerance* is sometimes established, and very large doses may be taken without poisonous effect.

What are the symptoms, pathological anatomy and treatment of acute arsenical poisoning?

The symptoms of the gastro-intestinal variety of poisoning are intense burning pain in the epigastrium, radiating over the abdomen, violent vomiting, great thirst, bloody and offensive stools, strangury and bloody urine, rapid feeble heart, oppressed breathing and collapse, sometimes simulating cholera. *Those of the cerebral variety* are sudden, profound insensibility and coma with collapse.

Pathological Anatomy.—*Alimentary mucous membrane* congested, eroded, softened and ecchymotic; *broncho-pulmonary mucous membrane* and lungs congested; fatty degeneration of the *liver, kidneys* and other organs.

Treatment, (1) give a *prompt emetic* or use the stomach pump; (2) give as antidotes *hydrated ferric oxide* or the *hydrated ferric oxide with magnesia*, and continue the emetics; (3) *mucilaginous*

drinks to protect the surfaces; (4) large amounts of diluents to favor elimination.

Prognosis.—Death often results long afterward, from the pathological conditions produced by the poison.

What are the medicinal uses of the arsenical preparations?

Arsenic is used as a *tonic* and *alterative* in *anæmia* and *chlorosis*, persistent *malaria*, *rheumatoid arthritis* and *chronic rheumatism*, and to delay the formation of connective tissue in the various *scleroses* and the growth of *carcinomata*. In minute doses it is used as a gastric tonic in *irritative dyspepsia*, *gastric ulcer* and *cancer*, and in *gastralgia* and *enteralgia*. (Fowler's Solution.) In *emphysema*, *chronic bronchitis* and *phthisis* it is of value. For its nervous action it is used in *chorea*, *neuralgic*, *paralysis agitans* and *melancholia*. In *chronic skin diseases*, as *psoriasis*, *eczema* and *acne*, it is of the greatest service, and in *diabetes* and *sexual debility* it is highly recommended. *Locally*, arsenous acid has been used as a caustic in *lupus*, *onychchia maligna*, *cancer*, and to destroy the nerves of carious teeth. Cigarettes made of bibulous paper saturated with a solution of sodium arsenate are smoked for the relief of *asthma*.

What are the preparations and doses of arsenic?

ACIDUM ARSENOSUM (*arsenous acid—white arsenic*), dose gr. $\frac{1}{16}$ – $\frac{1}{12}$ (0.004–0.005) in pills t. i. d.

LIQUOR POTASSII ARSENITIS (*solution of potassium arsenite—Fowler's Solution*) contains arsenous acid gr. $\frac{6}{10}$ in each fʒj of solution; dose $\mathfrak{m}\text{j}$ –x (0.061–0.616) freely diluted.

LIQUOR ACIDI ARSENOSI (*solution of arsenous acid*), dose $\mathfrak{m}\text{j}$ –x (0.061–0.616) freely diluted.

SODII ARSENAS (*sodium arsenate*) is milder in its action than the foregoing preparations; dose gr. $\frac{1}{12}$ – $\frac{1}{4}$ (0.005–0.016).

LIQUOR SODII ARSENATIS (*solution of sodium arsenate*), dose $\mathfrak{m}\text{ij}$ –x (0.123–0.616) freely diluted.

ARSENI IODIDUM (*arsenic iodide*), dose gr. $\frac{1}{8}$ (0.008) t. i. d.

LIQUOR ARSENI ET HYDRARGYRI IODIDI (*solution of arsenic and mercuric iodide—Donovan's Solution*) contains 1 per cent. of arsenic iodide and of mercuric iodide each, and is used in obstinate syphilis and in papular and squamous skin diseases; dose $\mathfrak{m}\text{ij}$ –x (0.123–0.616) freely diluted.

What medicinal substances are incompatible with these preparations?

The *salts of iron, magnesium and calcium* and the *astringents* are chemically incompatible with the preparations of arsenic. *Fowler's Solution* should not be prescribed with infusions or decoctions of *cinchona* nor *Donovan's Solution* with the alkalies, or the salts of the alkaloids.

How and when is arsenic administered?

As individual susceptibility to arsenic varies greatly,* it is better to give a small dose at first. When it is determined that the patient can take arsenic, large doses should be given until some constitutional symptoms occur, when the medicine is discontinued for a few days. Or full doses of the liquid preparations gradually decreased to the minimum and as gradually increased may be given when it is desirable to continue the medicine for some time. Arsenic is less apt to disorder the stomach when freely diluted and taken after eating.

OLEUM MORRHUÆ—COD-LIVER OIL.

What is cod-liver oil?

Cod-liver oil is a *fixed oil* obtained from the fresh *livers* of *Gadus morrhue*, or codfish (*Class Pisces; Ord. Teleostia; Fam. Gadida*).

What are its principal chemical constituents?

It contains various fatty acids, several biliary principles, *gaduin*, *iodine*, *chlorine* and traces of *bromine*.

What are its physiological effects?

Cod-liver oil is more of a *food* than a *medicine*. It is readily absorbed, whether rubbed into the skin or taken internally. Oil is digested in the intestines (not in the stomach), and is readily assimilated, increasing the deposit of fat in the tissues and the number of corpuscles in the blood. In suitable doses it *improves* the *appetite* and *digestion* and *increases nutrition*. In large doses, or in very hot weather, it causes nausea and diarrhœa.

* The author has seen gastro-enteric symptoms follow the administration of one dose of Fowler's Solution ℥j.

What are its therapeutic applications?

Cod-liver oil is used as a *nutrient* in chronic wasting diseases, particularly in *phthisis* (the earlier it is given, the more pronounced are its effects), also in *rickets*, *scrofula*, *chlorosis*, *rheumatoid arthritis*, *chronic rheumatism*, *chronic bronchitis*, *emphysema* and *chronic intestinal diseases* leading to mal-nutrition.

It is applied *locally for its constitutional effects* when it cannot be taken internally, and also in squamous skin affections.

What is the dose of cod-liver oil, and how should it be administered?

OLEUM MORRHUÆ (*cod-liver oil*) is given in doses of f3j-f3ss (3.696–15.0) t. i. d., and preferably $\frac{1}{2}$ to 2 hours after meals. The *pure oil* is the best form for administration. If it nauseates, it may be given in emulsion with the hypophosphites, or with mucilage of acaciæ and oil of lemon. The addition of a few drops of ether renders it more easy of digestion.

THE PHOSPHATES AND HYPOPHOSPHITES.

What is the chemical difference between a phosphide, a phosphate and a phosphite?

A *phosphide* is formed by the direct union of *phosphorus*; a *phosphate* by the union of *phosphoric acid*, and a *phosphite* by the union of *phosphorous acid* with a base.*

What are the physiological actions of the phosphates and hypophosphites?

They increase the nutrition of the body.

Calcium phosphate exists normally in all the tissues of the body, and hence is an important nutrient. It is to this salt that bone owes its hardness.

* In other words, whenever a salt ends in *ide* it is formed by the union of an element with a base (thus, iron *oxide*, iron and oxygen; iron *phosphide*, iron and phosphorus; iron *iodide*, iron and iodine; iron *sulphide*, iron and sulphur); when a salt ends in *ate*, it shows the union of an *acid ending in ic* with a base (thus, iron *phosphate*, iron and *phosphoric acid*; iron *sulphate*, iron and *sulphuric acid*; iron *nitrate*, iron and *nitric acid*); when a salt ends in *ite*, it is composed of an *acid ending in ous* and a base (thus, sodium *sulphite*, sodium and *sulphurous acid*; amyl *nitrite*, amyl and *nitrous acid*, etc.).

What are the medicinal uses of these substances?

They are used as *alteratives* and *nutrients* in diseases of mal-nutrition, especially when these salts are wanting in the system, as in *rickets*, *mollities ossium*, *caries* and *necrosis*, *delayed union* of fractures, *superlactation*, *prolonged suppuration*, *anæmia*, *chronic diarrhœa* and *chronic phthisis*. The *hypophosphites* have been lauded in the treatment of phthisis, but probably act only as nutrients.

What are the preparations of these salts and their doses?

CALCII PHOSPHAS PRÆCIPITATUS (*precipitated calcium phosphate*), a very insoluble preparation, and apt to form intestinal concretions; dose gr. v-x (0.324-0.648). It may be given in milk.

SYRUPUS CALCII LACTOPHOSPHATIS (*syrup of calcium lactophosphate*); the addition of lactic acid renders the calcium phosphate much more soluble; dose f3j-iv (4.0-15.0) t. i. d.

CALCII HYPOPHOSPHIS (*calcium hypophosphite*); POTASSII HYPOPHOSPHIS (*potassium hypophosphite*); SODII HYPOPHOSPHIS (*sodium hypophosphite*), may each be given in doses of gr. x-xxx (0.648-1.944) t. i. d. Usually administered as

SYRUPUS HYPOPHOSPHITUM (*syrup of hypophosphites*), dose f3j-ij (4.0-8.0) t. i. d.

SYRUPUS HYPOPHOSPHITUM CUM FERRO (*syrup of hypophosphites with iron*) contains 1 per cent. of ferrous lactate; dose f3j-ij (4.0-8.0) t. i. d.

The *ferric phosphates* and *hypophosphite* have been considered under iron; *sodium phosphate* will be mentioned with the cathartics.

CALCII CHLORIDUM—CALCIUM CHLORIDE.

What are the effects, uses, and doses of this salt?

Calcium chloride is a gastro-intestinal irritant, and is supposed to possess *alterative* powers in *strumous affections*, *glandular enlargements*, *uterine fibroid* and *ovarian tumors*. Dose gr. x-xx (0.648-1.296) in milk or water.

AMMONII CHLORIDUM—AMMONIUM CHLORIDE.

What are the effects, uses and doses of ammonium chloride?

In addition to the general effects of the ammonium salts (*q. v.*) it

is an *alterative* and *expectorant*. *Locally* it is an irritant. *Internally*, in *small doses*, it stimulates the mucous membranes, increasing their secretions and epithelial exfoliation. It stimulates the intestinal glands and, perhaps, the liver. It aids the excretion of waste products and increases all the solids in the urine except uric acid. It is said to increase the menstrual flow in women. In *large doses* it irritates the stomach and intestines and proves purgative.

It is used as an *expectorant* (*q. v.*) and as an *alterative* in *enlarged glands*, *plastic exudations* (given internally and used locally), in *uterine fibroids*, *functional amenorrhœa*, in *muscular* and *chronic rheumatism* and *myalgia*, and in the early stages of *cirrhosis of the liver*. In *albuminoid* or *waxy infiltration of the liver*, *kidneys* and *spleen* it is a very valuable agent. It is also used in *neuralgia* and *neuralgic headache*. *Externally* it is applied as a *refrigerant* (℥j (31.10) to water Oss (120.0)) in the early stages of inflammation and in the later stages as a sorbefacient. Dose gr. v—xxx (0.324–1.944) every three or four hours, in compressed pills, capsules, or in solution with fluid extract of liquorice, which somewhat disguises its unpleasant taste.

COLCHICUM.

What is colchicum?

Colchicum is the *corm* and *seed* of *C. autumnale* or meadow saffron (*Nat. Ord.* Liliacæe), a native of Europe.

What alkaloid does it contain?

It contains *colchicine*, which is converted by mineral acids into *colchicin*.

What are its physiological effects?

Locally applied it is *irritant*. *Internally*, *colchicum*, or its *alkaloid* in *small doses*, increase the secretions generally, particularly the urine, increasing greatly the amount of urea and uric acid eliminated; in *larger doses* it causes nausea and vomiting, frequent purging, and depresses the heart's action; in *excessive doses* it acts as an irritant poison, producing violent vomiting and purging, great depression, collapse and death by failure of respiration.

How should poisoning by colchicum be treated?

(1) Give emetics and cathartics; (2) tannic acid, as a partial anti

dote ; (3) demulcents, to protect the surfaces, and (4) opium and stimulants, to counteract the resulting depression.

What are the therapeutical uses of colchicum ?

Colchicum is used as an *alterative* and *diuretic* in *acute gout* (often combined with magnesia and magnesium sulphate (Scudamore's draught) or with potassium iodide or other alkali) and in *rheumatism*. As a *diuretic* it is employed in ascites and other varieties of dropsy. As a *cholagogue* it is given in congestions of the portal circulation.

What are the preparations and doses of colchicum ?

EXTRACTUM COLCHICI RADICIS (*extract of colchicum root*), dose gr. j-ij (0.065-0.130) in pill.

EXTRACTUM COLCHICI RADICIS FLUIDUM (*fluid extract of colchicum root*), dose ℥ij-iv (0.123-0.246).

EXTRACTUM COLCHICI SEMINIS FLUIDUM (*fluid extract of colchicum seed*), dose ℥ij-vj (0.123-0.40).

TINCTURA COLCHICI (*tincture of colchicum*), made from the seed ; dose fʒss-j (2.0-4.0).

VINUM COLCHICI RADICIS (*wine of colchicum root*), dose ℥v-xx (0.308-1.232) ; as a purgative fʒss (2.0).

VINUM COLCHICI SEMINIS (*wine of colchicum seed*), dose ℥v-xxx (0.308-2.0)—the best preparation.

Colchicine (not official), dose for hypodermic use gr. $\frac{1}{50}$ (0.001).

SARSAPARILLA.

What is sarsaparilla ?

Sarsaparilla is the *root* of *smilax officinalis* and other species of *smilax* (*Nat. Ord. Smilacæ*), prickly shrubs of Mexico, Central and South America.

What special chemical principle does it contain ?

A glucoside, called *smilacin* or *parillin*, which resembles saponin.

What are its effects and uses ?

It has a very slight diaphoretic action, and in large doses is emetic and purgative. It is popularly supposed to depurate the blood. It has been used as an alterative in tertiary syphilis and in chronic rheumatism and various cachectic conditions.

What are its preparations and doses?

EXTRACTUM SARSAPARILLÆ FLUIDUM (*fluid extract of sarsaparilla*), dose f3ss (2.0) t. i. d.

EXTRACTUM SARSAPARILLÆ FLUIDUM COMPOSITUM (*compound fluid extract of sarsaparilla*), containing also sassafras, mezereon, and glycyrrhiza; dose f3j (4.0).

DECOCTUM SARSAPARILLÆ COMPOSITUM (*compound decoction of sarsaparilla*), containing the above ingredients and guaiac wood; dose f3ij-vj (80.0–200.0) t. i. d.

SYRUPUS SARSAPARILLÆ COMPOSITUS (*compound syrup of sarsaparilla*), containing guaiac wood, pale rose, senna, glycyrrhiza, sassafras, anise, and gaultheria. It decomposes corrosive sublimate, converting it into calomel. Dose f3ss-j (15.0–30.0) t. i. d.

GUAIACI LIGNUM—GUAIAIC WOOD.

GUAIACI RESINA—GUAIAIC.

What are these substances?

Guaiacum wood, or *lignum vitæ*, is the *heart wood* of *G. officinale* and *G. sanctum* (*Nat. Ord. Zygophyllacæ*), evergreen trees of S. America.

Guaiac is the *resin* of the wood of *G. officinale*, and is composed of *guaiaconic*, *guaiac* and *guaiacresinic acids*, *guaiac beta-resin*, *guaiac yellow* and *guaiacín*.

What are the effects, uses, preparations and doses of these substances?

Guaiac wood and resin slightly increase the secretions of the skin and mucous membranes, and in large doses prove emetic and cathartic. They have been used as *alteratives* in syphilis and chronic rheumatism, and also as *emmenagogues* in amenorrhœa and congestive dysmenorrhœa. In *tonsillitis* the tincture (f3ss (2.0) every 3 or 4 hours) is of value.

The preparations are TINCTURA GUAIACI (*tincture of guaiac*) and TINCTURA GUAIACI AMMONIATA (*ammoniated tincture of guaiac*); dose of either, f3ss-ij (2.0–8.0), 3 or 4 times daily, in milk.

MEZEREUM—MEZEREON.

What is mezereon?

It is the *bark* of *daphne mezereum* and other species of *daphne*

(*Nat. Ord.* Thymelacææ), and contains daphnin (a bitter glucoside) and an acid resin. *Locally* it is irritant; *internally in medicinal doses* it increases the secretions, especially of the salivary glands, skin and kidneys; *in overdoses* it is an irritant poison and may cause death. It is *used* (generally with sarsaparilla) as an alterative in chronic rheumatism, tertiary syphilis and strumous affections, and as a *masticatory*, it has been chewed in paralysis of the muscles of the tongue and deglutition.

EXTRACTUM MEZEREI FLUIDUM (*fluid extract of mezercon*), dose \mathfrak{m}_x (0.616).

What is memispermum?

It is the *rhizome* and *roots* of *M. canadense*, yellow parilla or Canada moonseed (*Nat. Ord.* Menispermacææ), and contains *berberine*. It is supposed to be *tonic*, *diuretic*, *diaphoretic*, and *alterative*, and is *used* as a substitute for sarsaparilla. The only preparation is the *fluid extract* (*extractum menispermæ fluidum*), dose \mathfrak{m}_{v-xxx} (0.308–2.0).

What is calendula?

Calendula is the *fresh flowering herb* of *C. officinalis* or marigold (*Nat. Ord.* Compositæ). It contains *calendulin*, and is supposed to be *diuretic*, *diaphoretic* and *alterative*. It is not much used.

TINCTURA CALENDULÆ (*tincture of calendula*); dose \mathfrak{f}_{3ss-j} (2.0–4.0). It may be used externally (diluted) as an application to wounds and contusions.

What is sassafras?

Sassafras is the *bark* of the *root* of *S. variifolium* (*Nat. Ord.* Laurinææ), and contains a *volatile oil*. It is supposed to be a mild *alterative* and *diaphoretic*, and is *used* in combination with sarsaparilla. The *oil* is used as a flavoring ingredient and as a perfume.

OLEUM SASSAFRAS (*oil of sassafras*), dose gtt. $\mathfrak{ij-x}$ (0.06–0.308).

What is stillingia?

It is the *root* of *S. sylvatica* or Queen's Delight (*Nat. Ord.* Euphorbiacææ) and contains a *volatile oil* and a *resin*. It increases all the secretions and excretions, slightly stimulates the cardiac action and is believed to be alterative. It has been used in *secondary* and in *tertiary syphilis*, *scrofula* and *chronic rheumatism*.

EXTRACTUM STILLINGIÆ FLUIDUM (*fluid extract of stillingia*), dose f3ss (2.0).

What is dulcamara?

It is the *young branches* of *solanum dulcamara*, woody nightshade or bittersweet (*Nat. Ord.* Solanaceæ), and contains *solamine* and *dulcamarine*. It is feebly narcotic, diaphoretic and diuretic. In large doses it is a cardiac depressant. It sometimes causes nausea and vomiting. It is neither hypnotic nor anodyne. It is chiefly *used* as an alterative in chronic eczema, psoriasis, etc., and has been recommended in whooping-cough, bronchitis and asthma.

EXTRACTUM DULCAMARÆ FLUIDUM (*fluid extract of dulcamara*), dose f3ss-j (2.0-4.0).

What is lappa?

Lappa or Burdock is the *root* of *Arcium Lappa* and other species of *Arcium* (*Nat. Ord.* Compositæ). It contains a bitter extractive, perhaps a glucoside, tannin, resin, etc., and possesses tonic, laxative, and slight diuretic, diaphoretic, and alterant properties. It is *used* in chronic rheumatism, syphilis, and "scrofulous affections" and occasionally in chronic cutaneous diseases, as psoriasis and prurigo. Locally, it is used as an application to contusions.

EXTRACTUM LAPPÆ FLUIDUM (*fluid extract of lappa*), dose m_{xv}-f3j (0.924-4.0).

ORDER IV.—ANTIPYRETICS.

What are antipyretics, and how do they act?

Antipyretics are remedies which reduce temperature in fever, but in medicinal doses have little or no effect upon it in the normal condition. The temperature may be lowered (1) by *lessening the production of heat*, or (2) by *increasing the loss of heat*. Heat production may be diminished either (1) by *depressing nutrition* (*i. e.*, by retarding oxidation and cell-proliferation), or (2) by *depressing the circulation of the part* (as by local abstraction of blood, etc.), or of the system (cardiac sedatives); while the loss of heat may be increased (1) by *increased evaporation* from the surface (diaphoretics), (2) by *increasing heat radiation*, through dilatation of the cutaneous vessels (amyl nitrite, alcohol, etc.), or (3) by *direct withdrawal of heat* from the surface (cold applications, etc.). The antipyretic group of medicines act chiefly by their effects on nutrition.

Mention the principal antipyretics.

Antipyrine; acetanilid (antifebrin); phenacetin; phenacoll; quinine; salicin; salicylic acid and the salicylates; salol; salophen; guaiacol; resorcin; hydroquinone; pyrocatechin; chinoline; thalline and kairine.

ANTIPYRINE.**What is antipyrine?**

Antipyrine (not official) is an alkaloidal substance usually synthetically prepared, but also obtained from the destructive distillation of coal tar, and belonging chemically to the phenyl group of carbon compounds. According to the latest investigations, antipyrine is a derivative of pyzole, which is derived from pyrrole.

What are the physiological effects of antipyrine?

Locally applied it is *antiseptic*, *antifermentative*, and *anæsthetic*. *Internally* it exerts its influence principally on *febrile temperature* and the *nervous system*, as an antipyretic and analgesic. In *fevers*, when an antipyretic dose (gr. xx-xxx (1.0-2.0)) is taken, the temperature rapidly falls several degrees (due principally to its direct action upon the heat-regulating apparatus of the nervous system, also partly to interference with the oxygen-carrying function of the blood, thus diminishing heat-production, and partly to dilatation of the cutaneous vessels, thus regulating heat-distribution and increasing heat-radiation), and more or less profuse sweating occurs. After an interval of from 4 to 12 hours the fever returns, sometimes preceded by a chill (which is not as severe as that caused by many of the other antipyretics, as kairine) and sometimes (very rarely) by depression almost amounting to collapse. Given in the same doses to a *healthy man* it only causes tinnitus aurium, nausea, and a slight fall of temperature (about $\frac{1}{16}^{\circ}$); *large doses* (3j (4.0)) cause vomiting and giddiness.

Antipyrine is quickly absorbed, and probably interferes with nutrition by combining with the hæmoglobin of the red corpuscles, forming methæmoglobin (the face becomes livid and the lips bluish when large doses are taken). It does not permanently impair their function nor alter their form. Medium doses stimulate the heart and raise the arterial pressure (partly from spasm of the internal vessels); very large doses (in animals) directly paralyze the heart and lower

the blood-pressure. Small doses have little or no effect upon the respiration. Given in large doses it increases the rapidity of the respiration and in poisonous doses it causes death apparently by respiratory paralysis. This action appears to be centric. It is a cerebral sedative and analgesic, acting especially on the sensory nerves, and dilates the pupils. It is eliminated by the kidneys, and when the perspiration is slight increases the urine. It sometimes produces a cutaneous rash and (very rarely in therapeutic doses) induces vomiting. It is said to be hæmostatic.

What are the symptoms of poisoning by this drug?

When an overdose is taken there are often nausea and vomiting, a sense of fulness in the head, tinnitus aurium, relaxation of the muscular system and muscular weakness, cyanosis, cutaneous eruptions, sweating, rapid breathing, and feeble pulse. Occasionally the reflexes are absent. Consciousness is retained.

How should these symptoms be treated?

The treatment consists in keeping the patient in a recumbent position; maintaining the body-heat; administering alcoholic and diffusible stimulants; giving atropine hypodermically and, if the cyanosis is marked, inhalations of alcohol.

The profuse sweating which is sometimes seen is best combated by giving atropine or agaricine.

The cutaneous eruptions rapidly disappear when the drug is discontinued.

What are its therapeutic uses?

It is used as an *antipyretic* in all cases of *hyperpyrexia*, as in *typhoid fever*, in *sunstroke* with high temperature, in the *exanthemata*, *surgical fever*, *pneumonia*, *erysipelas*, and *influenza*. In *acute rheumatism* it not only lowers temperature but relieves the pain. It is not an antiperiodic, but may be used to reduce the temperature in *malarial fevers*.

As an *analgesic* it is much employed in *neuralgia*, the "lightning-pains" of *locomotor ataxia*, *muscular rheumatism*, *myalgia*, *dysmenorrhœa*, to allay the pains of the first stage of *labor*, and in *influenza*. In *acute coryza* it is useful as a spray to the nasal mucous membrane (4 per cent. solution), preceded by the local application of cocaine.

What is the dose of antipyrine, and how may it be administered?

The dose as an *antipyretic* is gr. xv-xxx (1.0-2.0), repeated as circumstances require; as an analgesic, gr. x-xv (0.648-1.0), repeated in from 1 to 4 hours, if necessary. It is given in capsules or in solution. As it is non-irritating, it may be used hypodermically.

It is incompatible with calomel and with sweet spirits of nitre, tannates, and solutions of chloral, mercury, arsenic, and carbolic acid.

ACETANILIDUM—ACETANILID.

What is acetanilid?

Acetanilid or *antifebrin* is a neutral chemical product, obtained by the action of glacial acetic acid and heat on aniline. It is an acetyl derivative of aniline and is known chemically as phenylacetamide (C_6H_5NH, C_2H_3O).

What are its physiological actions?

Like antipyrine, *acetanilid* is markedly *antipyretic* and *analgesic*.

78

As an antipyretic *in fevers* it acts more slowly than antipyrine, influencing the temperature in the same manner and being attended with the same symptoms and with more marked cyanosis. As the temperature declines the pulse becomes slower and stronger and the arterial tension is elevated. Gastric and intestinal disturbance and other unpleasant concomitants rarely result. Given to *healthy men* in ordinary doses it produces no apparent effect. In *decided doses* acetanilid lessens the conductivity of nerves, depresses the reflex centres, and acts as a hypnotic. In *poisonous amounts* it causes motor and sensory paralysis. Its effects are chiefly expended upon the sensory cord and nerves. Given to animals *in toxic doses*, beside cyanosis it causes general prostration, stupor, rapidly lowered temperature, with primary stimulation and afterward depression of the heart and respiration, coma, convulsions, and death. It is eliminated by the kidneys as para-amido-phenol sulphate.

What are its uses?

It is used as an *antipyretic* in all cases of high temperature, as *typhoid fever*, *acute rheumatism*, *scarlatina*, *phthisis*, *influenza*,

pneumonia, and *acute sepsis*. As an *anodyne* it is employed, also, in the same diseases in which antipyrine is indicated, and has been found useful in *epilepsy*, particularly in the lesser attacks (*petit mal*). It is also used *locally* as an *antiseptic*.

How may it be administered, and in what doses?

The dose of acetanilid is from gr. iv-x (0.25-0.64). It is quite insoluble in water, and is better given in capsules or in powder.

PHENACETIN.

What is phenacetin?

Phenacetin or *acetphenetidin* (not official) is an acetyl derivative of amido-phenol.

What are its physiological effects?

It is a nervous sedative, acting principally upon the sensory portions of the spinal cord.

In small doses it has no effect upon the circulatory system. Large doses produce a general cyanosis due to the formation in the blood of methæmoglobin.

It slightly depresses normal temperature, and reduces markedly that of fever, usually causing but slight diaphoresis. When administered in the evening its effects upon febrile temperature appear to be more marked than when given in the morning. After large doses, in addition to cyanosis, vomiting, incoördination, hurried breathing, and somnolence may occur.

What are its therapeutic uses and doses?

As an *antipyretic* in febrile conditions generally, as in *typhoid fever*, *pneumonia*, *pleurisy*, *tuberculosis*, *influenza*, *acute rheumatism*, etc.

As an *analgesic* in *migraine*, *headache* from any cause, *myalgia*, etc. It is better to administer this, as well as the other coal-tar products, cautiously in all cases of weak heart. Dose gr. iij-xx (0.2-1.29).

PHENOCOLL.

What is phenocoll?

Phenocoll (not official) or *amidophenacetin* is a coal-tar product prepared "by the action of glycocoll, which is amido-acetic acid, upon phenacetin."

It is a white crystalline powder and is soluble in 16 parts of water. It is incompatible with the alkalies.

It is said to reduce febrile temperature very rapidly by diminishing the heat-production without affecting the heat-radiation, and to have no unpleasant sequelæ, except occasionally profuse perspiration.

It should be used with care, however, in all cases of prostration, as it may then produce cyanosis and cardiac depression. It has been used as a hydrochloride for its *antipyretic* effect in various febrile conditions; as an *analgesic* in *acute rheumatism* and in *neuralgia*, and as a *sedative* in *whooping-cough*. Dose of the hydrochloride gr. v-xv (0.32-1.0).

QUININE.

Is quinine used as an antipyretic?

Quinine (*vid.* p. 00) was formerly much used for this purpose, but at the present time it has been largely superseded by other agents.

As an antipyretic it is given in doses of gr. x-xx (0.64-1.30) repeated as required.

SALICIN, SALICYLIC ACID, AND THE SALICYLATES.

What are these substances, and whence are they obtained?

Salicin is a neutral principle found in several species of *salix*. *Salicylic acid* is a derivative of salicin, is found also as a salt (of methyl) in oil of *gaultheria*, and has been prepared synthetically by combining pure carbolic acid with dry carbon dioxide by means of heat. The *salicylates* are the salts formed by the union of this acid with bases.

What are the physiological effects of these substances?

The effects of *salicin* resemble those of salicylic acid, but its action is slower and less powerful. It is *antipyretic*, *tonic*, *antifermentative*, *antiseptic*, and to a slight extent *antiperiodic*. It does not depress the heart to the same extent as does salicylic acid, and is much less toxic. It is eliminated by the urine (chiefly as salicylic and salicyluric acids) and by the perspiration, rendering both of these excretions alkaline.

Salicylic acid, when *locally* applied, is *antiseptic* and *antifermentative*. Administered *internally in full doses* it lowers the temperature

in fevers, causes profuse, sometimes exhausting sweats, and produces symptoms similar to those of *cinchonism*, such as tinnitus aurium, partial deafness, headache, etc. After very large doses these symptoms are all increased, and dyspnœa, with depression of the heart and nervous system and dilated pupils, is observed. If a *toxic dose* be given, delirium, irregular respiration, involuntary evacuations, stupor, and death from respiratory failure occur. In healthy subjects salicylic acid in full doses often causes nausea and vomiting, is a cardiac depressant, inhibits the movements of the leucocytes, quickens and deepens the respiratory movements (producing dyspnœa), is a diaphoretic, does not affect the normal temperature, and is eliminated chiefly by the urine, which, if large amounts have been taken, will be of a green color (due to an increase of indican). The urine frequently contains albumin.

The *salicylates* administered internally are less irritant to the stomach than is salicylic acid; in other respects their effects are identical.

What are the medicinal uses of these substances?

Salicin is not much employed. It has been used as an *antipyretic*, especially in rheumatism, for which purpose gr. iij-xv (0.2-1.0) may be given every few hours.

Salicylic acid has been given as an *antipyretic* in various fevers, but is now little used for this purpose except in *acute rheumatism* in robust subjects, where it not only lowers the temperature and moderates the pain, but materially aids in the elimination of uric acid, yet probably does not decrease the frequency of cardiac complications. It is also used successfully in *pleurisy with effusion*. It has also been used with success as an *anthelmintic*. *Externally* it is applied for the removal of corns and warts; to check sweating of the feet; in eczema; and as an *antiseptic* to wounds and ulcers.

The *salicylates* are used internally as *antipyretics* in *acute rheumatism* and as *antizymotics* to prevent *fermentation* of food in the stomach.

What are the preparations and doses of these remedies?

SALICINUM (*salicin*), dose gr. x-3j (0.648-4.0). In *acute rheumatism*, gr. iij-xv (0.2-1.0) may be given every few hours.

ACIDUM SALICYLICUM (*salicylic acid*), dose gr. x-3j (0.648-4.0).

In *acute rheumatism*, gr. x (0.648) may be administered every hour until six doses are taken. If more is necessary, decrease the frequency of the dose.

SODII SALICYLAS (*sodium salicylate*) (3j (0.648) contains of salicylic acid gr. 48 (3.12); dose gr. v–3j (0.324–4.0). It is more soluble and less apt to cause vomiting than the acid, and hence is preferred for internal use.

LITHII SALICYLAS (*lithium salicylate*) is used for the same purposes and in the same doses.

METHYL SALICYLAS (*methyl salicylate*—*artificial* (or synthetic) *oil of wintergreen*) is a product obtained by the distillation of salicylic acid, or its salts, and methyl alcohol in the presence of sulphuric acid. It is identical with volatile oil of betula and differs in no respect from the other salicylates either in physiological action or in therapeutic use. It is less irritating to the stomach than the oil of gaultheria. Dose m̄j–v (0.06–0.31).

Phenyl salicylate will be described under the head of “*salol*.”

Physostigmine salicylate will be spoken of in connection with physostigma.

What other official preparations are used for the salicylate which they contain?

OLEUM GAULTHERIÆ (*oil of gaultheria*), a volatile oil distilled from Gaultheria procumbens or wintergreen. It contains *methyl salicylate*, and acts similarly to salicylic acid upon the system. In large doses it is toxic. It is administered for the same purposes as are the salicylates in doses of m̄x–xv (0.61–0.92) in capsules, repeated as required.

OLEUM BETULÆ VOLATILE (*volatile oil of betula*—*oil of sweet birch*), a volatile oil obtained by distillation from the bark of Betula lenta (*Nat. Ord.* Betulacæ). It is identical with methyl salicylate and nearly identical with oil of gaultheria. Dose m̄j–v (0.06–0.31).

What should be done if, during the use of these medicines, tinnitus aurium or other symptoms of intoxication should occur?

The dose should immediately be decreased or the medicine discontinued.

SALOL.

What is salol?

Salol (*phenyl salicylate*) is the salicylic ether of phenol and occurs as a white crystalline, slightly aromatic, and nearly tasteless powder, insoluble in water, but soluble in alcohol.

What are its physiological effects?

Locally it is *antiseptic*. *Internally* it lowers febrile temperature and produces sweating. The circulation is practically uninfluenced; the respirations increase in frequency and become shallow, tinnitus aurium and other salicylic acid symptoms sometimes occur. It is somewhat analgesic. In the intestine the pancreatic juice converts it into salicylic and carbolic acids, which render antiseptic the remainder of the alimentary canal. It seems to make the bile more watery. It is eliminated chiefly in the urine, which after its use resists decomposition and becomes darker in color.

What are its uses and doses?

As an *antipyretic* it is used in *acute rheumatism*, in *typhoid* and other *fevers*; in *phthisis* and in *pneumonia*; as an *intestinal antiseptic* in *duodenal catarrh*, *catarrhal jaundice*, *typhoid fever*, and *cholera*; to *prevent decomposition of urine* in *cystitis* and *pyelitis*, and as an *analgesic* in *neuritis*, *neuralgia*, *nervous headache*, etc. It is also employed in *pharyngitis* and in *chronic sore throat* depending upon the uric acid diathesis. It may be given in any stage of *gonorrhœa* for its influence on the urine. It is also said to be useful in diabetes.

Locally it is used as an antiseptic in surgical dressings. Dose gr. v-3j (0.324-4.0) in pills, capsules, or in solution in alcohol or oil. It should be administered cautiously in disease of the kidneys, because of the carbolic acid which it contains.

SALOPHEN.

What is salophen?

Salophen (not official) is acetylpara-amidophenol salicylate. Like salol it is decomposed in the intestine by the pancreatic juice, being split into salicylic acid and acetylpara-amidophenol, which is harmless, thus possessing an advantage in point of safety over salol. It is

chiefly eliminated under these forms in the urine, a small quantity passing unchanged in the stools. It is an *antipyretic*, *intestinal antiseptic*, and *analgesic*. Its most marked antipyretic effect is seen in *acute rheumatism*. It has also been used as an *intestinal antiseptic* in *typhoid fever*, *cholera*, and *dysentery*, and as an *analgesic* in *sciatica*, *migraine*, and *neuralgia*. It appears to be non-toxic. Dose gr. x-xv (0.64-1.0) in compressed pill or capsule every 2 hours until gr. xc (6.0) are taken in 24 hours.

GUAIACOL—GUAIACOL CARBONATE.

What are guaiacol and guaiacol carbonate?

Guaiacol (not official) or methylpyrocatechin is a colorless volatile liquid, with a strong aromatic odor, prepared by the fractional distillation of beechwood creasote, of which it forms 60 to 90 per cent.

It is also prepared by the dry distillation of guaiac resin, and, synthetically by the action of methyl-sulphuric acid upon pyrocatechin. The latter product is the purest form of guaiacol.

It is slightly soluble in water (1 part in 85, Helbing), but more so in ether and alcohol.

Guaiacol carbonate (not official) is made "by decomposing guaiacol in a solution of soda with carbonyl chloride. It is a neutral, white, crystalline powder, containing 91 per cent. of chemically pure guaiacol in combination with 9 per cent. of carbonic acid." Insoluble in water and in cold alcohol, it dissolves readily in ether, in chloroform, and in boiling alcohol, sparingly in the fixed oils and in glycerin.

What are their effects?

When *guaiacol* is given internally or administered hypodermically to healthy persons it produces no effects beyond slight depression and slight diaphoresis. When given in suitable doses to febrile patients a gradual *reduction of the febrile temperature* takes place, the fever declining for from 2 to 4 hours. When the lowest point has been reached the temperature rises rather rapidly to its former height, half the time being occupied in the ascent which was required for the decline of the temperature. As the temperature falls the patient often complains of much depression, but in ordinary doses the effects upon the respiration and pulse are evanescent.

It is eliminated by the lungs and kidneys as guaiaco-sulphuric

ether, giving a dark color to the urine when a large dose has been taken.

There is but one case of poisoning on record. *When applied locally* it is *antipyretic* and *analgesic*. It rapidly lowers febrile temperature, the subsequent rise being, as when it is administered internally, much more rapid than the fall. There are also profuse sweating and feeble cardiac action approaching syncope.

The action is most pronounced when applied to the abdomen, not so marked when rubbed on the chest, and least marked when applied to the thigh (Da Costa), probably because of the relative absorptive powers of different parts of the body.

It appears to lower the temperature by direct action on the thermogenetic centres. Guaiacol is also an *antiseptic*.

The carbonate when given internally is antipyretic, and, being decomposed in the small intestine into pure guaiacol and carbonic acid, acts as a local antiseptic upon the alimentary canal.

For what purposes are they employed?

They are much used as substitutes for creasote in the treatment of *pulmonary tuberculosis*, acute or chronic, especially in the first and second stages of the disease. In suitable cases it lessens the fever, improves the appetite and digestion, and moderates the cough.

By some "it is believed that it forms compounds in the blood with the toxins produced by the presence of the bacillus tuberculosis in the organism, and that it aids in eliminating them from the system." It may be used internally, either by inhalation or by hypodermic injection, or it may be applied locally. The internal use of the carbonate is probably to be preferred in these cases.

In *typhoid fever* as an *antipyretic* when the cold bath cannot be employed or is contraindicated, and as an *intestinal antiseptic*, guaiacol is very useful, employed locally and given internally in the form of the carbonate.

It has also been used as an *antipyretic* in other acute infectious diseases accompanied with high temperature, as in *pneumonia*, *influenza*, *erysipelas*, etc., and has been used for its *diaphoretic* action in *pleurisy* with effusion, as a means of promoting absorption.

As an *analgesic* it has been successfully used in *sciatica* and in various *neuralgias* painted over the painful areas, and has also been

used locally in *arthritis deformans* as a pain-relieving agent. In the latter affection the author has not seen the beneficial effects which have been produced by this agent in the hands of other physicians.

In *acute rheumatism* with high fever its local application in selected cases is of value.

It has also been used locally in *follicular tonsillitis*, and while not curative, will relieve the pain and enable the patient to swallow more easily. In *gonorrhæal epididymitis* painted over the scrotum it is said to relieve the pain quickly.

It has also been recommended as a *local anæsthetic* in dental and minor surgical procedures.

It is used as an *inhalation* in *fetid bronchorrhœa* and in *gangrene of the lung*. Dose of guaiacol $\mathfrak{m}\text{ijj-v}$ (0.18–0.3) four times a day in capsule, pill, or in solution in oil or alcohol. Hypodermically gr. j–iv (0.06–0.2) in sterilized oil may be used. The smaller dose should always be administered at first and gradually increased.

Locally $\mathfrak{m}\text{v-xl}$ (0.3–2.0) may be employed. When used in this manner the skin should be thoroughly cleansed and the guaiacol applied with a camel's hair pencil and rubbed in until absorbed. The surface should then be covered with oiled silk or paraffin paper to prevent evaporation.

As an inhalation $\mathfrak{m}\text{v-x}$ (0.3–0.6) in a pint of hot water may be used, or an oxygen inhaler may be employed.

Guaiacol carbonate is preferable for internal administration, as it is free from the odor and taste of guaiacol, which are objectionable to many persons, and does not irritate the alimentary canal. It is used in *tuberculosis* and *typhoid fever*. Dose gr. v (0.3) t. i. d. in capsules increased gradually until gr. xlv (3.0) are taken in divided doses in 24 hours.

What other antipyretics are occasionally used?

Resorcin; hydroquinone; pyrocatechin; chinoline; thalline, and kairine, of which only the first is official.

Describe their effects and uses.

Resorcin (*resorcinum*), *hydroquinone*, and *pyrocatechin* are isomers, being respectively meta-, para-, and ortho-oxyphenol.

Locally they are antiseptic and antifermentative. *Internally* they reduce febrile temperature and cause profuse diaphoresis, followed

by a more or less well-marked chill and gradual return of the fever. Fulness of the head or headache, flushed face, and tinnitus aurium accompany these effects; the pulse and respiration are at first accelerated but soon depressed. Large doses given to animals cause tremor, epileptiform convulsions, weak and rapid pulse, and finally failure of respiration. They are eliminated by the urine.

Chinoline is antiseptic and antipyretic, acting very similarly to resorcin. It depresses the heart and respiration, and paralyzes the reflex centres.

Kairine is also antipyretic, often causing much depression and acting dangerously on the blood.

Thalline is a synthetically prepared alkaloid used in the form of sulphate or tartrate. It is antipyretic, but causes profuse diaphoresis with, sometimes, dangerous depression almost amounting to collapse.

These substances have been used in all cases of high temperature as antipyretics, but have been superseded by newer and less dangerous drugs.

Resorcin is used *locally* as an *antiseptic*, *analgesic*, and *caustic*. In purulent and offensive discharges from the *nose*, *ears*, or *vagina* an injection of the strength of gr. x-xx (0.64-1.29) to f $\overline{3}$ j (30.0) of water may be used. In *chronic otitis media* it may be applied diluted with 7 parts of boric acid. It has also been used locally to the larynx in *pertussis*, in *chronic aphonia*, and to stimulate *chronic ulcers of the larynx*. In the form of ointments varying in strength from 1-50 per cent. it has been used in *chronic cutaneous diseases*, especially those of bacterial origin.

Of *resorcin* the dose is gr. v-xxx (0.32-2.0) in solution; of *hydroquinone* and *pyrocatechin* gr. ij-x (0.13-0.64); of *chinoline* gr. vii-xv (0.32-1.0) in solution in oil; of *kairine* gr. vij-xx (0.32-1.3); and of *thalline* gr. ij-xv (0.13-1.0) in pills or capsules.

Class II.—Medicines acting on the body by their effects on the various organs.

(A) ON THE NERVOUS SYSTEM.

ORDER I.—HYPNOTICS.

What are hypnotics?

Hypnotics are remedies which are administered to induce sleep. Most of them are *narcotics*, or remedies which by a stupefying effect on the nerve centres, diminish our relationship with external nature; and many of them are also *anodynes*, or remedies which by lessening the excitability of the nerves and nerve centres, tend to relieve pain.

How do hypnotics act?

As *during sleep* the functional activity of the brain is depressed, while the nutritive function is active, and as it is necessary for the nutrition of an organ that the blood should pass through the capillaries in a slow, steady and continuous stream, removing the waste products and giving up substances capable of replacing the tissue destroyed, while *during functional activity* the blood is in larger amount, hurried through the vessels (which are dilated) in a tumultuous manner, and does not remove the waste as fast as it accumulates in the tissues, therefore a medicine to cause sleep must *depress the cerebral functions* or *regulate the blood supply* to the brain by either contracting its vessels or by determining more blood to other portions of the body.

Name the principal hypnotics.

Opium and its alkaloids; chloral; chloralamide; chloralose; the bromides; hyoscine; sulphonal; trional; tetranal; paraldehyd; urethan; cannabis Indica; humulus, and lactucarium.

OPIUM.

What is opium?

Opium is the *concrete milky exudation* of the unripe capsules of *Papaver somniferum* (*Nat. Ord.* Papaveraceæ), a native of Persia, but cultivated throughout the civilized world.

What are its principal chemical ingredients?

It contains the following *alkaloids*: Morphine, narcotine, codeine, narceine, paramorphine (thebaine), papaverine, pseudomorphine, laudamine, hydrocotarnine, cryptopine, porphyroxine, meconine and others.

These are in combination with meconic and thebolactic acids.

What are the chemical tests for the presence of opium?

With even very diluted *solutions of opium, ferric chloride or sulphate* produces a blood-red ferric meconate. With *morphine and its salts, ferric chloride or sulphate* gives rise to a deep blue color, while *concentrated nitric acid* strikes with them a rich orange-red, slowly fading to yellow.

What are the physiological actions of opium?

Opium is hypnotic, anodyne, antispasmodic and diaphoretic. It also depresses the circulation and respiration. A *moderate dose* at first slightly stimulates, then calms, and finally depresses the *cerebral* functions, consciousness finally being lost in sleep, which is often disturbed by dreams; the reflex centres are depressed, and if the dose be toxic, finally paralyzed, death occurring from paralysis of the respiratory centre. The pupil is contracted from stimulation of the 3d pair of nerves. It depresses the *cardiac* motor ganglia, the pulse becoming slower and fuller and the arterial tension elevated. *Respiration* is slowed. The *secretions* are all diminished, except that of the skin, which is increased. Peristalsis is retarded. *Elimination* takes place chiefly through the kidneys. Occasionally, intense itching of the skin, and sometimes cutaneous rashes occur. Headache, nausea and constipation commonly follow a medium dose of opium.

What are the symptoms and treatment of opium poisoning?

The *symptoms* of opium-poisoning may be divided into three stages:

The first stage—not always present, and when present usually of short duration—in which there is mental exultation or excitement, with some confusion of ideas; a sensation of heat and often itching of the nose and face; a dry skin; and sometimes nausea and vomiting. This quickly passes into

The second stage, in which there is overpowering drowsiness, pass-

ing into stupor; contraction of the pupils; dryness of the mouth and throat; sweating; full, regular respirations, diminishing in frequency as the poisoning proceeds; and congested face. This stage gradually passes into

The third stage, with deepening coma; minutely contracted pupils (pin-point pupils); very slow, stertorous breathing; abolition of reflexes; pale, cyanosed face; rapid, feeble circulation, and death from respiratory failure.

Before death occurs the pupils often become widely dilated.

Death has resulted, in an adult, from morphine gr. $\frac{1}{4}$, hypodermically (rare), and from opium gr. ivss (rare). Children are very susceptible to its action.

Treatment.—(1) Remove any poison present in the stomach by the stomach pump or emetics (zinc sulphate gr. xx-xxx, or cupric sulphate gr. v-x in a glass of tepid water, or apomorphine gr. $\frac{1}{8}$, hypodermically); (2) Catheterize the patient to prevent resorption of the opium from the urine; (3) Maintain respiration and circulation by hypodermics of atropine sulphate gr. $\frac{1}{64}$, repeated in half the amount at the end of 15 to 30 minutes if necessary, the guide to its administration being the respiration. Strychnine, caffeine or cocaine are also recommended. Faradization, cold effusions and artificial respiration are useful. Potassium permanganate has been, of recent years, strongly recommended as an antidote to opium-poisoning. While it is not probable that it can act on the opium after the latter has been absorbed, yet it is well to employ a solution of the permanganate in washing out the stomach.

Post-mortem examination shows no characteristic changes in the organs.

How does the action of its alkaloids differ from that of opium?

The action of *morphine* differs from opium in that the former is less constipating, less of a diaphoretic, and about four times more powerful in other respects. *Codeine* is uncertain as an anodyne and hypnotic, but appears to possess a sedative effect on the vagus, thus quieting cough and bronchial irritation. The results of experiments with the other alkaloids are so conflicting that no reliance can be placed in them.

In what diseases is opium used medicinally?

As an *anodyne* opium or morphine is used in almost all diseases of which pain is a prominent and persistent symptom, and is only contraindicated when acute inflammation of the brain is present.

As a *hypnotic* in *delirium tremens* (often combined with the bromides or chloral) and in *sleeplessness* of fevers and other acute diseases; to *allay irritation*, especially when accompanied with *painful spasm*, as in the various forms of *colic* or in *asthma* (morphine hypodermically); in *severe acute vomiting* (by suppositories); to *check morbid discharges* or *restrain peristalsis*, as in *cholera*, *diarrhœa*, *dysentery*, and *enteritis*. In *peritonitis* for the latter effect and as an *anodyne*, large doses are required, a tolerance to opium being established.

In *puerperal septicæmia*, to *support the system* and as an *anodyne* it is of great value. As a *motor depressant* in *cerebro-spinal fever* very large doses are tolerated; also in *uræmic convulsions* (dangerous if the renal structure is much diseased), *puerperal eclampsia*, *tetanus*, *hydrophobia* and other affections with increased excitability of the reflex centres. In the *early stages of various inflammations*, for its influence on the circulation and tendency to retard effusion and relieve pain, and as a *diaphoretic* (Dover's powder) in muscular rheumatism, the early stages of acute catarrhs, etc., it is much employed. In *diabetes* opium and codeine are most reliable drugs to decrease the amount of sugar in the urine. *Great caution* should be exercised in prescribing opium in chronic diseases, as the *opium-habit* is almost certain to result. *Locally* opium is applied in solution as anodyne and sedative.

What are the preparations and doses of opium and its alkaloids?

OPIUM PULVIS (*powdered opium*) contains not less than 13 nor more than 15 per cent. of morphine. It is used in making the other preparations.

OPIUM DEODORATUM (*deodorized opium*) contains 14 per cent. of morphine; dose gr. ss-ij (0.032-0.130).

PILULÆ OPII (*opium pills*), each pill contains of powdered opium gr. j (0.065).

EXTRACTUM OPII (*extract of opium*), dose gr. ss (0.032).

EMPLASTRUM OPII (*opium plaster*) contains the extract with Burgundy pitch and lead plaster.

TROCHISCI GLYCYRRHIZÆ ET OPII (*troches of glycyrrhiza and opium—Wistar's cough lozenges*); each lozenge contains of extract of opium gr. $\frac{1}{20}$ (0.003). Very useful in laryngeal and pharyngeal coughs.

PULVIS IPECACUANHÆ ET OPII (*powder of ipecac and opium—Dover's powder*) is a diaphoretic and anodyne; dose gr. x (0.648), containing of opium and ipecac each gr. j (0.06) with sugar of milk.

TINCTURA OPII (*tincture of opium—laudanum*), dose ℥xij or 25 drops (0.74).

TINCTURA OPII DEODORATI (*tincture of deodorized opium*), dose ℥xij (0.74).

TINCTURA OPII CAMPHORATA (*camphorated tincture of opium—paregoric*) contains opium, diluted alcohol, oil of anise, benzoic acid, glycerin, and camphor; dose f3j-iv (3.696–14.786). To a young child gtt. ij-xx (0.185–1.232), according to age.

TINCTURA IPECACUANILÆ ET OPII (*tincture of ipecac and opium*). This is Dover's powder in a fluid form. Dose ℥xij-xx (0.74–1.232).

ACETUM OPII (*vinegar of opium—black drop*), dose ℥xij (0.74).

VINUM OPII (*wine of opium—Sydenham's laudanum*), dose ℥xij (0.74).

MORPHINÆ SULPHAS (*morphine sulphate*); MORPHINÆ HYDROCHLORAS (*morphine hydrochlorate*); MORPHINÆ ACETAS (*morphine acetate*). The salts are preferred to the pure alkaloid because they are more soluble. Dose of either gr. $\frac{1}{8}$ – $\frac{1}{4}$ (0.010–0.016) is equal to opium gr. j (0.065). They are well adapted to hypodermic use.

PULVIS MORPHINÆ COMPOSITUS (*compound powder of morphine—Tully's powder*) contains of morphine sulphate 1 part, with 20 parts each of camphor, liquorice, and calcium carbonate; dose gr. v-xv (0.324–0.972).

TROCHISCI MORPHINÆ ET IPECACUANHÆ (*troches of morphine and ipecac*). Each lozenge contains of morphine sulphate gr. $\frac{1}{40}$, (0.0016) and of ipecac gr. $\frac{1}{12}$ (0.005).

Magendie's solution (not official) contains of morphine gr. xvj to the ounce. It is sometimes used hypodermically, but it is better to make a fresh solution of morphine for each injection, as bacteria

frequently develop in solutions of the alkaloids when kept for some time, and also because otherwise there is some risk of confusion, as the French Magendie's solution contains about gr. xijss to the ounce. Also the solutions of morphine official in the British Pharmacopœia do not correspond in strength to either of the above solutions.

CODEINA (*codeine*), dose gr. $\frac{1}{4}$ –ij or even gr. v (0.016–0.130–0.32).

What medicines are incompatible with opium ?

The alkalies precipitate morphine from solutions of opium and vegetable infusions containing tannic acid form a less soluble tannate of morphine ; many mineral salts are also decomposed by solutions of opium.

How is opium administered ?

Opium is often administered in the form of the officinal preparations, either alone or combined with other substances. The bromides, chloral or other hypnotics are often added to increase its soporific influence ; belladonna, etc., to enhance its anodyne effects ; lead or other astringent to act more powerfully in intestinal disorders ; potassium nitrate, or minute doses of antimony, to increase, while atropine is often combined with morphine to diminish, its action as a diaphoretic. It may be administered by the mouth or rectum (suppositories or laudanum enemata), and its alkaloids may also be given hypodermically.

CHLORAL.

What is chloral ?

Anhydrous chloral is an oily liquid made by saturating absolute alcohol with dry chlorine gas, and is *trichloraldehyd*. By uniting with water, a hydrate, the *official chloral*, is formed, which occurs as a crystalline solid.

What are its physiological effects ?

Locally applied, chloral is antiseptic and irritant. Given *internally*, it is a hypnotic, motor depressant and cardiac sedative. In *medicinal doses*, it causes quiet, natural sleep, contraction of the pupil and deep, regular breathing. After *large doses*, the sleep is deeper and may even pass into coma, respiration is slowed, the cardiac action is weakened and diminished, the temperature falls, and sensibility and reflex action are depressed. It is not an anodyne.

Nausea and vomiting sometimes occur from its irritant action on the mucous membrane, and in some persons excitement and even delirium may take the place of the sleep which it usually produces. The *continued use* of chloral causes *chloralism* or the *chloral habit*, and produces an anæmic condition, with weak, irregular heart, irritable temper and loss of self control: a state similar to delirium tremens has been observed.

What are the symptoms and treatment of chloral poisoning?

After toxic doses, profound narcosis, great muscular relaxation, slow and very weak pulse, diminished frequency of breathing, marked contraction of the pupil and lividity of surface, with great reduction of body heat, occur. Death is due to paralysis of respiration or to sudden failure of the heart from paralysis of its motor ganglia.

The smallest recorded fatal dose is gr. xxx (2.0).

The *treatment* for chloral poisoning consists in (1) maintaining the temperature by artificial heat; (2) maintaining the cardiac action by *atropine* in small doses, repeated as required, and (3) maintaining the respiration (partly by the atropine or by morphine, and partly by artificial manipulations). While chloral is the best antidote for strychnine poisoning, the converse does not hold good.

What are its therapeutical uses?

As a *hypnotic*, chloral is employed in *insomnia*, *hysterical* or *maniacal excitement*, and in *delirium tremens* and *mania-a-potu*, but is *contraindicated* in any condition where the heart is weak or has undergone fatty degeneration. As a *motor depressant* in *uræmic*, *infantile* or *puerperal convulsions*, in tetanus, and in poisoning by *strychnine*, it is of great service, and has been used in *whooping-cough* and *chorea*. It is also recommended in sea-sickness (gr. v t. d.) and to relax a rigid cervix during the first stage of labor. *Locally*, in diluted solution, it has been used as an *anodyne* and *antiseptic* dressing, and, triturated with equal parts of camphor, it forms a valuable liniment in neuralgic and muscular pains.

How is chloral administered?

The dose of chloral is gr. ij-xx (0.130-1.296), which may be repeated in an hour or two if sleep is not produced. It is better given in syrup. It should not be given with alkalis, which decompose it, chloroform and formic acid being formed.

What is croton-chloral hydrate?

Croton-chloral hydrate (not official), or butyl-chloral hydrate, is made by the action of chlorine gas on ethylic aldehyd, and occurs in small, glittering crystals. Its *effects* and *uses* are similar to those of chloral, and as it possesses a very decided anæsthetic influence over the filaments of the trigemini, it is much used in the neuralgiæ affecting the branches of these nerves. Dose gr. v-xx (0.324-1.296), in glycerin or syrup.

CHLORALAMIDE.

What is chloralamide?

Chloralamide (not official) is a compound prepared by the action of formamide upon chloral. It is soluble in water, but more so in alcohol. Its *action* is similar to that of chloral, but it is not so certain a hypnotic, and neither depresses the heart nor irritates the stomach and kidneys to the same extent as does the latter drug. It is excreted as urochloralic acid.

Occasionally, after the exhibition of large doses, it causes *slight poisoning*, the symptoms being chilliness followed by a sensation of heat, sweating, arrhythmia, nausea, and sometimes vomiting, with respiratory depression. These symptoms are rarely severe and usually pass off rapidly.

It is used as a *hypnotic* and *nervous sedative* in the same conditions for which chloral is prescribed, and is especially useful in *insomnia* due to nervous causes, but fails usually when *insomnia* is due to pain. It is also recommended in sea-sickness.

Dose gr. x-xxx (0.65-2.0) given in capsule or solution one hour before its effects are desired. It may be administered by the rectum.

CHLORALOSE.

What is chloralose?

Chloralose (not official) is formed by heating a mixture of anhydrous chloral and glucose.

It occurs as fine white crystals having an unpleasant, rather sharp taste, and being readily soluble in hot water and alcohol, but only slightly soluble in cold water.

While it appears to be a reliable hypnotic, it induces toxic symptoms readily, apparently by increasing the reflex excitability of the

medulla oblongata, and would at present seem rather difficult to handle, as the toxic dose varies with different individuals, and is smaller in nervous and hysterical patients and in alcoholic subjects.

In sufficient doses general convulsions are produced, frequently accompanied by involuntary urination.

The toxic symptoms rapidly disappear on discontinuing the drug and appear to leave no serious consequences behind them. It has been *used* as a *hypnotic* and *analgesic*.

Dose gr. ij-ijj or iv (0.13-0.2-0.25), always beginning with the smaller dose.

THE BROMIDES.

What are the physiological effects of the bromides?

The *bromides* are hypnotic and depressors of the reflex spinal centres. Applied *locally* to the pharynx, they diminish its reflex irritability. Given *internally* they cause drowsiness and often sleep by inducing a condition of cerebral anæmia, depress the reflex spinal centres and the functions of the end-organs of peripheral nerves, thus lessening nervous irritability, reflex activity and cutaneous sensibility. They also depress the functions of the sympathetic nervous system. In very large doses *potassium bromide* lessens the cardiac frequency and force, lowers the arterial tension (while contracting the blood vessels), and finally in animals paralyzes the heart in diastole. This seems to be due to the potassium only (which is a cardiac and nerve poison), for the other bromides do not arrest the cardiac action. They slow the respiration and lower the temperature of the body. They are eliminated by the kidneys, skin and mucous membrane.

What is bromism?

When taken for a long time they induce a condition known as *bromism*, the symptoms of which are acne, anæmia, drowsiness, defective memory, malaise, impaired reflexes, malnutrition, loss of sexual power and desire, and often a tendency to melancholia. These symptoms pass away when the drug is discontinued. No case of *acute poisoning* is on record.

What are the medicinal uses of the bromides?

They are used as *hypnotics* in all forms of *insomnia due to cerebral hyperæmia*, but are contraindicated when anæmia is present. In

the early stages of *delirium tremens* they are of great service. They are also given as *depressor motors* in local or general convulsions as *chorea*, *uræmic*, *puerperal*, *hysterical* or *infantile convulsions*, *tetanus*, and as an *antidote* to *strychnine poisoning* (inferior to chloral). In the treatment of *epilepsy* they rank among the best remedies. In reflex vomiting, as of sea-sickness or pregnancy, they are of use. In *whooping cough* ammonium bromide is often efficacious. In *migraine* they are sometimes beneficial. For their *influence over the sexual functions* they are used to control *nymphomania*, *priapism*, and *masturbation*. Conjoined with ergot they are used to decrease the uterine blood supply in *menorrhagia*, and are of utility in *subinvolution of the uterus*. Strontium bromide is said markedly to diminish the amount of sugar in the urine in *glycosuria*.

What are the official bromides?

POTASSII BROMIDUM (*potassium bromide*), dose gr. v-3j (0.32-4.0).

AMMONII BROMIDUM (*ammonium bromide*) has less influence on the circulation, respiration, and temperature than the potassium salt; dose gr. v-xxx (0.32-2.0).

SODII BROMIDUM (*sodium bromide*), dose gr. v-3j (0.32-4.0).

CALCI BROMIDUM (*calcium bromide*), dose gr. v-3j (0.32-4.0).

ZINCI BROMIDUM (*zinc bromide*), dose gr. j-x (0.065-0.648) (not much used).

STRONTII BROMIDUM (*strontium bromide*), dose gr. iij-x (0.2-0.6).

ACIDUM HYDROBROMICUM DILUTUM (*diluted hydrobromic acid*), containing 10 per cent. of the absolute acid, is said to possess similar effects to the bromides. It is chiefly used as a hypnotic, and added to quinine mixtures to prevent cinchonism. Dose f3ss-j (2.0-4.0).

What medicines are incompatible with the bromides?

Acids and the acidulous and metallic salts are chemically incompatible with the bromides. They are antagonized by cardiac and vaso-motor stimulants.

How may they be administered?

They may be given in powder or in solution in syrup. Zinc bromide may be used in pilular form. They are often given together in small doses, and to enhance their hypnotic effect may be combined with opium or its alkaloids, chloral or cannabis indica.

HYOSCINÆ HYDROBROMAS.

What is hyoscine hydrobromate?

Hyoscine is one of the alkaloids found in *Hyoscyamus niger*, or henbane (*q. v.*). It is an efficient hypnotic when used hypodermically, and is employed in the form of *hydrobromate* in *acute mania*, *alcoholic delirium*, and *hysteria*. Dose gr. $\frac{1}{80}$ (0.0008) by the mouth, or hypodermically gr. $\frac{1}{120}$ (0.0005).

PARALDEHYDUM.

What is paraldehyd?

Paraldehyd is a colorless liquid with a disagreeable odor and taste, prepared by heating ethyl aldehyd with a mineral acid or with zinc chloride.

What are its physiological actions?

In medicinal doses paraldehyd produces sleep closely simulating natural sleep, without affecting the cardiac action. It depresses the reflex centres, especially those of the upper part of the cord, and in *toxic amounts* depresses and finally paralyzes the respiratory centre, diminishing the amount of carbonic acid exhaled, and lowering the body temperature. It has no anodyne virtues. It often irritates the mucous membranes, and cutaneous eruptions sometimes follow its use.

What are its medicinal uses and dose?

It may be given as a *hypnotic*, where sleeplessness is not induced by pain, as in *delirium tremens*, *hysteria*, etc. As a *reflex depressor* it has been employed in *epilepsy*, and in animals poisoned by strychnine has proved antidotal. It is given in doses of f3j (4.0), and should be freely diluted.

SULPHONAL.

What is sulphonal?

Sulphonal or *dional* (not official) is a substance prepared synthetically and described chemically as diethylsulphon-dimethylmethane. It occurs as heavy, prismatic crystals, colorless, odorless, and nearly tasteless, and is soluble in 100 parts of cold water, but freely soluble in hot water and also in ether and alcohol.

What are its physiological and its toxic effects?

The physiological action of sulphonal is not thoroughly understood. It acts chiefly on the cerebral centres, depressing them and

causing sleep. Sometimes, especially after large doses have been taken, it depresses the reflexes, disorders digestion, and causes cutaneous eruptions. In large doses it depresses respiration, rendering it slow, weak, and, when excessive doses have been taken, stertorous, and finally producing death by cessation of the respiratory act.

It has but little effect on the circulation, and experimenters differ as to its effect upon the blood, the majority believing that it causes no change in that fluid which can be detected by chemical or spectroscopic analysis.

It is probably eliminated in great part by the urine, and frequently changes the color of that excretion to a reddish-brown—a change supposed to be due to the presence of hæmatoporphyrin, although the coloring-matter differs from the latter on spectroscopic analysis.

Disagreeable after-effects sometimes follow the use of sulphonal, as drowsiness, giddiness, headache, and occasionally incoördination of gait, sometimes cutaneous eruptions or disordered digestion, perhaps nausea and vomiting, occur.

Sulphonal is by no means an absolutely innocuous drug, and *death* has resulted from large doses in many recorded cases, from respiratory failure.

The symptoms of poisoning are various, the most common being drowsiness, stupor, muscular incoördination, mental and physical hebetude, depression of reflexes, nausea and vomiting, and usually constipation. Sometimes ataxic nervous symptoms, “diplopia, muscular tremor or paresis, ptosis, œdema of the eyelids, slow and weak (possibly stertorous) respiration, slow pulse, elevation of temperature, general anæsthesia, urine changed in color to reddish-brown, diminished in quantity or suppressed, aphasia, and cyanosis” (M. L. Foster).

In several cases with recovery which the author has seen the symptoms were drowsiness almost amounting to stupor; mental hebetude; slow speech; slight hallucinations; slow, shallow respiration; slow, weak pulse, and staggering gait with slight cyanosis. In non-fatal cases recovery is usually rapid. Sulphonal is *cumulative* in its action. During its administration it is of the utmost importance that the bowels should be made to act daily (Schulz) and that the urinary secretion should be abundant.

If symptoms of poisoning appear, the drug should at once be discontinued; the bowels should be freely opened, saline diuretics administered, and the patient kept in the recumbent position. If there is much depression of the circulation or respiration, stimulants may be administered.

What are its therapeutic uses?

Sulphonal is much used as a *hypnotic* and *cerebral sedative* in *nervous insomnia*, and even in *melancholia*, *delirium tremens*, and *acute mania*. In the *insomnia* of low fevers, as that of *typhoid fever*, it sometimes gives good results.

It has been recommended as a *sedative reflex depressant* in *asthma*, *hiccough*, *trismus neonatorum*, and in *convulsions due to dentition*, and has even been used in *epilepsy* and *chorea*. It is said that in small doses (gr. viij (0.5)) it is effective in arresting the *night-sweats* of *phthisis*, and that in *diabetes* it will temporarily diminish the amount of sugar excreted in the urine.

It is said to be *contraindicated* in *angina pectoris*. Dose gr. v-xxx (0.3-2.0), which may be given in capsules, but is better administered dissolved in hot water or milk, two hours at least before the time when its effects are desired.

As it is cumulative in its action, it is better not to give more than gr. xxx (2.0) in 24 hours.

TRIONAL.

What is trional?

Trional (not official) is a sulphone differing from sulphonal in that it contains one atom more of carbon and one less of hydrogen than the latter. Its chemical name is diethylsulphon-methylethylmethane.

It occurs as white, odorless, almost tasteless crystals, not very soluble in cold, but readily soluble in boiling water, and also in alcohol and milk.

What are its effects and uses?

Trional is a hypnotic, acting more quickly than sulphonal, being easily decomposed in the body, sometimes slightly lowering the pulse and blood-pressure, and rarely leaving any disagreeable after-effects. It is not cumulative. Occasionally after its long-continued use the urine is colored reddish-brown from the presence of hæmaturia.

Toxic doses given to animals cause death from respiratory paralysis.

Occasional *after-effects* are loss of appetite, epigastric pain, nausea and vomiting, vertigo, and unsteadiness of gait. These symptoms usually soon disappear.

It is supposed to act directly on the cerebral cortex.

Several cases of *acute poisoning* by trional have been reported; the symptoms being drowsiness, staggering gait, vertigo, vomiting, diarrhoea, loss of muscular power, and lowered temperature, sometimes attended with rapid pulse and respiration, excitement, and collapse.

Trional is *used* as a *hypnotic* in *insomnia* due to organic cerebral affections, neurasthenia, or other functional psychosis, worry or mental over-work.

As a sedative in cerebral excitement it is decidedly inferior to tetronal, and in sleeplessness due to pain it usually fails entirely.

It is a "safe and efficient hypnotic for children" when pain is not present (Claus).

It has also been recommended in *epilepsy*, either alone or combined with the bromides.

Dose xv-xxx (1.0-2.0) about half an hour before its effect is desired.

If it is administered for any length of time, it is better to intermit the dose occasionally, in order to guard against any possible cumulative effect, and it is important also to insure a daily evacuation of the bowels.

TETRONAL.

What is tetronal?

Tetronal (not official) belongs chemically to the disulphone group, to which also sulphonal and trional belong. It is diethylsulphon-diethylmethane. These three products differ from each other chemically only in the number of molecules of ethyl and methyl which they contain. Thus *sulphonal* contains *two* molecules each of ethyl and of methyl, and is sometimes spoken of as dional; *trional* contains *three* molecules of ethyl and one of methyl, and *tetronal* contains *four* molecules of ethyl and none of methyl.

Tetronal is formed synthetically, and occurs in colorless, shining plates, without odor or taste, soluble in about 500 parts of cold, more soluble in hot water; soluble also in alcohol and in ether.

What are its effects and uses?

Its *effects* and *uses* are similar to those of trional, but it has a more decided sedative effect. Occasionally its administration is followed by disagreeable after-effects, as headache, anorexia, nausea, vomiting, vertigo, and even incoördination.

It is not decomposed in the system, and is excreted under its own form in the urine.

Dose gr. v-xv (0.3-1.0), repeated in an hour if necessary. The best vehicle for its administration is hot milk.

When given for a length of time it is better to intermit its use every 3d or 4th night.

URETHAN.

What are the effects and uses of urethan?

Urethan (not official) or ethyl carbamate is a synthetical product formed by the interaction of ethyl carbonate and ammonia, or by that of urea nitrate and ethyl alcohol, and occurs in crystalline masses readily soluble in water and alcohol. *In medicinal doses* it causes tranquil sleep, preceded by a short period of excitement. It depresses the reflex centres, at first quickens but soon slows the respiration, and somewhat reduces the temperature, and in overdoses causes, in animals, death from respiratory paralysis. The effect upon the heart is disputed. It is not analgesic. It has been *used* as a *hypnotic* in simple insomnia with varied success, and appears to be more applicable to cases where the action of chloral on the heart contraindicates the use of that agent. As a *motor depressant* it has been employed in *tetanus*, *infantile convulsions*, and *puerperal eclampsia*. Dose gr. ij-xxx (0.130-2.0). It has been used locally as an antiseptic dressing for wounds.

CANNABIS INDICA—INDIAN CANNABIS.

What is this drug?

Cannabis Indica is the *flowering tops* of the female plant of *C. sativa* or hemp (*Nat. Ord.* Urticaceæ), grown in the East Indies.

What are the active chemical constituents of hemp?

The alcoholic extract contains a *resin* (cannabin) and a volatile oil. It is claimed that several alkaloids have been discovered, but their existence is, as yet, doubtful.

What are the physiological actions of cannabis?

In full *medicinal doses* it at first stimulates and then depresses the cerebro-spinal functions without affecting respiration or circulation to any marked extent. The mental exhilaration is usually of a pleasurable kind, the pupils are dilated, ideas flow rapidly and conception of time and space are so altered that seconds seem hours and objects a few feet off seem miles away; an active, busy delirium usually occurs and sometimes increase of the sexual appetite; this is followed by a sense of weight in the extremities, depression of the reflexes, some loss of muscular power and cutaneous anæsthesia, succeeded by sleep. It increases but does not originate uterine contractions. No fatal case from its use is recorded.

What are its therapeutic uses?

It is used as a *hypnotic* in *mania* and in *mania-a-potu*; as an *anodyne* in *neuralgia*, *migraine*, *dysmenorrhœa*, *rheumatism*, and *gout*, and to relieve pain in *acute* and *chronic Bright's disease*; as an *antispasmodic* in *tetanus*, *chorea*, *hysteria*, and *spasm* of the neck of the bladder; as an *addition to cough medicines* to allay irritation of the throat, and to promote a condition of euthanasia in the later stages of *phthisis*. It is highly recommended in *dysentery*, and is of undoubted service in checking *uterine hemorrhages* not due to abortion or following labor. In *sexual impotence*, when functional, it is sometimes of use. In *gonorrhœa* it is said to lessen the discharge and prevent *chordee*. As a *local application* it enters into the composition of many *corn-cures*.

What are its preparations and doses?

EXTRACTUM CANNABIS INDICÆ (*extract of cannabis Indica*), dose gr. $\frac{1}{4}$ -j (0.016-0.065).

EXTRACTUM CANNABIS INDICÆ FLUIDUM (*fluid extract of cannabis Indica*), dose m̄j-x (0.06-0.6).

TINCTURA CANNABIS INDICÆ (*tincture of cannabis Indica*), dose m̄xxx (2.0). In using a new sample of *cannabis Indica* always begin with the minimum dose, as different samples vary much in strength.

HUMULUS.

What are the effects, uses, preparations, and doses of humulus?

Humulus is the *strobiles* of *H. lupulus* or hop vine (*Nat. Ord.* Urticaceæ). *Lupulin* is a yellow powder found near the base of the strobiles. It contains a volatile oil, *choline* (formerly called lupuline, a strongly alkaline liquid), *lupamaric acid* (a bitter principle), resin, etc.

Hops are tonic, very feebly narcotic, anaphrodisiac, and diuretic. They are used as a *hypnotic* in mild forms of *delirium tremens*, and in *restlessness* and *sleeplessness* from overwork; as an *anaphrodisiac* in *priapism*, *chordee*, *nymphomania*, and *satyriasis*; as an *anti-spasmodic* and *sedative* in *hysteria* and *neurasthenia*; as a *diuretic* and *anodyne* in *irritation of the bladder*, and locally as an anodyne poultice, and as a sedative injection in acute vaginitis.

Hop pillows are a popular remedy for sleeplessness due to nervousness.

TINCTURA HUMULI (*tincture of humulus*), dose fʒj-iv (4.0-15.0) or more.

LUPULINUM (*lupulin*), dose gr. v-x (0.324-0.648), but not much used in this form.

EXTRACTUM LUPULINI FLUIDUM (*fluid extract of lupulin*), dose fʒss-ij (2.0-8.0).

OLEORESINA LUPULINI (*oleoresin of lupulin*), dose m℥x-xxx (0.61-2.0).

LACTUCARIUM.

What are the effects, uses, and preparations of lactucarium?

Lactucarium is the *concrete milk-juice* of *Lactuca verosa*, the garden lettuce (*Nat. Ord.* Compositæ), and contains *lactucin*, which is said to be even more feebly hypnotic than the crude drug. *Lactucarium* is supposed to possess *anodyne* and *hypnotic* powers, and has been used as a substitute for opium when the latter is indicated, but cannot, for any reason, be given. Its action, if any, is very mild.

TINCTURA LACTUCARII (*tincture of lactucarium*), dose m℥x-fʒj (0.6-3.6).

SYRUPUS LACTUCARII (*syrup of lactucarium*), dose fʒij-iv (8.0-15.0).

ORDER II.—MYDRIATIC ANODYNES.

What are the mydriatic anodynes ?

The *mydriatic anodynes* are remedies employed as analgesics and antispasmodics, which possess no hypnotic effect, but cause dilatation of the pupil and, in large doses, a restless delirium.

What medicines are included under this head ?

Belladonna, stramonium, hyoseyamus, duboisia, coca, and their alkaloids, and antipyrine.

BELLADONNA.

What is belladonna ?

Belladonna is the leaves and root of *Atropa Belladonna* or deadly nightshade (*Nat. Ord. Solanaceæ*), a European plant, cultivated in this country.

What are the chemical constituents of belladonna ?

Its properties depend upon an alkaloid called *atropine*, which exists in the plant as a malate. Both the leaves and the root also contain *belladonnine* (probably oxyatropine), *hyoscyamine* (sometimes the predominating alkaloid in the leaves and often the only alkaloid in young roots), and *atropamine*.

What are the tests for atropine ?

A delicate test is the addition of hydrobromic acid, which produces a yellow, amorphous precipitate, soon becoming crystalline. The *physiological test* (viz., the dilatation of the pupil in the lower animals after the local application of the suspected solution) should always be resorted to.

What are the physiological effects of belladonna and its alkaloid ?

Applied *locally* to the skin, it is absorbed, lessens the sensibility of the sensory nerves, and produces constitutional effects. *Small doses, administered internally*, cause dryness of the mouth and throat, with some headache ; after *larger amounts*, there is difficulty in swallowing, from arrest of the mucous and salivary secretions, the pupils dilate widely, the skin becomes dry and a scarlatinoid rash appears, the cardiac action is accelerated, and an active delirium occurs. In *moderate doses*, they act as *cerebral excitants*; *larger*

doses produce hallucinations and impair the excitability of the motor and sensory nerves, thus producing a peculiar delirium in which there is a great desire for constant motion associated with lassitude. The *spinal reflex centres* are first stimulated, afterward paralyzed, if the amount taken be sufficiently large. It paralyzes the 3d nerves and stimulates the end organs of the sympathetic, thus dilating the *pupil* whether used locally or internally; the power of accommodation is also paralyzed, and after large doses the intraocular tension is lessened.

The *cardiac action* is at first slowed, if the dose be small (from stimulation of the inhibitory centre), but is soon much quickened (from progressive paralysis of the peripheral filaments of the vagus and stimulation of the cardio-accelerator centres or nerves). Large doses paralyze the cardiac muscle. Atropine stimulates the vaso-motor centres and raises the arterial pressure, but after toxic doses the arterioles dilate, from paralysis of their muscular walls, and the blood pressure falls. It first stimulates and then paralyzes the respiratory centre. The *temperature* rises after moderate doses, probably from stimulation of the thermo-genetic spinal centres; but after toxic amounts have been taken a decided fall of temperature occurs, probably from vaso-motor paralysis. Small doses probably increase *intestinal peristalsis*, by depressing the inhibitory fibres of the splanchnics, while large doses paralyze the non-striated muscular fibres of the intestines and bladder. It arrests the *secretions* of the skin, mammary, salivary and muciparous glands, and probably of the liver and pancreas, by paralyzing the terminal filaments of the secretory nerves. The intestinal secretions are probably increased, as is that of the kidneys (usually). After large doses, the urine may be entirely suppressed.

Atropine is eliminated principally by the kidneys.

What are the symptoms and treatment of atropine poisoning?

The pupils dilate so widely that the iris can scarcely be seen; swallowing is impossible, from dryness of the throat and mouth; the muscular system is relaxed; maniacal delirium occurs, with impaired sensation, failing pulse, quickened followed by failing respiration, stupor, coma, sometimes convulsions, diminished body heat, and death usually from asphyxia. The *treatment* consists of: (1)

evacuation of the contents of the stomach (by emetics or stomach pump), *bowels* (by cathartics) and *bladder* (by the catheter), to prevent further absorption of the poison; (2) *giving tannic acid* freely as a chemical antidote; (3) *giving pilocarpine, physostigmine or morphine* hypodermically, as partial physiological antagonists, and (4) maintaining the circulation and respiration by electricity and artificial respiration.

What are the medicinal uses of belladonna and atropine?

As an *anodyne* in the various *neuralgic*, either locally or given by the mouth, rectum or subcutaneously, it is of value, and is often combined with morphine. In *muscular rheumatism* and in *myalgia* it is also of service. In *conjunctivitis* and in *iritis* it is used locally, not only to dilate the pupil but to lessen pain.

As an *antispasmodic* it is used to relax spasm and allay pain in *lead colic*, *spasmodic dysmenorrhœa*, and *cramps* of the bowel, bladder and urethra; also to relax the *rigid cervix uteri* in labor. In *asthma*, in *laryngismus stridulus* and *whooping cough* it often proves of use. In *habitual constipation*, a small amount combined with purgatives increases their action by relaxing spasm of the intestinal muscular layer and stimulating peristalsis. It is a valuable remedy in the treatment of *incontinence of urine in children* when given in doses sufficient to produce some constitutional effect.

As a *cardiac and vaso-motor stimulant* in all cases of threatened collapse from *heart failure* it is advantageously employed, and has been highly recommended to contract the blood-vessels in *acute inflammatory affections* generally. To check *excessive secretion*, as in *salivation*, *excessive sweating*, especially the *night-sweats of phthisis*, and in *colliquative diarrhœa*, it is of great value, and is a well-known *galactafuge*, being used internally and applied locally for this purpose. As a *respiratory stimulant* in *opium poisoning* atropine is of great service, and may be used in cases of poisoning by other substances attended with failure of respiration.

Locally, belladonna is used as an ointment, liniment or plaster for the relief of pain in *neuralgia*, *myalgia* and *muscular rheumatism*; to *allay cardiac palpitation*; to *relax spasmodic stricture* of the urethra (applied on a bougie or given by suppository), and as a *galactafuge*. In *ophthalmic practice* a solution of atropine (gr. $\frac{1}{20}$ -iv

(0.003–0.25) to water f3j (4.0)) may be dropped into the conjunctival sac to *dilate the pupil* and to *paralyze accommodation*, so that an examination can be readily performed or the refraction of the eye determined. In various inflammations of the ocular apparatus, as conjunctivitis, keratitis, iritis, etc., it is used to lessen pain and diminish the blood-supply to the inflamed parts.

What are the preparations and doses of belladonna?

EXTRACTUM BELLADONNÆ FOLIORUM ALCOHOLICUM (alcoholic extract of belladonna leaves), dose gr. $\frac{1}{8}$ –ss (0.008–0.032) in pill or suppository.

EXTRACTUM BELLADONNÆ RADICIS FLUIDUM (fluid extract of belladonna root), dose ℥j–ij (0.061–0.12).

TINCTURA BELLADONNÆ (*tincture of belladonna*), dose ℥v–xxx (0.3–2.0).

LINIMENTUM BELLADONNÆ (*liniment of belladonna*) contains 95 per cent. of the fluid extract and 5 per cent. of camphor.

UNGUENTUM BELLADONNÆ (*ointment of belladonna*) contains 10 per cent. of fluid extract.

EMPLASTRUM BELLADONNÆ (*plaster of belladonna*).

ATROPINA (*atropine*), usually employed in the more soluble form of ATROPINÆ SULPHAS (*atropine sulphate*), dose gr. $\frac{1}{120}$ – $\frac{1}{60}$ (0.0005–0.001).

What substances are incompatible with belladonna?

Alkalies precipitate atropine from solutions of belladonna, and tannic acid forms with it atropine tannate, a less soluble salt.

What is homatropine hydrobromate?

It is a non-official salt of an alkaloid derived artificially from atropine. *It differs from atropine* in dilating the pupil more rapidly and much more transiently, in not causing much constitutional disturbance when used locally for this purpose, and in slowing, not accelerating, the heart's action. *It is used* as a substitute for atropine in ocular therapeutics.

STRAMONIUM.

What are the source, composition, action and uses of stramonium?

Stramonium is the leaves and seeds of *Datura Stramonium*, thorn-apple, Jamestown weed or gypsum (*Nat. Ord.* Solanaceæ), a common indigenous plant. It contains an alkaloid called *daturine*, isomeric with atropine and having the same action. As this is the active principle of stramonium, the effects of the latter are identical with those of belladonna, and it may be used for the same purposes. The leaves have been made into cigarettes and smoked to avert a threatened paroxysm of *asthma*, and it has been used externally as a poultice or as ointment to painful wounds, external inflammations, hemorrhoids and fissure of the anus. The official preparations (all made from the seed) are :—

EXTRACTUM STRAMONII SEMINIS (*extract of stramonium seed*), dose gr. ss (0.03).

EXTRACTUM STRAMONII SEMINIS FLUIDUM (*fluid extract of stramonium seed*), dose ℥j-v (0.06-0.3).

TINCTURA STRAMONII SEMINIS (*tincture of stramonium seed*), dose ℥v-xv (0.3-1.0).

UNGENTUM STRAMONII (*ointment of stramonium*) contains 10 per cent. of the extract.

HYOSCYAMUS.

What is hyoscyamus?

Hyoscyamus is the leaves and flowering tops of *H. niger*, or henbane (*Nat. Ord.* Solanaceæ), a native of Europe.

What does it contain?

It contains two alkaloids, *hyoscyamine* and *hyoscine* (identical with scopolamine) both of which are isomers of atropine and daturine.

What are the physiological effects of hyoscyamus and its alkaloids?

Hyoscyamine appears to be identical in action to its isomer atropine, but feebler and less irritant. It does not dilate the pupil as rapidly; sometimes retards, sometimes quickens the heart's action; and at first quickens, then slows the respiration.

Hyoscine causes dryness of the mouth and throat, flushing of the

face, sleep, slow respiration, and in large doses slow, full pulse, muscular weakness, incoördination, diaphoresis, and, it is said, a rise in temperature. The treatment of poisoning by hyoscine consists in the administration of alcoholic stimulants and coffee or caffeine; in artificial respiration; in the use of faradism, and perhaps the cautious use of physostigmine.

Hyoscyamus combines the action of its alkaloids, differing from that of belladonna in proving hypnotic.

What are the uses of hyoscyamus and its alkaloids?

Hyoscyamus has been used as a substitute for belladonna in the diseases in which the latter is employed. It has also been used as a *hypnotic* in various forms of *insanity*, in *delirium tremens*, and in *hysteria*. Added to *purgatives* it enhances their efficacy and prevents griping.

Hyoscyamine has been found useful as a *sedative* to the nervous system in *insanity*; to lessen spasm and allay pain in *vesical tenesmus*, and in *irritable bladder*; and, to prevent griping, as an addition to purgatives. It has sometimes proved useful in *chorea*.

Hyoscine is used as a *cerebral sedative* and *hypnotic* in maniacal excitement, but should only be employed temporarily, and then only in cases where *excitement* is a prominent symptom. In melancholia it has proved useless or injurious. Cases do not do well under its continued use. It may be employed to produce sleep when morphine or chloral is contraindicated in advanced renal disease and in cases of weak heart. To produce hypnotic effects it should be administered hypodermically. It has been found efficacious in *paralysis agitans* and in various forms of tremor. Disagreeable after-effects are rare.

What are the preparations and doses of these substances?

EXTRACTUM HYOSCYAMI (*extract of hyoscyamus*), a preparation of uncertain strength; dose gr. j-ij (0.065-0.13).

EXTRACTUM HYOSCYAMI FLUIDUM (*fluid extract of hyoscyamus*), dose ℥v-x (0.3-0.6).

TINCTURA HYOSCYAMI (*tincture of hyoscyamus*), dose fʒss-j (2.0-4.0).

HYOSCYAMINÆ SULPHAS (*hyoscyamine sulphate*), dose gr. ʒi-ʒi

$\frac{1}{50}$ (0.0006-0.0013) to begin with; much larger doses may be given with impunity.

HYOSCYAMINÆ HYDROBROMAS (*hyoscyamine hydrobromate*), dose gr. $\frac{1}{100}$ — $\frac{1}{50}$ (0.0006-0.0013).

HYOSCINÆ HYDROBROMAS (*hyoscine hydrobromate*), dose gr. $\frac{1}{100}$ — $\frac{1}{80}$ (0.0006-0.0008); if given hypodermically, dose gr. $\frac{1}{150}$ — $\frac{1}{100}$ (0.0001-0.00064) to begin with, but much larger doses may be used.

DUBOISIA.

What is duboisia?

Duboisia (not official) is the *leaves* of *D. myoporoides* (*Nat. Ord.* Solanaceæ), a shrub of Australia. It *contains* an alkaloid, called *duboisine*, which is isomeric with atropine. *Locally* applied to the conjunctival sac it paralyzes accommodation and dilates the pupil more rapidly but for a shorter time than does atropine, and is less irritating. Given *internally* it dilates the pupil, stimulates the respiratory centre, and depresses the inhibitory cardiac ganglia, thus causing increase in frequency both of the respiration and cardiac action. It is *used* as a substitute for atropine in ophthalmic practice.

COCA.

What is coca?

Coca is the *leaves* of *Erythroxylon Coca* (*Nat. Ord.* Lineæ), a shrub of Peru.

What does it contain?

It contains an alkaloid called *cocaine* combined with *cocatannic* acid.

What are the physiological effects of erythroxylon?

Coca is a cerebral stimulant, producing a sense of calm and well-being, followed, if the dose is large, by a pleasant delirium. It lessens the sense of fatigue and of hunger, but does not take the place of food. Small doses appear to slow, while large quicken the pulse and cause an elevation of temperature. It diminishes the amount of urea excreted. Large doses cause, also, muscular weakness, drowsiness, and finally deep sleep. When taken *in large doses habitually* it causes disordered digestion, enfeeblement of the intellectual powers, emaciation, ascites, and death from marasmus.

Cocaine produces similar effects. It is a *cerebral* stimulant, causing delirium in large doses followed by epileptiform convulsions and coma if the dose has been excessive. It first stimulates, then depresses the *reflex spinal centres*. Toxic doses depress and finally paralyze the *nerves*, these effects being more marked on the nerves of sensation. The effect on the *circulation* is disputed, but in small doses it increases the strength and frequency of the cardiac contractions. It acts as a powerful stimulant upon the *respiratory* centre, but in large doses paralyzes it, the heart continuing to beat after the breathing has stopped. *Injected hypodermically* into the superficial tissues it causes local anæsthesia; if a deep injection is given a slight general lessening of sensibility, dilated pupils, a slight rise in temperature with strong cardiac contractions result.

Locally applied it causes marked anæsthesia of the surface with which it comes in contact, with some diminution of sensibility of the deeper structures. There is at first local pallor from contraction of vessels, followed by redness. Applied to the conjunctiva it dilates the pupil (which may, however, be contracted by eserine) transiently, lessens without entirely suspending the power of accommodation, and lowers ocular tension.

In *toxic doses* cocaine causes symptoms varying much in different persons. The most common symptoms are stimulation of the mental faculties; loquacity; incoherence; depression of temperature accompanied by profuse sweating; shallow, irregular respiration; feeble pulse; dilated pupils and disturbed vision; nausea and collapse. Sometimes there is intellectual torpor with tonic or clonic convulsions. Sometimes the mental excitement is almost maniacal.

Respiratory disturbances are always marked, in severe cases being of the Cheyne-Stokes type, and the sufferer may die from respiratory paralysis.

Individual susceptibility to the drug varies greatly, and even persons who have used it before without inconvenience may experience a toxic effect from a dose of the same size as that which they had before taken. Even weak solutions applied to a mucous membrane occasionally produce alarming symptoms.

The *treatment of poisoning* by cocaine consists in inhalations of amyl nitrite and chloroform; ammonia should be given internally, and artificial respiration resorted to and continued for a long time.

Opium, or morphine, and chloral may be used if convulsions occur.

What are the medicinal uses of coca and its alkaloid?

Coca is not much employed in medicine. It may be given as a stomachic tonic in wasting diseases, as in *phthisis*.

Cocaine is principally used as a *local anæsthetic* (2 to 10 per cent. solution), in operations on the various mucous membranes, particularly those of the eye, nose, mouth, pharynx and larynx, but also to those of the vagina, rectum and urethra. Injected into the superficial tissues it diminishes sensation sufficiently to allow of the performance of many operations in minor surgery, particularly if its diffusion is prevented by compressing the circulation of the part. In ophthalmology it is useful both for this purpose and from its rapid and transient mydriatic effect.

To *relieve pain* and *contract the blood vessels* in the early stages of acute catarrhal inflammations, as *coryza*, *hay fever*, *laryngitis* and *bronchitis*; to deaden sensation and thus *lessen reflex cough* in *pertussis*, *influenza*, advanced *phthisis* and *chronic laryngitis*, it may be used in spray, or directly applied to the part by a brush. It has also been used as a *hemostatic*, applied to the mucous membranes, especially to the nasal mucous membrane, previous to an operation. It is highly recommended to allay the burning and itching of rhus poisoning. *Internally* it has been employed in *gastrodynia*, *nervous dyspepsia*, and to check *reflex vomiting*, as that of pregnancy or of sea-sickness (by the mouth or injected into the epigastric region), as a cardiac tonic in low fevers, and as a substitute for morphine in persons accustomed to the habitual use of the latter alkaloid. When used for this purpose the cocaine habit is apt to be formed, but can apparently be broken off suddenly without danger to the patient.

What are the preparations and doses of coca?

EXTRACTUM COCÆ FLUIDUM (*fluid extract of erythroxylon*), dose fʒss-ij (2.0-8.0).

COCAINÆ HYDROCHLORAS (*cocaine hydrochlorate*), dose gr. ¼-j (0.016-0.06).

ANTIPYRINE.

Is antipyrine a mydriatic anodyne?

Antipyrine and other members of the antipyretic group are anodynes of considerable power, as has already been stated, and in large doses dilate the pupil, but are not used as mydriatics.

ORDER III.—ANÆSTHETICS.

What are anæsthetics?

This name is applied to a group of medicines, the vapor of which when inhaled causes a temporary loss of sensibility to painful impressions (*analgesia*), with diminution and finally abolition of general sensation and reflex action, and lastly, complete loss of consciousness.

Name the agents of this group.

Ether, chloroform, nitrous oxide, methylene bichloride, ethyl bromide and bromoform.

ÆTHER—ETHER.

What is ether?

Ether is ethyl oxide, obtained by the distillation of alcohol with sulphuric acid. It is a colorless liquid, with a strong, disagreeable odor, and a hot, burning taste, and evaporates rapidly at ordinary temperatures. Its vapor is heavy—more than two and a half times as heavy as the atmosphere—and is very inflammable.

What are its physiological effects?

When applied *locally* ether produces a feeling of coldness by its rapid evaporation, thus obtunding the sensibility of the part, and if the application be continued the part may be entirely frozen.

When the vapor of ether is *inhaled* a sense of burning is experienced in the throat, followed by a feeling of suffocation and an irritative cough; then comes a *stage of stimulation*, during which the action of the higher nerve centres is more or less interfered with, there is a feeling of lightness in the head and buzzing in the ears, and the circulation is quickened; this passes into the *narcotic stage*, the patient becoming delirious and laughing, weeping, moaning, talking, singing or fighting. The face is suffused, the eyeballs injected and often protruding, the veins turgid, sensibility is dimin-

ished, and tonic contractions of the muscles are often present. If the inhalation is continued the *stage of anæsthesia* soon supervenes, in which the functions of the spinal cord are influenced; thus, there is not only complete insensibility, but also entire abolition of the reflexes. This stage may be continued indefinitely by carefully regulating the inhalation. If, however, too much be given, the *stage of paralysis* ensues, in which the reflex centres of the medulla oblongata are paralyzed and the respiration and circulation cease. Stertorous respiration, from paralysis of the muscles of the palate, is the first indication of the approach of this stage.

During the inhalation of ether the pupils at first contract slightly, afterward slightly dilate, but always respond to light. Sudden, wide dilatation is a sign of danger.

The pulse is increased in frequency during the early stages of ether inhalation, afterward returning gradually to about the normal. The arterial pressure is increased, due principally to contraction of the peripheral capillary vessels and to the increased power of the cardiac beat. During the stage of paralysis the pressure falls when respiratory failure begins.

In the first stage the respiration is increased both in frequency and depth, becoming much more shallow in the second stage, but again becoming deep and even stertorous in the third stage, and finally ceasing if the etherization be continued into the fourth stage from paralysis of the respiratory centre, the heart continuing to pulsate after the arrest of the respiratory movement.

The *after-effects* are headache, vomiting, and a taste of ether in the mouth. Some hours elapse after profound anæsthesia before the patient fully regains the use of his mental faculties.

Taken *internally* ether increases the secretions and is a cerebral and cardiac stimulant, being similar in its action to alcohol.

It is rapidly *eliminated* by the lungs.

What are the therapeutical uses of ether?

1. As an *anæsthetic*, ether is employed by inhalation for the prevention of pain during *surgical operations* and examinations, and during the passage of *renal* or *biliary calculi*. For these purposes the inhalation should be carried to the third stage—that of complete anæsthesia. To alleviate suffering during labor ether is often ad-

ministered, but here it should only be inhaled during the pains, and the stage of complete anæsthesia is rarely required, unless some obstetric manipulations, as version, the application of the forceps, or craniotomy is necessary, or when it is desirable to prevent the patient from "bearing down," either from fear of laceration of the soft parts, or of the strain on the circulation in cardiac disease, or where thorough relaxation of a rigid cervix uteri is desired. 2. As a *reflex depressant* it is inhaled in various forms of *convulsions* during the paroxysms, and to promote the reduction of a hernia, or aid in diagnostic manipulations by relaxing tensely contracted muscles. In persistent hiccough it is sometimes successful when given internally, or may be used by inhalation to check the paroxysm. 3. As a *cardiac stimulant* it may be given hypodermically or by the mouth in cases of syncope or threatened cardiac failure. 4. It is sometimes used as a *tæniacide*; and, 5. it has sometimes proved useful as an *anodyne* in *neuralgia* (injected in the vicinity of the nerve), and in *nervous headache* (given internally). 6. *Locally* it is employed to relieve pain in neuralgia, myalgia, muscular rheumatism and before the performance of minor surgical operations.

What cautions should be observed in administering ether?

As ether is highly inflammable, it should not be administered by inhalation at night near a candle or gas light, nor near an open fire or very hot stove. If a light is in the room it should be placed as high as possible above the patient, as the vapor of ether sinks toward the floor. Ether is contraindicated in advanced pulmonary disease, and should be cautiously administered, if at all, in cases of cerebral tumors, grave renal disease and excessive exhaustion of the nerve centres.

In cardiac diseases it is the safest anæsthetic and may be cautiously employed.

How should alarming symptoms be treated?

(1) Stop the inhalation and lower the head; (2) dash cold water on the face and chest and allow the patient plenty of fresh air; (3) use artificial respiration, make rhythmic tractions of the tongue at the rate of about 20 per minute, give atropine, strychnine, or nitroglycerin hypodermically; and (4) apply the faradaic current to the thoracic muscles.

What are the official preparations of ether and how are they administered?

ÆTHER (*ether—æther fortior*, U. S. P. 1880) contains about 96 per cent. by weight of absolute ether or ethyl oxide in alcohol and water; dose $\text{m}_{\text{v}}-\text{f}_{3\text{j}}$ (0.31–3.70).

SPIRITUS ÆTHERIS (*spirit of ether*) contains 32.5 per cent. of ether in alcohol; dose $\text{m}_{\text{xv}}-\text{f}_{3\text{j}}$ (0.92–3.70).

When ether is *administered by inhalation* the patient should be, if possible, in the recumbent position, and, the nose and throat having been sprayed with a 10 per cent. solution of cocaine (Rosenberg of Berlin), $\text{f}_{3\text{ss}}$ (15.0) may be poured on a napkin or sponge placed in a cone of stiff paper, which is then applied to the patient's mouth and nose, and kept there continuously, only removing it momentarily from time in time to add more ether, until the desired effect is produced.

The anæsthetizer should pay the strictest attention to the condition of his patient, watching closely the effect on the respiration and on the circulation as shown by the pulse and by the color of the skin or conjunctival mucous membrane.

Ether inhalations are better administered on an empty stomach, as the after-effects, especially the tendency to vomiting, are thus lessened.

Spraying the nose and throat with cocaine, as mentioned above, is said to increase the safety of the patient during the anæsthesia, and certainly prevents the coughing, gagging, and vomiting which otherwise usually accompany the action of the anæsthetic, and lessens the vomiting, headache, and persistent taste of ether which so often follow its use.

CHLOROFORMUM—CHLOROFORM.

What is chloroform?

Chloroform is methenyl chloride or formyl terchloride, and is obtained by distilling alcohol with chlorinated lime and slaked lime.

It is a limpid, colorless liquid, having a pleasant ethereal odor and an acrid, sweetish taste.

It is soluble in alcohol and the oils, but is not very soluble in water.

What are its physiological effects?

Applied *locally* it is to some extent anæsthetic; if evaporation be prevented it is an irritant and even a painful vesicant. It assists the absorption of organic alkaloids by the skin, which otherwise would not penetrate the epidermis. When the vapor of chloroform is *inhaled* anæsthesia and unconsciousness rapidly result.

The *stage of stimulation* is not as well marked as in the case of ether inhalation; the *narcotic stage* is usually short in duration and the struggles of the patient are, as a rule, not violent. The pulse is quickened and apparently strengthened until the *stage of anæsthesia*, when with the abolition of consciousness and sensation and muscular relaxation it diminishes slightly in frequency and is more or less weak. The *stage of paralysis* is one of imminent danger. In addition to the symptoms of the anæsthetic stage, there is stertorous breathing, entire abolition of the reflexes, with a weak and rapid pulse, and finally death, usually from cardiac paralysis.* When *swallowed* in small doses it stimulates the secretions and increases peristalsis. In large amounts it is an irritant, causing gastro-enteritis, and after its absorption producing anæsthesia and coma. It is eliminated by the lungs and kidneys.

How do the physiological actions of ether and chloroform compare?

Ether is the safer anæsthetic; chloroform the more rapid, more transient in its effects, which are produced by smaller amounts, pleasanter, less irritating, and with less disagreeable after-effects. It does not cause so much vomiting, and its vapor is not inflammable. Death from ether (rare) is usually the result of respiratory paralysis and is gradual, while death from chloroform (much more frequent) is usually sudden, and is due to cardiac failure.

What are the therapeutical uses of chloroform?

It has been used as an *anæsthetic* by inhalation in the same range of cases in which ether is given, but its employment for this pur-

* The Hyderabad commission concluded, from their very elaborate experiments upon dogs, that death always occurs from respiratory paralysis, a conclusion which at present is not in accord with clinical observation. The author has therefore adhered to the older view as to the mode of death under chloroform.

pose is now very generally limited to diseases like puerperal eclampsia, in which it is necessary rapidly to produce anæsthesia and depress reflex action. In *obstetric practice* it is largely used, but even here ether is safer. Administered *internally* it has proved serviceable in *cholera*, *protracted diarrhœa*, and various forms of *colic*. It has also been used as a *tœniacide*. Given before the expected chill of *malarial fevers* it frequently aborts the paroxysm. Deep *injections* in the vicinity of the affected nerve sometimes afford relief in obstinate neuralgia. *Locally* as an anodyne it is employed in the form of liniment in muscular pains and in neuralgic affections. Its *contraindications* as an anæsthetic are identical with those of ether, and in addition *it should never be administered when any disease of the circulatory system exists or if the heart is feeble from any cause whatsoever*.

How should the occurrence of dangerous symptoms be met?

Stop the inhalations; invert the patient; artificial respiration; faradization; and atropine, strychnine, or nitroglycerin hypodermically. If it has been administered internally, treat the case as one of irritant or narcotic poisoning.

What are the official preparations, and how may they be administered?

CHLOROFORMUM (*chloroform*), dose $\text{m}_{\text{xv}}\text{--f}_{3\text{j}}$ (1.0–4.0) in emulsion. As an *inhalation*, 20 drops of chloroform may be poured on a folded napkin and placed cautiously over the nose and mouth in such a manner that the vapor is freely diluted with atmospheric air. *Not more than 3½ per cent. of chloroform should be contained in the air inspired.*

AQUA CHLOROFORMI (*chloroform water*), dose $\text{f}_{3\text{j}}\text{--jv}$ (4.0–15.0).

SPIRITUS CHLOROFORMI (*spirit of chloroform*), dose $\text{f}_{3\text{ss}}\text{--ij}$ (2.0–8.0).

EMULSUM CHLOROFORMI (*emulsion of chloroform*), dose $\text{f}_{3\text{j}}\text{--ij}$ (4.0–8.0).

LINIMENTUM CHLOROFORMI (*liniment of chloroform*) contains chloroform 30 per cent. in soap liniment.

NITROUS OXIDE.

What is nitrous oxide?

Nitrous oxide (not official) or nitrogen monoxide is a colorless, almost odorless gas, made by the action of heat on ammonium nitrate.

What are its physiological actions?

When nitrous oxide is inhaled in from one to three minutes complete anæsthesia occurs with loss of consciousness, often preceded by a stage of excitement; the face is livid and the patient presents the appearance of asphyxia. The anæsthesia is probably due to diminution in the relative amount of oxygen in the blood. When the inhalation is discontinued, consciousness and sensation speedily return, and as a rule no after-effects are manifested. When death results it is generally due to asphyxia.

What are the therapeutics and the mode of administration of nitrous oxide?

As an *anæsthetic* in minor surgical operations of short duration, as the extraction of teeth or the opening of abscesses, it is advantageously employed, but when the operation necessitates the maintenance of anæsthesia for some time, ether is a safer agent.

It is best *administered* from an india-rubber bag, connected with an iron reservoir, containing at least eight gallons, furnished with a mouth-piece having two valves, so that the expired air may not pass back into the bag.

METHYLENE BICHLORIDE: ETHYL BROMIDE: BROMOFORM.

What are these substances?

Methylene bichloride (not official) is a neutral, colorless liquid, resembling chloroform in odor. Its vapor is inflammable. When *inhaled* it rapidly produces anæsthesia and unconsciousness, and these effects as rapidly disappear when the inhalation is discontinued. Death is due to cardiac paralysis. It is not much employed in this country.

Ethyl bromide (not official) is a colorless, neutral liquid, having an ethereal odor and a disagreeable, sweetish taste. When inhaled

it is anæsthetic, acting on the heart like chloroform, and seems to be fully as dangerous.

Bromoform (not official) is a heavy liquid with an agreeable taste and smell. Its effects resemble chloroform, but are slower. It is not much used as an anæsthetic. As an antispasmodic it has been used in *pertussis* in doses of gtt. ij-v (0.12-0.3) t. i. d.

LOCAL ANÆSTHETICS.

What are local anæsthetics?

They are remedies which produce an anæsthetic effect limited to the particular part to which they are applied.

What are the indications for their employment, and what are the objections to their use?

They are employed whenever it is necessary to obtund the pain in a part, either caused by disease or induced by a surgical operation, and when it is not necessary for any reason that the patient should lose consciousness.

They are generally made use of prior to minor surgical operations for the avoidance of pain, although they have also been employed for the same purpose before major operations, as amputations.

In such cases, however, it is often advantageous to have the patient unconscious, as the knowledge of what is being done, even in the absence of pain, may prove injurious by causing nervous perturbation or by allowing the patient to interfere with the operator.

They are objectionable in some cases on account of the anæmia which they produce, as during the reaction the resulting hyperæmia may be sufficiently intense to cause hemorrhage, or occasionally interfere with the healing process.

How is local anæsthesia produced?

Either by freezing the part which it is desired to anæsthetize, or by the application to, or injection into it of an anæsthetic drug.

How may a part be frozen for such a purpose, and what are the agents employed?

The part may be frozen by the direct application for a few minutes of a freezing mixture such as powdered ice and salt, or by spraying it with a very volatile liquid, such as rhigolene, ether, or ethyl chloride.

What are these substances, and how are they used ?

RHIGOLENE (not official) is a very volatile and inflammable liquid obtained in the distillation of petroleum, and is used in the ordinary hand-atomizer.

ETHER has already been spoken of as a local anæsthetic.

ETHYL CHLORIDE (not official), or *hydrochloric ether*, is a very volatile liquid, which is sold in glass bottles having a minute orifice against which a tightly fitting stopper, moved by a small lever, is kept constantly pressed by means of elastic bands. When it is desired to use it the lever is pressed down, the orifice of the bottle passed rapidly over the part to be anæsthetized at a short distance from it, and the heat of the hand volatilizing the liquid in the bottle, the part is soon frozen by the spray.

These agents are *used* as local anæsthetics in all *minor surgical operations*, such as opening abscesses or felons, the extraction of ingrowing toenails, paracentesis thoracis or abdominis, and are sometimes used to *numb* the *pain* of headache and neuralgia.

What drugs are employed topically or by hypodermic injection for this purpose ?

COCAINE (*vid. Mydriatic Anodynes*) is used locally as has already been mentioned, applied to the conjunctival, to the nasal, and to other mucous membranes as a local anæsthetic.

It is also used by injection into the skin for the same purpose. When this is desired the part is first sprayed with ethyl chloride, and as soon as sensibility is obtunded the needle of the hypodermic syringe is inserted and a few drops of the cocaine solution instilled just beneath the integument. By injecting the solution in front of the point of the needle, which is then pushed further into the anæsthetized area, and a few more drops inserted, it is possible gradually to anæsthetize quite a large surface. By casting a ligature around the part and thus preventing the cocaine from passing out by the blood current, this anæsthetic action may be prolonged.

By this or similar means local anæsthesia has been maintained for periods considerably over an hour, during which operative procedures of some magnitude have been performed. It is probably better, however, to employ general anæsthesia for major surgery unless

there is some contraindication to its employment, and to restrict the anæsthetic use of cocaine to minor surgery.

EUCAINE (not official) has been highly recommended as a local anæsthetic. The hydrochloride is used, and occurs as brilliant colorless crystals, readily soluble in water (which is not the case with eucaine itself).

It is much more stable than cocaine, not even decomposing when kept in solution. Locally applied to mucous membranes in 2 to 5 per cent. solutions it causes anæsthesia, but it neither dilates the pupil nor causes ischæmia, as does cocaine, when applied to the conjunctiva. In fact, it appears to favor hyperæmia and does not interfere with the pupil reflex. It is said to produce a primary reduction in the cardiac frequency when given internally. In animals after large doses there is general nervous excitement followed by paralysis, sometimes preceded by tonic and clonic convulsions. The blood-pressure falls and finally death occurs from respiratory failure.

No toxic symptoms have been reported as occurring in the human subject.

It has been used as a substitute for cocaine to produce local anæsthesia, and may be employed in solutions of the strength of 2 to 5 per cent. It has also been injected subcutaneously in the same manner as cocaine prior to the opening of abscesses or the removal of small tumors. It is said that as much as gr. xxx (1.9) has been used subcutaneously for this purpose without ill effect.

MENTHOL (official) is a "*stearopten* (having the character of a secondary alcohol), *obtained from the official oil of peppermint* (from *Mentha piperita*), or from Japanese or Chinese oil of peppermint (from *Mentha arvensis*" "and *Mentha canadensis*," *Nat. Ord. Labiatae*). It occurs in the form of colorless, acicular or prismatic crystals, having a strong and pure odor of peppermint, and a warm, aromatic taste, followed by a sensation of cold when air is drawn into the mouth, and being only slightly soluble in water, but freely so in alcohol, ether, and chloroform. Menthol is a superficial, local anæsthetic. Given internally to animals in sufficient doses it depresses both the sensory and motor nerves, lowers the temperature, and depresses the respiration, which becomes slow and shallow.

It is chiefly *used* as a *local anæsthetic* painted or sprayed on the part in ethereal or alcoholic solutions varying in strength from 10 to 50 per cent., in cases of neuralgia, toothache, and other superficial pains. The author has seen the burning pains which often follow herpes zoster quickly relieved and often permanently cured by a few applications of a 50 per cent. alcoholic solution. Urticaria and senile pruritus may be temporarily relieved by the application of a 10 per cent. solution in albolene. It is also used with lanolin as an ointment, and has been used as an antiphlogistic remedy in superficial inflammations of the skin and mucous membranes.

GUAIACOL (not official), which has been considered among the antipyretics (q. v.), is also used as a *local anæsthetic* in *neuralgiæ*, especially in *sciatica*, and in rheumatic pains.

Various other remedies, among which may be mentioned *Tincture of Aconite* (*vid. Cardiac Sedatives*), have been used locally in neuralgia for their local anæsthetic effect.

ORDER IV.—ANTISPASMODICS.

What are antispasmodics?

Antispasmodics are remedies which prevent or allay spasmodic action of voluntary or involuntary muscles by increasing the inhibiting power of the higher over the lower nerve centres.

Name the antispasmodics.

Musk, valerian, the valerianates, asafoetida, camphor, ethereal oil, and Hoffman's anodyne, cypripedium, and scutellaria.

MOSCHUS—MUSK.

What is musk?

Musk is the *dried secretion* from the preputial follicles of *Moschus moschiferus*, or musk deer.

What are its effects and uses?

Musk is a mild cerebral stimulant and antispasmodic, slightly increasing the cardiac frequency. In cases of nervous exhaustion, however, it acts more powerfully, and when a pure article can be obtained no better nervous stimulant can be employed in the later stages of *typhoid fever* with great depression of vital forces, and in *typhoid pneumonia*. It is also of use in *hysteria*, *hiccough*, and *chorea*.

What are the preparations of musk?

MOSCHUS (*musk*) is very expensive and is often adulterated, and imitations are not infrequently substituted for the genuine article; dose gr. x-xv (0.65-1.0) in capsule or emulsion. It is often given per rectum.

TINCTURA MOSCHI (*tincture of musk*), dose f3j-ij (4.0-8.0).

VALERIANA—VALERIAN.

What is valerian, and what does it contain?

Valerian is the *rhizome* and *roots* of *V. officinalis* or wild valerian (*Nat. Ord. Valerianæ*). It contains a volatile oil and a colorless volatile acid (*valerianic*).

What are the effects and uses of valerian?

Large doses of valerian may cause nausea, vomiting, and sometimes colicky pains. It slightly stimulates the brain and circulation, and increases the secretions of the skin and kidneys. Its effects are more decided in nervous excitement due to depression of the cerebral functions, as in *hysteria* or "*nervousness*," and in *hypochondriasis*. It has also been used as a cerebral stimulant in the *delirium* of low fevers and in *delirium tremens*.

What are the preparations and doses of valerian?

EXTRACTUM VALERIANÆ FLUIDUM (*fluid extract of valerian*), dose f3j (4.0).

TINCTURA VALERIANÆ (*tincture of valerian*), dose f3j (4.0).

TINCTURA VALERIANÆ AMMONIATA (*ammoniated tincture of valerian*), dose f3j-ij (4.0-8.0).

AMMONII VALERIANAS (*ammonium valerianate*) is much used in *hysteria*, *neuralgia*, and *nervous headache*; dose gr. ij-x (0.13-0.65) in capsules or elixir.

Iron valerianate and *quinine valerianate* have already been considered; *zinc valerianate* will be noticed under the zinc preparations.

ASAFÆTIDA—ASAFETIDA.

What is asafetida, and of what is it composed?

Asafetida is a *gum-resin* obtained from the *root* of *Ferula foetida* (*Nat. Ord. Umbelliferæ*), a native of Afghanistan. It is composed of a gum-resin united to a volatile oil.

What are its effects and uses?

It is a gastric stimulant and carminative, and produces a sense of exhilaration. It increases the mucous secretions and peristalsis, and in large doses quickens the pulse and respiration. Very large doses may cause nausea and vomiting.

It is *used* as an *antispasmodic* in *hysteria* and *nervous exhaustion*; as a *carminative* in *tympanites* (by the rectum); as a *stomachic tonic* in *dyspepsia* with flatulent colic, and as an *expectorant* in *chronic bronchitis* and in *whooping-cough*, where it also acts as an antispasmodic.

What are its preparations and doses?

EMULSUM ASAFÆTIDÆ (*emulsion of asafetida—lac asfætida*), dose fʒss-j (15.0–31.0), or as an enema fʒij–iv (62.0–124.0).

PILULÆ ASAFÆTIDÆ (*pills of asafetida*), each pill containing gr. ij (0.2) of asafetida; dose No. ij–iv.

TINCTURA ASAFÆTIDÆ (*tincture of asafetida*) contains 20 per cent. of asafetida; dose fʒss-j (2.0–4.0).

PILULÆ ALOES ET ASAFÆTIDÆ (*pills of aloes and asafetida*), each pill contains of aloes, asafetida and soap gr. j½ (0.09) each.

CAMPHORA—CAMPHOR.

What is camphor?

Camphor is a *stearopten*, derived from *Cinnamomum Camphora*, or camphor laurel (*Nat. Ord. Laurineæ*), an evergreen tree of China and Japan. It is obtained by boiling chips of the tree in water and skimming off the camphor as it rises to the surface. This *crude camphor* is afterward *refined* by sublimation with quicklime.

What are the physiological effects of camphor?

Locally camphor is an irritant. *Internally in moderate doses* it is to some extent an anodyne and a cerebral exhilarant, sometimes quickening the pulse and raising arterial tension, and occasionally proving aphrodisiac. In *large doses* (gr. xx (0.13)) it depresses the circulation, and produces lassitude and diaphoresis. It is also anaphrodisiac. In *toxic doses* (ʒss-j (2.0–4.0)) it causes faintness, vertigo, headache, delirium, violent epigastric pain, unconsciousness, and sometimes convulsions. No case is recorded to have proved fatal in an adult.

What are the medicinal uses of camphor?

It is employed as an *antispasmodic* in “*nervousness*,” *nervous headaches*, *hysterical complaints*, *dysmenorrhœa*, “*after-pains*,” simple *diarrhœa*, and combined with other remedies in *cholera*. It is also used in *low forms of fever*, to quiet delirium and sustain the system. As an *anaphrodisiac* to quiet abnormal sexual excitement in *chordee*, *nymphomania*, and *priapism*. *Externally* it is employed as a liniment in myalgia, muscular rheumatism, and sprains. Dissolved in oil of turpentine it forms an efficient liniment in mild bronchial affections.

What are its preparations and their uses?

CAMPORA (*camphor*): dose gr. j–x (0.06–0.65), or even gr. xx (1.3) given in emulsion, or in pill or capsule. Not much used in this form.

AQUA CAMPHORÆ (*camphor-water*) contains 1 part of camphor in 125 parts of the preparation; dose f̄ss–ij (15.0–60.0). It is used chiefly as a vehicle or where a mild antispasmodic is needed.

SPIRITUS CAMPHORÆ (*spirit of camphor*) contains 10 per cent. of camphor. It is used as an embrocation, and is also given internally where a stronger preparation than the water is desired; dose m̄v–f̄j (0.3–4.0).

LINIMENTUM CAMPHORÆ (*camphor-liniment*) contains 20 per cent. of camphor in cottonseed oil.

CERATUM CAMPHORÆ (*camphor-cerate*) contains 1 per cent. of camphor-liniment.

LINIMENTUM SAPONIS (*soap-liniment*) is much used externally, either alone or combined with stronger liniments.

CAMPORA MONOBROMATA (*monobromated camphor*); dose gr. v–x (0.32–0.65) in gelatin-coated pill or capsule. It is useful as an antispasmodic in *hysteria*, and as an anaphrodisiac in *chordee* and *priapism*.

Hope's Camphor Mixture (not official) consists of nitrous acid f̄j (4.0); laudanum m̄xij (0.74); camphor-water f̄viiij (240.0). Dose, f̄ss (15.0) repeated as required. Much used in diarrhœa and dysentery.

OLEUM ÆTHEREUM—ETHEREAL OIL.

What is ethereal oil?

Ethereal oil, or oil of wine, is made by distilling alcohol with an excess of sulphuric acid and diluting with an equal volume of ether, and is a nearly colorless volatile liquid. It is antispasmodic, and in small doses a cardiac and vaso-motor stimulant, while in large doses it paralyzes the cardiac muscle and the vaso-motor nerves. It is only used medicinally as

SPIRITUS ÆTHERIS COMPOSITUS—COMPOUND SPIRIT OF ETHER.

What is this substance?

Compound spirit of ether, or *Hoffman's anodyne*, is a solution of ethereal oil (2.5) in ether (32.5) and alcohol (65.0), and is a good antispasmodic, circulatory stimulant and carminative, and is useful in the class of diseases to which antispasmodics are applicable, especially when a cardiac stimulant is required; dose f3j-ij (4.0-8.0) repeated, and best administered in iced water.

CYPRIPEDIUM: SCUTELLARIA.

What are these remedies?

Cypripedium is the *rhizome* and *roots* of *C. pubescens* and of *C. parviflorum* (*Nat. Ord. Orchideæ*), indigenous plants, commonly called ladies' slipper and moccasin plant. *Scutellaria* is the *herb* *S. laterifolia* or skullcap (*Nat. Ord. Labiatae*), an indigenous herb. They possess mild antispasmodic properties, and have been used as substitutes for valerian.

EXTRACTUM CYPRIPEDII FLUIDUM (*fluid extract of cypripedium*), dose ℥x-xx (0.6-1.2).

EXTRACTUM SCUTELLARIÆ FLUIDUM (*fluid extract of scutellaria*), dose f3j (4.0).

ORDER V.—EXCITO-MOTORS.

What are excito-motors?

Excito-motors are medicines which by stimulating the reflex centres of the spinal cord excite muscular action.

Name the excito-motors.

Nux vomica and its alkaloids; *rhus toxicodendron* and *picrotoxin*.

NUX VOMICA.

What is nux vomica ?

Nux vomica is the seed of the *Strychnos Nux-vomica* or poison nut (*Nat. Ord.* Loganiaceæ), a tree of India.

What active principles does it contain ?

It contains two alkaloids, *strychnine* and *brucine*, combined with *igasuric acid*, and according to some investigators, a third alkaloid, *igasurine*, has been discovered.

How may the presence of these alkaloids be detected ?

Strychnine may be detected by the addition of a crystal of potassium bichromate to a solution of the alkaloid in concentrated sulphuric acid on a porcelain plate, when a deep, rich blue color passing through purple into red rapidly makes its appearance.

The *physiological test* is the production of tetanic spasms in a small animal (preferably a frog) to whom a minute amount of the suspected substance has been administered.

Brucine with concentrated nitric acid strikes a blood-red color, fading into yellow.

What are the physiological effects of nux vomica ?

The effects of *nux vomica* are identical with those of its alkaloids. The action of *brucine* is similar to, but much weaker than, that of *strychnine*.

In *small doses* it increases the appetite and digestion and the peristaltic movements of the intestines. When continued for some time in small doses it is a cardiac and vaso-motor stimulant, and increases the sensibility of the sensory nerves and nerves of special sense, so that sight, hearing and touch are more acute, and also stimulates the respiratory centre. In *large doses* it causes malaise, restlessness, anxiety, muscular twitchings, and finally clonic convulsions. *Strychnine* acts principally on the *spinal cord*, stimulating especially its reflex, but also its motor, vaso-motor and sensory functions. After large doses, the reflex functions are so exalted that the slightest impression conveyed by a sensory nerve produces general convulsions. It is eliminated unchanged in the urine.

What are the symptoms and treatment of strychnine poisoning ?

Sudden shuddering, quickly followed by tetanic convulsions of all

the voluntary muscles, accompanied with muscular pains, opisthotonos, the *risus sardonicus* (a ghastly grin), a pale face, becoming livid from embarrassed respiration. After a time the muscles relax and an interval of quiet occurs, to be succeeded again and again by convulsions, if a liberal dose has been taken and is not antagonized by antidotes, until the sufferer dies of asphyxia from fixation of the respiratory muscles during a paroxysm ; or the patient may die of paralysis and collapse during one of the intervals. The mind is clear to the end.

It differs from *tetanus* in history, in clonic (not tonic) spasms, in the muscles of the jaw not being affected before other voluntary muscles.

The *treatment* consists in (1) evacuating the contents of the stomach by emetics or the stomach-pump before the convulsive stage, or washing out the stomach under ether or chloroform if not seen until the convulsions have commenced ; (2) giving tannic acid as a chemical antidote and chloral (hypodermically) as an antagonist ; and (3) the frequent use of the catheter to prevent resorption of the poison from the bladder.

What are the therapeutical uses of *nux vomica* and *strychnine* ?

It is employed (1) as an *excito-motor* in all cases of *paralysis* due to depression of the spinal or other motor centres, as in *diphtheritic*, *alcoholic*, or *lead paralyses*, *paraplegia*, *infantile paralysis* and *hemiplegia* after the subsidence of irritation. As it stimulates the nerve centres, it is *contraindicated* whenever inflammation or excitation of these centres is present ; hence, in paralyses due to the presence of blood-clots it may do great injury if too early exhibited. In *constipation* due to atony and relaxation of muscular fibres, or in *diarrhœa*, *dysentery*, *impotence*, or *incontinence of urine* from the same cause, it is very useful. (2) In *amaurosis*, especially when due to the abuse of tobacco or of alcohol, and in *amblyopia* from disease, strychnine pushed to the verge of tolerance is very efficacious. It is more useful before atrophy of the optic nerve fibres has occurred. (3) As a *cardiac* and *respiratory stimulant* in *dyspnœa* depending on chronic bronchitis, emphysema, phthisis, or due to cardiac disease, as fatty or dilated heart, it is of value, and is often combined or alternated with digitalis in the latter affection. (4) As a *general*

tonic it is advantageously given in *atonic dyspepsia*, *gastric catarrh*, *anæmia* and *chlorosis* (combined with iron), as an adjunct to quinine in obstinate *malarial fevers*, in various *neuralgiæ* occurring in debilitated subjects, and in *convalescence* from acute diseases (combined with iron and other tonics). Minute doses of *nux vomica* (tincture) will often arrest the *vomiting of pregnancy*.

What are the official preparations and their doses?

EXTRACTUM NUCIS VOMICÆ (*extract of nux vomica*), dose gr. $\frac{1}{4}$ -j (0.016-0.065) given in pill. It is frequently combined with purgatives to enhance their effect.

EXTRACTUM NUCIS VOMICÆ FLUIDUM (*fluid extract of nux vomica*), dose ℥ij-xx (0.12-0.61).

TINCTURA NUCIS VOMICÆ (*tincture of nux vomica*), dose ℥j-xx (0.06-1.2). Chiefly used as a bitter tonic.

STRYCHNINA (*strychnine*), preferably given as—

STRYCHNINÆ SULPHAS (*strychnine sulphate*), dose gr. $\frac{1}{30}$ - $\frac{1}{20}$ (0.002-0.003) internally, or gr. $\frac{1}{60}$ (0.001), cautiously increased, if used hypodermically. As a cardiac tonic strychnine is often given in large doses, frequently repeated. It is not safe, however, to begin with a larger dose than gr. $\frac{1}{30}$ - $\frac{1}{20}$ (0.002-0.003). As a tonic, minute doses may be given in pill, alone or combined with other medicines. In nervous affections it should be given in solution, the dose being gradually increased until slight muscular twitchings are produced.

Iron and strychnine citrate and *syrup of iron, quinine and strychnine phosphates* have been noticed with the iron preparations.

RHUS TOXICODENDRON.

What is this substance?

It is the *fresh leaves* of the *Rhus radicans*, poison-oak or poison-ivy (*Nat. Ord. Anacardiæ*), which contains *toxicodendric acid*. *Locally* applied it produces in many persons a very distressing eczematous eruption attended with much inflammation and swelling of the skin. Some persons are so susceptible that these symptoms occur if they are in the vicinity of the growing plant, even without coming in contact with it.

The treatment consists in alkaline washes, lotions of lead or corro-

sive sublimate (1 to 8000 or even weaker) or the application of carbolic acid in glycerin. *Internally* it is a gastro-intestinal irritant, causing also drowsiness, stupor, delirium and convulsions. Its precise use is not clear, but it has been given as a substitute for nux vomica in *paralysis* and in *incontinence of urine*. *Dose* gr. j-ij (0.06-0.19), perhaps, if given at all; best administered in the form of a tincture.

PICROTOXINUM—PICROTOXIN.

What is picrotoxin?

Picrotoxin is a *neutral principle* prepared from the seeds of *Anamirta paniculata* (*Nat. Ord.* Menispermaceæ), a climbing shrub of India. Its *action* on the brain resembles that of belladonna, while it affects the spinal centres somewhat like nux vomica: thus, it causes nausea, vertigo, delirium, incoördination, muscular twitching, followed by epileptiform convulsions, alternating tonic and clonic spasms, coma and cardiac paralysis. It stimulates all the motor and inhibitory centres of the medulla, particularly the respiratory and vagus centres, and excites the motor and reflex spinal centres. It is at present chiefly used to check the *night-sweats of phthisis*, but has also been recommended as an *excito-motor* in *functional paralysis* and in *paralysis or relaxation of the sphincters*. It is said to be of service in *chorea* when given in full doses. *Locally* it is used as an ointment (gr. v-x to $\overline{3j}$ (0.3-0.6 to 31.0)) in parasitic skin diseases and to destroy pediculi, but care should be used in applying it, as constitutional effects and even death have followed its employment externally. *Dose* gr. $\frac{2}{100}$ - $\frac{1}{60}$ (0.0003-0.001).

ORDER VI.—DEPRESSO-MOTORS.

What are depresso-motors?

They are remedies which are used to depress the functions of the spinal cord, motor nerves or muscles, thus lessening motivity.

Name the members of this group.

Physostigma; chloral; the bromides; the nitrites; gelsemium; prussic acid; the cyanides; oil of bitter almonds; lobelia; conium; woorara, and tobacco.

PHYSOSTIGMA.

What is physostigma, and what active principles does it contain?

Physostigma, or Calabar bean, is the seed of *P. venenosum* (Nat. Ord. Leguminosæ), a plant found on the western coast of Africa. It contains an alkaloid called *eserine* or *physostigmine*, and it is claimed that a second alkaloid (a tetanizing agent), called *calabarine*, has also been discovered, but this is probably a derivative alkaloid of eserine.

What are the physiological effects of physostigma?

Calabar bean is a reflex depressant, causing also, in full doses, giddiness, slow and often irregular pulse, great muscular weakness, nausea and sometimes vomiting, generally free purgation, and usually contraction of the pupils. When poisonous doses are taken, there are, in addition, muscular tremors and abolition of reflex action, slow pulse and respiration, myosis and motor paralysis, and death from respiratory failure.

It depresses and finally paralyzes the *reflex and motor centres of the spinal cord*, having but slight effect upon the nerves themselves. It slows the heart (by greatly lengthening the diastolic pause) and increases its power (acting either on the cardiac motor-ganglia or muscle). The *blood pressure* at first falls, but soon rises and remains high until death approaches. It stimulates *intestinal peristalsis* and increases the secretions of the part. On the *eye* its action is probably directly opposed to that of atropine, for no matter how used it contracts the pupil, probably by paralyzing the peripheral sympathetic nerve-fibres in the iris and at the same time stimulating the end organs of the 3d pair of nerves. It is probably *eliminated* by the urine.

How should poisoning by this drug be treated?

(1) Give emetics and cathartics, and use a catheter to prevent further absorption of poison; (2) use atropine hypodermically as a physiological antagonist, and (3) give other cardiac and respiratory stimulants as digitalis, ammonia, and alcohol.

For what purpose is it employed medicinally?

As a *depresso-motor* it is used in *tetanus*, *trismus neonatorum*, *epilepsy* and *chorea* with some success. It has also been given as a

physiological antagonist in *strychnine poisoning*, but is probably inferior to chloral for this purpose.

From its influence over *intestinal peristalsis*, it is advantageously used in *habitual constipation*, especially when associated with chronic intestinal dyspepsia and dry, hard stools. The *alkaloid* is used in ocular therapeutics, to prevent or overcome adhesions of the iris, lessen intra-ocular tension and contract the pupil.

What are its preparations and doses?

EXTRACTUM PHYSOSTIGMATIS (*extract of physostigma*), dose gr. $\frac{1}{12}$ – $\frac{1}{2}$ (0.005–0.03).

TINCTURA PHYSOSTIGMATIS (*tincture of physostigma*), dose $\text{m}\bar{\text{v}}$ –xxx (0.3–2.0).

PHYSOSTIGMINÆ SULPHAS (*physostigmine sulphate*), a very deliquescent salt; dose $\frac{1}{60}$ – $\frac{1}{12}$ (0.001–0.005).

PHYSOSTIGMINÆ SALICYLAS (*physostigmine salicylate*), preferred to the sulphate because of its greater permanency; dose gr. $\frac{1}{60}$ – $\frac{1}{12}$ (0.001–0.005).

CHLORAL and THE BROMIDES have been discussed under the head of *hypnotics* (*vid.* pp. 95–99).

THE NITRITES.

What are the nitrites?

The *nitrites* are salts formed by the union of *nitrous acid** with a base.

Amyl nitris (*amyl nitrite*) is the only official organic nitrite (except the sweet spirits of nitre, which contains ethyl nitrite and will be considered elsewhere). It is a yellowish, oily, very volatile liquid, with a pleasant, fruity odor. *Nitroglycerin* (*trinitrin*, *trininitro-glycerin* or *glonoin*) is a colorless, oily liquid, without odor, soluble in ether and alcohol. It forms the basis of various explosives. It is really a *trinitrate* (not a *nitrite*) of glyceryl or propenyl [$\text{C}_3\text{H}_5(\text{NO}_3)_3$]. *Potassium and sodium nitrites* are also used.

What is the physiological action of the nitrites?

They cause vertigo, headache, excited cardiac action, great dilata-

* See foot-note on p. 71.

tion of the arterioles and capillaries (from paralysis of their muscular coats), and consequently greatly diminished blood pressure, slowed respiration, lowered temperature, and diminished reflexes. After toxic doses, the pupils dilate, the muscular system relaxes, the blood becomes darker (probably from the formation of methemoglobin), the respiration and circulation become slow and irregular, and finally death occurs from respiratory failure, sensation and consciousness being preserved to the last. They depress both the *motor and reflex spinal* centres without influencing either the sensory nerves or centres. They first stimulate the *heart*, but soon depress and finally paralyze the cardiac muscle by direct action. The *temperature* is lowered, owing to lessened oxydation. The nitrites are all *muscle poisons*. They are *eliminated* by the urine, increasing greatly the amount of water and the uric acid and urea. The action of *amyl nitrite* is the most rapid, but most transient; *nitroglycerin* is the slowest, but its effects are more lasting,—the headache, however, which it causes, is very persistent, and often constitutes a great objection to its use. It sometimes causes nausea and vomiting.

What are the therapeutical uses of the nitrites?

The nitrites are used (1) as *depresso-motors* to relax spasm in *angina pectoris*, *asthma*, *tetanus*, puerperal and other forms of *convulsions*, *strychnine poisoning*, congestive and nervous *dysmenorrhœa*, and in all cases of very *high arterial tension* from spasm of the arterioles; thus, in *renal asthma* the inhalation of a few drops of amyl nitrite quickly relieves the spasm, and *migraine* with capillary spasm is frequently benefited by it. Given just before the expected chill of *malarial fevers* it will abort it, and sometimes avert the paroxysm. If amyl nitrite is inhaled when the “*aura epileptica*” is experienced, it will often prevent the *epileptic* attack. Nitroglycerin has proved of value in *chronic interstitial nephritis*, lessening the albuminuria and relieving the arterial spasm. Obstinate *hic-cough* has been benefited by nitroglycerin.

(2) As *cardiac stimulants* the nitrites are employed in *syncope*, *cardiac dyspnœa*, and in *threatened cardiac failure* during the administration of anæsthetics, especially chloroform.

(3) They have also been found useful in the *nausea and vomiting* of sea-sickness and of pregnancy.

What are the doses of the nitrites, and how are they administered?

AMYL NITRIS (*amyl nitrite*), given internally or by inhalation; dose mij-v (0.1–0.3).

SODII NITRIS (*sodium nitrite*), dose gr. ij–iij (0.1–0.2).

Potassium nitrite is not official. It may be used in the same doses as the sodium salt.

SPIRITUS GLONOIINI (*spirit of glonoin—spirit of nitroglycerin*) is a 1 per cent. alcoholic solution of nitroglycerin, the dose of which is mss (0.03) cautiously increased.

Triturates of nitroglycerin, each containing gr. $\frac{1}{100}$ (0.0006), are also employed.

Amyl nitrite is usually given by *inhalation*, but may be administered internally on sugar, or even hypodermically. The other preparations are given by the mouth. If a speedy action is required amyl nitrite should be used, and after the immediate danger is over, if it is desired to maintain the effect, the other nitrites or nitroglycerin may be given.

GELSEMIUM.

What is gelsemium, and what chemical principles does it contain?

Gelsemium is the *rhizome* and *roots* of *G. sempervirens*, the yellow or Carolina jasmine (*Nat. Ord. Loganiaceæ*), a climbing plant of the Southern States. It contains two alkaloids, *gelsemine* and *gelseminine*, combined with *gelseminic acid*, a volatile oil, resin, etc.

What are its physiological effects?

Small doses produce languor, dizziness, and perhaps slight lowering of the pulse rate; *large doses* diminish the force and frequency of the pulse and lower arterial pressure, depress respiration, cause frontal headache, and disordered, usually double vision; *toxic doses* cause extreme muscular weakness, partial blindness, dilated pupil, often internal strabismus, ptosis, incoördination, the jaw drops, articulation is impaired, general sensibility is much diminished, respiration is slowed, the pulse is very feeble, the temperature declines, a profuse cold sweat breaks out, and death occurs from respiratory failure, preceded in animals by convulsions followed by

general paralysis. The mind remains clear until the end approaches. The paralyzes of sensation and of motion appear to be of *spinal origin*, and the convulsions in animals have been attributed to excitation of the reflex spinal centres from *gelseminic acid*, which is supposed to be a tetanizing agent. The *respiratory centre* is depressed and finally paralyzed. The poison appears in toxic doses to depress the *heart* directly. It paralyzes the end-organs of the motor oculi, thus causing mydriasis with paralysis of accommodation and ptosis, and must also act on the 6th nerve, as the action of the external rectus muscle is much enfeebled.

What are its therapeutic uses?

It has been used as a *motor depressant* in *asthma*, *laryngismus stridulus*, *pertussis*, and *nervous cough* and *tic douloureux*; as a *sensor-depressant* in *trigeminal neuralgia*, especially when associated with spasm, and in various other *neuralgic affections*; as a *calmative* in *mania* (full doses); as a *cardiac sedative* in *sthenic fevers* and *inflammations*, and in *ocular therapeutics*. Its chief use is in *neuralgia*, especially of the 5th nerve.

What are its preparations and their doses?

EXTRACTUM GELSEMII FLUIDUM (*fluid extract of gelsemium*), dose m_{V-x} (0.3–0.6) (f3j (4.0) has proved fatal), repeated every 2 or 3 hours until constitutional effects are produced.

TINCTURA GELSEMII (*tincture of gelsemium*), dose m_{V-xx} (0.3–1.2).

ACIDUM HYDROCYANICUM DILUTUM—DILUTED HYDROCYANIC ACID.

What are the sources of hydrocyanic acid?

Hydrocyanic or *prussic acid* exists in bitter almonds, peach kernels and leaves, wild cherry, etc., but is used in medicine only in a very diluted solution. The *diluted acid* is formed by the reaction of sulphuric acid and water upon potassium ferrocyanide or of hydrochloric acid on silver cyanide, and is a colorless, volatile liquid with an odor resembling that of bitter almonds. It contains 2 per cent. of the absolute acid.

How may the presence of prussic acid be detected?

1. The addition of a solution of silver nitrate produces a white

precipitate entirely soluble in boiling concentrated nitric acid. 2. The addition of a small amount of a mixed solution of ferrous and ferric sulphate, and afterward a little liquor potassæ, gives a dirty, greenish-blue precipitate, which changes to Prussian blue when slightly acidulated with hydrochloric acid.

What are its physiological effects?

Locally applied it passes through the cuticle and paralyzes the end-organs of the sensitive nerves beneath. When its vapor is inhaled or when the acid is taken in sufficient doses, it kills almost immediately, from arrest of circulation. If the toxic dose has not been large enough to arrest the circulation instantly, dyspnœa and depression of the heart occur, followed by dilated pupils, convulsions and unconsciousness, and death from asphyxia and collapse.

The convulsions are of *cerebral* origin. Prussic acid is a direct nerve and muscle poison. In large doses it causes diastolic arrest by direct action either on the *heart muscle* or its contained ganglia: in small doses it slows the heart by stimulation of the inhibitory centres. It stimulates at first the vaso-motor system and for a short time raises arterial tension, but it soon depresses and finally paralyzes the vaso-motor apparatus and the blood pressure falls. The venous blood at first becomes of an arterial hue but soon darkens, and after death even the arterial blood has lost its bright color. The cause is not well understood. Asphyxia is due to paralysis of the respiratory centre. It is rapidly eliminated by the lungs and kidneys.

How should poisoning by prussic acid be treated?

Evacuate, if possible, the contents of the stomach; give atropine hypodermically as a respiratory stimulant; give ammonia freely by inhalation, by the mouth and by intra-venous injection; practice artificial respiration assiduously and use alternate hot and cold douches. Death usually results so rapidly that nothing is of much service.

What are the medicinal uses of diluted hydrocyanic acid?

It is used (1) as a *depresso-motor* and *nerve sedative* in various forms of *cough*, as that of phthisis, also in *asthma* and in *whooping cough*. In obstinate *vomiting* it is sometimes successful. In *gastro-dynia* and in *enteralgia* it has proved beneficial, probably by a local

action on the nerves of the parts. When successful in these cases, it is so speedily.

(2) *Locally* (fʒj-ij to Oj (4.0-11.0 to 480.0)) it is used as an *anodyne* in neuralgia and in many cutaneous affections attended with pruritus. Great caution should be used when it is thus applied, as fatal results have attended its absorption through an abraded surface.

How is it administered?

ACIDUM HYDROCYANICUM DILUTUM (*diluted hydrocyanic acid*), dose gtt. j-ijj (0.06-0.18) diluted.

THE CYANIDES.

What are the effects, uses and doses of the cyanides?

POTASSII CYANIDUM (*potassium cyanide*). *Locally* it is an irritant. It may be absorbed through the skin, when the application is prolonged, and cause poisoning. When administered *internally* it reacts with the acids of the gastric juice and prussic acid is set free, consequently the systemic effects are identical with those of that acid, as a substitute for which it is used. Dose gr. $\frac{1}{12}$ — $\frac{1}{8}$ (0.005-0.008).

ARGENTI CYANIDIUM (*silver cyanide*) is only used in medicine in making prussic acid.

Mercury cyanide has been noticed among the mercurial preparations.

OLEUM AMYGDALÆ AMARÆ—OIL OF BITTER ALMOND.

What are the effects, uses, and dose of this oil?

Oil of bitter almond is a volatile oil obtained by distillation from the kernel of the fruit of *Prunus Amygdalus* (*var. amara*, *Nat. Ord. Rosaceæ*). It contains *benzoic aldehyd* and *hydrocyanic acid* (due to the reaction of *amygdalin* with water, in the presence of an albuminous ferment called *emulsin*). Its effects and uses are the same as those of prussic acid. Dose m $\frac{1}{4}$ -ss (0.015-0.03), in emulsion.

AQUA AMYGDALÆ AMARÆ (*bitter almond water*) contains $\frac{1}{10}$ of 1 per cent. of the oil, and is used as a vehicle; dose fʒj-iv (4.0-15.0).

SPIRITUS AMYGDALÆ AMARÆ (*spirits of bitter almond—essence*

of bitter almond) contains 1 per cent. of the oil; dose $\text{m}\text{j}-\text{xx}$ (0.06–1.2).

SYRUPUS AMYGDALÆ (*syrup of almonds*), made from both sweet and bitter almonds, is used as a vehicle.

EMULSUM AMYGDALÆ (*almond emulsion—milk of almonds*), made from the sweet almonds alone, is also used as a vehicle.

LOBELIA.

What is lobelia, and what are its active ingredients?

Lobelia is the leaves and tops of *L. inflata*, or Indian tobacco (*Nat. Ord.* Lobeliaceæ), an indigenous plant. It contains an *alkaloid*, *lobeline*; also *lobelic acid*, fixed and volatile oils, etc.

What are its effects and uses?

In large doses it causes nausea, violent vomiting with great depression, sometimes purging, feeble pulse, almost complete muscular relaxation, and, if a *lethal dose* has been taken, collapse, coma, sometimes preceded by convulsions, and death from paralysis of the respiratory centre.

Large doses appear to paralyze the motor nerves, the vaso-motor centre and the peripheral vagi. Its action has not been fully studied.

The *treatment of lobelia poisoning* consists in emetics, tannic acid, morphine, alcohol, ammonia and counter-irritation to the surface.

It is used as a *depresso-motor* and *expectorant* in *asthma* and *acute bronchitis* with spasmodic dyspnoea. In other spasmodic diseases it has been superseded by more reliable and less dangerous remedies. It is too depressing in its action to be used as an emetic.

Locally, in infusion ($\text{f}\text{3j}$ to Oj (30.0 to 480.0)), it is highly recommended as an application to *poison- oak eczema*.

What are the preparations of lobelia and their doses?

EXTRACTUM LOBELIÆ FLUIDUM (*fluid extract of lobelia*), dose as an *expectorant* $\text{m}\text{j}-\text{v}$ (0.06–0.3); as an *emetic* $\text{m}\text{xxv}-\text{xx}$ (1.0–1.2).

TINCTURA LOBELIÆ (*tincture of lobelia*), contains 20 per cent. of lobelia; dose as an *expectorant* $\text{m}\text{x}-\text{xx}$ (0.6–1.2); as a *motor-depressant* in *asthma* $\text{f}\text{3j}$ (4.0) repeated every $\frac{1}{2}$ –1 hour until nausea is experienced.

CONIUM.

What is conium, and what active principles does it contain?

Conium is the *full-grown fruit* of *C. maculatum*, or hemlock (*Nat. Ord. Umbelliferae*), gathered while yet green and carefully dried. The *active principle* is *conine*, a *volatile liquid alkaloid*. It also contains *conydrine*, *methyl-conine*, and *pseudoconydrine*.

What are its physiological effects?

Full medicinal doses cause great muscular weakness, first felt in the lower extremities, difficulty in keeping the eyelids raised, and disordered vision from paralysis of accommodation. Sometimes the pupils dilate, and sometimes there is nausea and vomiting. After a *poisonous amount* has been taken, these symptoms are intensified, and difficulty of speech, delirium or coma, paralysis, and death from asphyxia, preceded by convulsions of cerebral origin, occur. Conium has little if any action on the *cerebral hemispheres*, and but little is known of its influence on the *spinal cord*. It paralyzes the peripheral ends of *nerves*, acting much more powerfully on the motor than on sensory fibres. The effects on the *ocular apparatus* are due to peripheral oculo-motor paralysis. The *pulse* is at first accelerated (paralysis of the vagi), but afterward slowed (depression of cardiac-motor nerves), and the blood pressure, which was primarily lowered, is raised when symptoms of asphyxia take place. It is eliminated by the urine. The *treatment of poisoning* consists in evacuating the contents of the stomach, giving tannic acid as an antidote, maintaining artificial respiration and administering, as antagonists, atropine, and strychnine or picrotoxin.

What are the medicinal uses of conium?

Conium has been used as a *depresso-motor* in *whooping-cough*, *paralysis agitans*, *chorea*, *blepharospasm*, *asthma* and other spasmodic affections. It has been used as a *calmative* and *relaxant* in *maniacal* and *hysterical excitement*, and also in *melancholia*. As an *anodyne* it is employed internally and locally in *cancer*, and in *neuralgia*, especially *sciatica*.

What are the preparations and doses of conium?

EXTRACTUM CONII (*extract of conium*), dose gr. j (0.06).

EXTRACTUM CONII FLUIDUM (*fluid extract of conium*), dose m_j-v (0.06–0.3).

As different samples of the drug vary much in the amount of conine contained, it is safer to begin with a small dose, to be gradually increased until some effect is obtained.

WOORARA.

What is woorara?

Woorara (not official), called also woorali and curare, is an extract prepared by the South American Indians from *Strychnos toxifera* and other species of *Strychnos* (*Nat. Ord. Loganiaceæ*), and is used by them as an arrow-poison. It contains an alkaloid called *curarine*.

What are its effects and uses?

It is eliminated so rapidly that usually when administered internally no symptoms result. When it enters the system through a wound or by hypodermic injection it paralyzes the peripheral endings of the *motor nerves*, and if the dose be sufficiently large the *vagus* and the ends of the *sensory nerves* are also paralyzed. After a time the *nerve trunks* and the *spinal cord* are also affected. The *voluntary muscles* are uninfluenced. Large amounts lower the *blood pressure* and finally depress the *heart*. Death results from paralysis of the motor respiratory nerves. Small doses increase all the secretions. It is *eliminated* chiefly by the urine.

The *treatment of woorara-poisoning* consists in frequent catheterization to prevent resorption, artificial respiration, and atropine and strychnine, hypodermically as antagonists. They can scarcely be relied on.

Woorara is *used* as a *depresso-motor* in *tetanus* and in *hydrophobia*, and has also been employed in *epilepsy* and in *chorea*. It is given in doses of gr. $\frac{1}{10}$ (0.006), or its alkaloid, curarine, may be used in doses of gr. $\frac{1}{200}-\frac{1}{100}$ (0.0003–0.0006).

TABACUM—TOBACCO.

What is tobacco, and what does it contain?

Tobacco is the commercial, *dried leaves* of *Nicotina Tabacum* (*Nat. Ord. Solanaceæ*). It contains *nicotine*, a volatile *liquid alkaloid*; *nicotianin* (tobacco-camphor) and an empyreumatic oil.

What are its effects and uses?

To persons unaccustomed to its use, tobacco or nicotine is a depressant, nauseant and emetic, and causes dizziness and weakness. When a toxic dose is taken, purgation, diuresis, cramps or convulsions, followed by great muscular weakness and, finally, paralysis, a rapid, feeble and compressible pulse, collapse and death from paralysis of the respiratory muscles.

It first stimulates the *spinal* centres (thus causing convulsions), but probably finally paralyzes them. It first excites and then abolishes the functions of the peripheral endings of motor nerves, the nerve trunks being afterward affected, but has no influence on voluntary muscles. The *cardiac* action is first depressed from some unexplained cause, afterward much increased, probably from paralysis of the peripheral inhibitory nerves, and the blood pressure, which at first falls, is afterward elevated. Nicotine contracts the pupil, whether applied locally or administered internally.

Tobacco increases intestinal peristalsis, and is probably eliminated chiefly by the urine, which it greatly increases.

The *treatment of nicotine poisoning* consists in emetics or washing out the stomach, tannic acid and strychnine with artificial respiration.

Tobacco is not much *used* at present as a *depresso-motor*, but may be employed for this purpose in *tetanus* and in *strychnine poisoning*. In persons unaccustomed to its use, smoking a strong cigar may relieve an attack of *spasmodic asthma*. As a *local anodyne*, it has been used as a poultice to *painful ulcers*, as an ointment to inflamed *hemorrhoids*, and as a lotion to relieve *pruritus*. Great caution should be observed, as its local use has resulted in poisoning and death. There are no *official preparations*. Dose gr. j–iij (0.06–0.19), in infusion, which in poisoning by strychnine or in tetanus must be frequently repeated until some results are obtained.

(B) ON THE ORGANS OF CIRCULATION.**ORDER I.—CARDIAC STIMULANTS.****What are cardiac stimulants?**

They are medicines which are administered with a view of increasing the power of the heart or the force of the circulation.

Mention the remedies of this order.

Alcohol, ether, the ammonium preparations, amyl nitrite, nitroglycerin, atropine, digitalis, caffeine, adonidine, strophanthus, sparteine, strychnine, cimicifuga.

ALCOHOL.**What is alcohol?**

Alcohol (ethyl hydrate) results from the fermentation of substances containing grape sugar. It is a colorless, inflammable liquid, uniting in all proportions with water. *Alcohol* contains 94 per cent. by volume of absolute alcohol, which is not used in medicine.

What are the physiological effects of alcohol?

Locally applied it is an astringent. It evaporates rapidly and produces a feeling of cold, but if evaporation is prevented it acts as an irritant. When its vapor is *inhaled* it causes anaesthesia, coma and death. Taken *internally in small doses* it increases appetite and digestion, temporarily increases the force and frequency of the cardiac systole, dilates the vessels, thus causing a sensation of warmth, and diminishes the amount of urea and of carbonic acid eliminated. The *continued use* of small doses disorders the digestion and the hepatic functions, produces gastric catarrh and hyperplasia of the connective tissues, resulting in sclerosis of different viscera, and tends to the production of fatty degeneration of the walls of the vessels and various organs.

Large doses disorder digestion and cause the well-known alcoholic intoxication with muscular incoördination and weakness, hallucinations, lowered temperature, depression of the heart and reflexes, and stupor.

After toxic doses death from respiratory or cardiac failure may rapidly result, preceded by insensibility, stertorous breathing and coma, and occasionally by convulsions.

In small doses it stimulates the *cardiac* motor ganglia and the

heart muscle, followed, if a large dose be taken, by depression and paralysis of the cardiac muscle. Large doses also paralyze the vaso-motor system and dilate the vessels.

Small doses slightly increase the *temperature* of the body, but large doses reduce it by interfering with the oxygen-carrying function of the red corpuscles and with the nutritive process, and by dilating the cutaneous vessels. *Respiration* is first quickened, then slowed, and after fatal doses, paralyzed. It is a *cerebral* stimulant, but in toxic doses depresses the encephalic mass, the *spinal* reflex centres, the conductivity of both motor and sensory nerves, and finally the reflex functions of the medulla. It diminishes the *excretion* of nitrogenized waste, and is *eliminated* principally by the kidneys, lungs and skin, although a certain amount (more in proportion to the smallness of the dose) disappears in the system.

How should acute alcohol-narcosis be treated?

The contents of the stomach should be evacuated, ammonia or strychnine should be given, and the hot and cold douche alternately used.

What are the medicinal uses of alcohol?

As a *cardiac stimulant*, alcohol is used in *syncope* from exhaustion, loss of blood, surgical shock and other forms of fainting, and where *cardiac failure* is threatened, as in typhoid and typhus fevers.

As a *general stimulant* to the system in all *wasting diseases*, as *phthisis*, *scrofula* and prolonged *suppuration*, and in all low states of the system, as *typhoid* and *typhus fevers*, *diphtheria*, *pyæmia* and *septicæmia*. In these fevers a tolerance to alcohol is present, and it may be given in large quantities without causing intoxication. It not only aids the digestion of food in these conditions, but as very little of what is taken can be found in the excretions, it seems to be used by the system, and consequently may be classed *as a food* in these diseases. The dose in these diseases must be regulated by the pulse principally. When the pulse is quickened and more bounding, the face flushes, the temperature rises, and the breath smells strongly of alcohol, the patient is taking more than the system can use, and the dose should be decreased.

As an *antidote* to *snake bite* or other cardiac depressant, as *aconite* or *veratrum viride*, it is of great service. In poisoning by snake bite

it is given in frequently repeated doses almost *ad libitum*, conjoined with intravenous or, at least, hypodermic injection of ammonia.

To improve *digestion* and *nutrition*, it is useful in many chronic diseases, but the danger of forming the alcohol habit should always be borne in mind when it is thus prescribed.

Locally, alcohol is used as an *antiseptic* and *stimulant* dressing for wounds, to harden the cuticle when *bed sores* are imminent, and as a *tonic* and *cooling lotion*.

What are the official preparations of alcohol?

ALCOHOL ABSOLUTUM (*absolute alcohol*) is ethyl alcohol containing not more than 1 per cent., by weight, of water. It is not used medicinally.

ALCOHOL DEODORATUM (*deodorized alcohol*) is a liquid composed of about 92.5 per cent. by weight, or 95.1 per cent. by volume, of ethyl alcohol and about 7.5 per cent. by weight of water.

It is used in making various medicinal preparations into which alcohol enters.

ALCOHOL (*alcohol*) contains 94 per cent. by volume of absolute alcohol; sp. gr. 0.820.

ALCOHOL DILUTUM (*diluted alcohol*) contains 48.6 per cent. by volume of absolute alcohol; sp. gr. 0.936.

SPIRITUS FRUMENTI (*whiskey*), an alcoholic liquid obtained by the distillation of the mash of fermented grain (usually of mixtures of corn, wheat, and rye), and at least 2 years old; it contains 50 to 58 per cent. by volume of alcohol; sp. gr. 0.917–0.930).

SPIRITUS VINI GALlici (*brandy*) an alcoholic liquid obtained by the distillation of fermented, unmodified juice of fresh grapes, and at least 4 years old; it contains from 46 to 55 per cent. by volume of alcohol; sp. gr. 0.925–0.941. It contains some tannic acid, and is astringent. It is, therefore, preferable to whiskey if there is looseness of the bowels. It also is a stomachic sedative, and, next to champagne, tends to check vomiting more than any other alcoholic liquid.

VINUM ALBUM (*white wine*), an alcoholic liquid, made by fermenting the juice of fresh grapes, the fruit of *Vitis vinifera* (*Nat. Ord. Vitaceæ*), freed from seeds, stems, and skins, contains 10 to 14 per cent. by weight of absolute alcohol.

VINUM RUBRUM (*red wine*), an alcoholic liquid made by ferment-

ing the juice of fresh colored grapes, the fruit of *Vitis vinifera* (*Nat. Ord. Vitaceæ*), in the presence of their skins, and contains from 10 to 14 per cent. of absolute alcohol. It is somewhat astringent.

How is alcohol administered?

Wines may be given pure or mixed with water. Brandy and whiskey are given pure or diluted with water, especially when administered as cardiac stimulants, or in the form of milk punch or egg-nog, when used as a food to support the system.

They may also be injected per rectum, and, if cardiac failure is imminent, brandy may be used hypodermically.

ÆTHER—ETHER.

Is this substance ever used as a cardiac stimulant?

The use of *ether* as a cardiac stimulant has already been mentioned. It acts rapidly, but transiently, and may be used internally, or better still, hypodermically in cases of sudden cardiac failure.

AMMONIUM.

What are the sources of ammonium?

Ammonium is abundantly present in nature as the result of animal and vegetable decay, but is chiefly obtained for commercial purposes from the ammoniacal liquor of gas works.

What are the general physiological actions of ammonium preparations?

AMMONIA GAS in solution is an irritant. Kept in contact with the skin it reddens and finally vesicates. When the gas is *inhaled* it irritates, producing a sense of suffocation, spasm, and, if continued, inflammation of the air passages. When *injected* into the veins of animals in sufficient quantities, it damages the blood and causes convulsions, rapid respiration and death.

Ammonia temporarily stimulates the accelerator nerves of the heart, thus increasing the rapidity of its action. After intravenous injection, the *blood pressure* at first falls, but rapidly rises as the pulse becomes quicker. It aids in maintaining the fluidity of the *blood* by retaining the fibrogenous materials in solution. In toxic amounts, the oxygen-carrying function of the *red corpuscles* is interfered with. Intra-venous injections of ammonia quicken *respiration*

by powerfully stimulating the respiratory centres. The convulsions are due to stimulation of the *motor and reflex spinal centres*. It is *eliminated* chiefly by the urine as uric acid and, perhaps, as urea. Its salts are also eliminated by the broncho-pulmonary mucous membrane. When taken internally in medicinal doses it is antacid and carminative, but if continued for too long a period, it disorders digestion and causes emaciation.

What are the symptoms and treatment of poisoning by ammonia?

When a strong solution is swallowed it causes inflammation of the lips, mouth, fauces, pharynx, œsophagus and stomach. Œdema of the glottis may occur, and secondary strictures of the œsophagus or even of the pylorus may result. Death may be due to asphyxia or to collapse preceded by symptoms of corrosive poisoning. The *treatment* consists in neutralizing the alkali by vegetable acids (vinegar); protecting the injured surfaces by oils and demulcents and the occasional passage of an œsophageal bougie, for at least a year, to prevent secondary stricture. The occurrence of œdema of the glottis demands immediate tracheotomy.

What are the therapeutical uses of the ammonium preparations?

Ammonia is a rapid and powerful *cardiac stimulant*, but its effects are of brief duration. It is used for this purpose both internally and by inhalation in all cases of acute *cardiac failure*, and may be used to antagonize the action of *snake poison* and *prussic acid* on the heart, but should here be employed as an adjuvant to alcohol.

For *sudden cardiac failure* during *anæsthesia* or from *surgical shock* it may be given hypodermically.

It has been used by intra-venous injection to *maintain the fluidity of the blood* in cases of threatened *cardiac thrombosis*; and as a *stimulating expectorant* in *pneumonia*, especially when typhoid symptoms are present. It is given *as an alkali* in *acidity*, *flatulence* and in *nervous headache*.

Externally it is an ingredient of stimulating and irritating liniments and may be used as a vesicant.

What preparations of ammonium are used as cardiac stimulants?

AQUA AMMONIÆ FORTIOR (*stronger ammonia water*) contains 28 per cent. by weight of gaseous ammonia. It is a powerful corrosive poison. It is best adapted for hypodermic use in doses of $\mathfrak{M}\text{x-xxv}$ (0.6–1.5), diluted with four times its volume of water, and repeated in 15 minutes if necessary, and may be used externally as a rapid (but painful) vesicant by inverting a watch crystal full of it on the skin.

AQUA AMMONIÆ (*ammonia water*) contains 10 per cent. by weight of ammonia-gas; *dose* for internal use, gtt. x-xxx (0.6–2.0) freely diluted.

SPIRITUS AMMONIÆ (*spirit of ammonia*) contains 10 per cent. by weight of the gas in solution in alcohol; *dose* $\mathfrak{M}\text{x-xxx}$ (0.6–2.0) freely diluted.

SPIRITUS AMMONIÆ AROMATICUS (*aromatic spirit of ammonia*) contains also ammonium carbonate and aromatics; *dose* $\text{f}\mathfrak{3}\text{ss-j}$ (2.0–4.0), freely diluted.

AMMONIÏ CARBONAS (*ammonium carbonate*) is used as a cardiac stimulant, and also in threatened thrombosis. It is very valuable as a stimulating expectorant in capillary bronchitis, typhoid pneumonia, and other low forms of chronic pulmonary disease where an expectorant is required. *Dose* gr. v-x (0.3–0.6).

The *iodide*, *bromide* and the *acetate*, *tartrate*, and *citrate of iron* and *ammonium*, and *valerianate of ammonium* have already been noticed; the *acetate*, *benzoate* and *chloride* will be discussed hereafter. The *nitrate of ammonium* is only used in making other official preparations.

AMYL NITRIS—AMYL NITRITE— NITROGLYCERIN.

Are these substances used as cardiac stimulants?

Inhalations of amyl nitrite have been employed in cases of *sudden heart-failure*, as in *fatty heart*, or *during anæsthesia*. *Nitroglycerin* has been used also as a cardiac stimulant, especially in cases associated with high arterial tension. These remedies have been discussed under the head of Depresso-Motors.

ATROPINÆ—ATROPINE.

Is atropine used as a cardiac stimulant?

Atropine (*vid. belladonna*, p. 107) is used as a cardiac stimulant in cases of sudden or threatened heart-failure. Dose gr. $\frac{1}{20}$ – $\frac{1}{60}$ (0.0005–0.001), given hypodermically.

DIGITALIS.

What is digitalis?

Digitalis is the leaves of *D. purpurea*, or purple foxglove (*Nat. Ord. Scrophularinæ*), obtained from plants of the second year's growth. It is a European plant, cultivated in this country.

What are its chemical constituents?

The constituents of digitalis* are, *digitalin*, *digitalosmin* (stearopten), *digitaleic acid*, *resin*, *mucilage*, etc. "Commercial digitalin is a mixture of several compounds of which the following have been investigated to some extent:" *Digitoxin*; *Digitonin*, a crystallizable glucoside, is decomposed by warm hydrochloric acid into dextrose, galactose and digitogenin; *Digitalein*; *Digitalin*, and *Digitin*. All these principles except the last are cardiac poisons.

What are the physiological effects of digitalis?

Its effects are much more slowly produced, but are much more lasting than those of the medicines previously considered, and it is therefore more of a tonic than a stimulant to the heart. In moderate doses it slows and strengthens the cardiac beat, elevates the blood pressure and causes diuresis. It sometimes interferes with the appetite and digestion. After very large doses, or when too long continued, nausea and vomiting may occur; the pulse becomes dicrotic, weak, and sometimes rapid, and diaphoresis more or less profuse takes place. Toxic doses cause the same symptoms, and in addition vertigo, disordered vision, lowered reflexes, muscular tremors, lassitude, reduced temperature, delirium, stupor, usually convulsions, and death from cardiac paralysis.

In moderate doses it stimulates the peripheral cardio-inhibitory fibres of the vagi (thus slowing the heart by increasing the length of

* A Manual of Organic Materia Medica, by John M. Maisch, Ph. M., Pharm. D., 6th ed., 1895, pp. 228, 229.

the diastole), the cardiac muscle and its contained motor ganglia (thus increasing the strength and force of the systole), and the vaso-motor centres (thus raising the arterial tension). It also acts directly on the muscular walls of the vessels, and thus lessens their calibre.

When full medicinal doses are taken, or when the pulse has been greatly slowed by it, a sudden change from the recumbent to the erect posture may give rise to a small, weak, rapid pulse, with lowered arterial tension.

When *fatal doses* are taken the heart is arrested in systole.

In poisonous doses it lowers *reflex action* by stimulating Setschenow's inhibitory reflex centres and, later, by paralyzing directly the spinal cord. It also lowers the contractility of *muscles*. It usually proves *diuretic*, probably from elevating the arterial tension. It is *eliminated* chiefly by the urine.

How should poisoning by digitalis be treated?

(1) Give emetics and cathartics to remove the poison from the alimentary canal; (2) tannic acid as a chemical antidote; alcohol, ammonium carbonate and opium, or saponin, as physiological antagonists.

What are the medicinal uses of digitalis?

As a *cardiac tonic* digitalis is used in all cases of loss of cardiac power, as in *dilatation* or in *weakness of the cardiac muscle*. It *should never be prescribed* in simple hypertrophy, nor when hypertrophy is sufficient to compensate for valvular lesion. In advanced cases of fatty heart, it should be used very cautiously, if at all, as the increased systolic power which it induces may lead to rupture of the friable cardiac walls. In *valvular diseases*, when the hypertrophy is not compensatory, or when dilatation is taking place, it is invaluable. In "*irritable heart*," *functional palpitation* and *venous engorgements* (by lengthening the diastole it allows the veins more time to empty their contents into the auricles), it is much used. The tincture (Mx-xx (0.6-1.2)) may be employed hypodermically in cases of *sudden cardiac failure*. To sustain the action of a depressed heart in various acute diseases, particularly in *pneumonia*, it is often administered.

In *delirium tremens*, where the pulse is soft and compressible, digitalis is often given in large doses with benefit.

It is also used as a *physiological antidote* to poisoning by *aconite*, *delphinine* and *muscarine*, all of which arrest the heart in diastole.

To *contract the blood vessels* it has been used in *acute inflammations* and in *hemorrhages*, and as it is supposed to stimulate involuntary muscle-fibres generally, it is given in *menorrhagia* and in *spermatorrhœa*.

As a *cardiac tonic* and *diuretic* it is used in all forms of *cardiac* and of *renal dropsy*, in *acute suppression of urine* and in *acute desquamative nephritis*.

What are the preparations of digitalis, and how are they administered?

Digitalis leaves are sometimes given in powder or pill, dose gr. j (0.06) t. i. d., gradually increased. The official preparations are—

EXTRACTUM DIGITALIS (*extract of digitalis*), dose gr. $\frac{1}{4}$ (0.016), gradually increased.

EXTRACTUM DIGITALIS FLUIDUM (*fluid extract of digitalis*), dose ℥j (0.06).

TINCTURA DIGITALIS (*tincture of digitalis*), dose ℥v-x (0.3-0.6), gradually increased.

INFUSUM DIGITALIS (*infusion of digitalis*), one of the best preparations; dose f3ss-jv (2.0-15.0).

Occasionally after digitalis has been taken for some time continuously, especially when free diuresis has not taken place, its action becomes suddenly manifested by the pulse rapidly falling and the supervention of other effects characteristic of the drug. This is probably due to some delay in the absorption or elimination of the medicine, and when it occurs the remedy should be promptly discontinued. So-called *commercial digitalin* should never be prescribed because of its variable composition. *Merck's German digitalin* is a stable and valuable remedy, representing the full value of digitalis, without its irritating effects upon the digestive tract. Dose gr. $\frac{1}{10}$ - $\frac{1}{2}$ (0.006-0.03).

CAFFEINA—CAFFEINE.

What is caffeine?

Caffeine (theine) is a feebly basic, *proximate principle*, obtained from the *dried leaves* of *Thea sinensis* or tea plant (*Nat. Ord.* Ternstræmiaceæ), from the *dried seeds* of *Coffea arabica* or coffee plant (*Nat. Ord.* Rubiaceæ), or from *guarana* (a paste prepared from the crushed or ground seeds of *Paullinia cupana* (*Nat. Ord.* Sapindaceæ). *Guarana* is official.)

What are its physiological actions?

In *medicinal doses*, caffeine is a cerebral stimulant, causing nervous restlessness, increased mental activity and wakefulness; it also increases the cardiac force, raises the arterial pressure, and acts as a decided diuretic. After *poisonous amounts*, there is also confusion of thought, with delirium, muscular tremblings, cardiac oppression, quickened respiration, cold extremities, symptoms of collapse, rise of temperature, and (in animals) tonic and clonic convulsions, muscular weakness, general paresis, and death from paralysis of respiration.

It increases the power of the *heart* and raises *arterial tension*, probably by a direct stimulant effect upon the heart muscle. It is *diuretic*, not only indirectly by increasing the blood pressure, but also by some direct action on the secreting cells of the kidney itself. The *convulsions* are probable due to spinal stimulation. In moderate doses, caffeine is said to be entirely destroyed in the system; in toxic doses, it is partly destroyed and partly eliminated by the kidneys. Its effect on the elimination of waste products is at present uncertain.

What are its medicinal uses?

As a *cardiac tonic* (more rapid in action, but not so enduring as digitalis), it is used to meet the same indications which call for the use of digitalis. In cases of *sudden heart failure*, it may be used hypodermically combined with equal parts of sodium salicylate, which renders it more soluble in water.

As a combined *cardiac stimulant* and *diuretic* it is very useful in the latter stages of *chronic Bright's disease* with weak heart.

As a *diuretic* in all *renal* and *cardiac dropsies* it is very serviceable. It is also employed as a *cerebral stimulant* in nervous and neuralgic *headaches*, and combined with other antagonists in opium poisoning.

How is caffeine administered?

CAFFEINA (*Caffeine*) may be given in pill, capsule or solution, dose gr. j-v (0.06-0.3) or more. For administration in fluid form *caffeine citrate* (so-called) is preferred as more soluble.

When given as a *cerebral stimulant*, one dose of gr. ij-v (0.13-0.3) is usually sufficient, but as a *cardiac tonic* and *diuretic* this dose should be repeated twice daily, and gradually increased if necessary up to gr. x-xij (0.6-0.8).

CAFFEINA CITRATA (*citrated caffeine*), dose gr. ij-v (0.13-0.3).

CAFFEINA CITRATA EFFERVESCENS (*effervescent citrated caffeine*), dose ʒj-iiij (4.0-12.0).

GUARANA (*Guarana*) contains *tannic acid* as well as *caffeine* in considerable quantities, and is principally used in *neuralgic* and in *sick-headache*. In Brazil, whence it comes, it is also used as an astringent in diarrhoeal affections, dose ʒj (4.0).

EXTRACTUM GUARANÆ FLUIDUM (*fluid extract of guarana*), dose m℥x-xx (0.6-1.2).

ADONIDIN.

What is adonidin?

Adonidin (not official) is a glucoside, obtained from the root of *Adonis vernalis* (*Nat. Ord.* Ranunculaceæ), a plant of northern and central Europe.

What are its effects, uses and modes of administration?

The *effects* of adonidin are almost identical with those of *digitalis* (but are much more quickly produced), as it stimulates the cardiac muscle and contained motor ganglia, thus increasing the force of the cardiac contractions at the same time that, by stimulating the inhibitory fibres of the vagi, it increases the length of the diastole and thus slows the heart. The arterial tension is also elevated, probably by stimulation of the vaso-motor centres. In large doses it paralyzes both the heart and vessels, the arterial tension falls and the heart is finally arrested in diastole. It sometimes (not as frequently as does *digitalis*) causes nausea, vomiting and diarrhoea. It sometimes proves diuretic. It is *used* as a *cardiac tonic* in the same class of diseases in which *digitalis* is employed, and may be substituted for it where the latter disagrees or is too slow in its action.

It may be alternated with digitalis for a few weeks during a course of the latter. Dose gr. $\frac{1}{8}$ – $\frac{1}{2}$ (0.008–0.03) every 3 or 4 hours, in pill or capsule.

STROPHANTHUS.

What is strophanthus?

Strophanthus is the seed of *S. hispidus* (Nat. Ord. Apocynaceæ), an African plant which has been used by the natives as an arrow-poison, and called by them Kombé or Inéc. It contains a crystalline glucoside, called *strophanthin*.

What are its effects, uses and modes of administration?

The effects of *strophanthus* are more rapid in action, but much more evanescent, than those of digitalis. It appears to act upon the muscles directly, and to have little or no effect upon the nervous system.

In moderate doses it greatly increases the strength of the cardiac systole, prolongs the diastole and increases the arterial tension probably by acting on the muscular walls of the vessels. In large doses it increases the contractibility of all the striated muscles and finally arrests the heart in systole.

It proves diuretic by increasing the arterial tension, and has no cumulative effects. It is used as a substitute for digitalis in cardiac diseases when a temporary effect or a temporary change of medicines is desirable. From its diuretic action it has been successfully employed in various forms of œdema depending on a weak circulation and general cardiac and renal dropsies. The tincture of *strophanthus* (*tinctura strophanthi*—5 per cent. by weight) is used; dose ℥j–x (0.06–0.6) t. i. d.; or *strophanthin* may be cautiously administered hypodermically; dose gr. $\frac{1}{100}$ – $\frac{1}{60}$ (0.0006–0.001).

SPARTEINÆ SULPHAS—SPARTEINE SULPHATE.

What is sparteine?

Sparteine is a liquid alkaloid obtained from *Cytisus Scoparius*, or broom (Nat. Ord. Leguminosæ).

What are its effects and uses, and how is it administered?

Sparteine is a cardiac tonic, in moderate doses increasing the force and persistence of the cardiac contractions by direct action upon the

heart or its contained motor ganglia, and at first increasing its frequency by paralyzing the vagi, afterward decreasing the pulse-rate by its direct action on the heart. When the cardiac rhythm is disturbed it renders its contractions regular. In *large doses* it depresses the heart and causes death from paralysis of the respiratory centre. It depresses the cerebral and spinal centres. It has no diuretic properties. It has been *used* as a substitute for digitalis in *cardiac diseases*, and in cases of feebleness and irregularity of cardiac action. It is inferior to digitalis as a cardiac tonic, but may be used as a succedaneum to it. It is administered in the form of *sulphate* (official), of which the dose is gr. $\frac{1}{20}$ – $\frac{1}{4}$ (0.003–0.01), hypodermically; or if given internally, gr. $\frac{1}{4}$ –j or ij (0.01–0.06–0.13) in capsule or pill.

STRYCHNINA—STRYCHNINE.

Is strychnine prescribed for its cardiac effects?

The use of strychnine as a cardiac tonic has been noticed under the head of *nux vomica* (*q. v.*); dose gr. $\frac{1}{30}$ – $\frac{1}{20}$ (0.002–0.003) internally, or gr. $\frac{1}{60}$ (0.001) if used hypodermically.

CIMICIFUGA.

What is cimicifuga?

Cimicifuga is the *rhizome* and *roots* of *C. racemosa*, black snake-root or cohosh (*Nat. Ord. Ranunculaceæ*), a common indigenous plant. It contains a *volatile oil*, to which its virtues are probably due. A crystalline principle has also recently been obtained from it. Its *effects* on the circulation are supposed to resemble those of digitalis, while it acts on involuntary muscular fibres similarly to ergot. Recently it has been said, however, that it is a direct depressant to the cardiac muscle or its ganglia. It depresses the reflexes and occasionally causes nausea and vomiting. Its action is mild and not well understood. It is also diuretic and expectorant. It has been *used* as a *cardiac tonic* in fatty heart, as a *diaphoretic* in *rheumatism*, as a *diuretic* in *dropsies*, as an expectorant in *chronic bronchitis*, and as an *antispasmodic* in *chorea*, in which disease it is sometimes of value, combined with tonics and laxatives. It is not often prescribed.

EXTRACTUM CIMICIFUGÆ FLUIDUM (*fluid extract of cimicifuga*), dose fʒss–j (2.0–4.0).

TINCTURA CIMICIFUGÆ (*tincture of cimicifuga*), dose fʒss-ij (2.0-8.0).

EXTRACTUM CIMICIFUGÆ (*extract of cimicifuga*), dose gr. iij-v (0.2-0.3).

CONVALLARIA.

What is convallaria?

Convallaria is the *rhizome* and *roots* of *C. majalis*, or lily of the valley (*Nat. Ord. Liliaceæ*), a European plant naturalized in this country.

It contains two glucosides, *convallamarin* and *convallarin*, to the former of which it probably owes its action on the heart, while the latter appears to cause nausea, diarrhœa and gastric pain.

The results of the investigations regarding the *physiological action* of this plant are very contradictory, but it seems to slow the heart and increases the blood-pressure in small doses. In toxic doses the cardiac action becomes rapid, the arterial pressure gradually falls, and death occurs from heart-failure.

It is *used* in all cases of *cardiac weakness* whether accompanied by valvular lesion or not, as in *dilatation* or *fatty heart*. It is also used in *palpitation*.

EXTRACTUM CONVALLARIÆ FLUIDUM (*fluid extract of convallaria*), dose ℥v-xv (0.3-0.9).

ORDER II.—CARDIAC SEDATIVES.

What are cardiac sedatives?

Cardiac sedatives are medicines which are used to diminish the force and frequency of the cardiac contractions when they are morbidly increased, and thus to depress the circulation.

Name the principal cardiac sedatives.

Antimonials, aconite and its alkaloid, veratrum viride and its alkaloids, veratrine, gelsemium, pulsatilla, arnica, the vegetable acids.

ANTIMONIUM—ANTIMONY.

What are the sources of antimony?

Antimony is a metallic element found native in the form of black antimonious sulphide.

Mention some of the tests for antimony.

Sulphuretted hydrogen in solution forms with solutions of antimonial salts an orange-colored precipitate. The metal should also be reduced by Marsh's test, and chemically and microscopically distinguished from arsenic.

What are the general physiological effects of antimonials?

When applied *locally* to the skin, the salts of antimony produce an eruption of papules, which become pustules, somewhat resembling the eruption of smallpox. After prolonged use, the fauces sometimes present a similar eruption. *Internally*, in *small doses*, they are diaphoretic. In somewhat *larger doses*, they cause nausea, depressed circulation, weakness, and increase of the mucous and cutaneous secretions. In *large doses*, they prove emetic, with very weak pulse, great general depression and relaxation of the muscular system. In *poisonous doses*, they are gastro-intestinal irritants, causing emesis, catharsis with cramps in the extremities, and collapse, often resembling Asiatic cholera, and death from asthenia, sometimes preceded by tonic and clonic spasms.

Antimonials slow the *heart*, increase the duration of its diastole and diminish the force of its systole by direct depression of the cardiac functions; they lower the blood pressure partly by cardiac depression and partly by paralyzing the peripheral vaso-motor system. The pulse finally becomes rapid and very feeble, the pressure sinks almost to *nil* and the heart is arrested in diastole. They depress and paralyze the reflex, sensory, and, to a less extent, the motor functions of the *spinal cord*. The *temperature* is reduced by poisonous doses. The *emesis* is probably partly from local and partly from centric irritation.

When *given for a length of time*, they cause fatty degeneration of various viscera.

Antimony is *eliminated* by the mucous membranes, especially of the stomach, by the liver, kidneys and skin.

What is the treatment of poisoning by antimony?

Wash out the stomach to remove any poison present; give tannic acid as a chemical antidote, and opium and stimulants to overcome the resulting depression.

What are the medicinal uses of antimonials?

They are used as *cardiac sedatives* in the early stages of *acute sthenic inflammations*, but are probably inferior to aconite or veratrum viride for this purpose. They are very useful in the early stages of *acute bronchitis* and in *acute laryngitis* from their combined sedative, diaphoretic and expectorant qualities. As *diaphoretics* they are used in minute doses, frequently repeated, in fever mixtures, or combined with minute amounts of morphine, which increases their action. The depression which they produce should always be remembered.

As *nauseants* and *emetics* they are rarely resorted to, because of the great depression which they produce.

Sulphurated antimony, or the old Kermes mineral, is sometimes used as a nauseant in membranous croup.

Locally, as a *counter-irritant* it is rarely employed.

What are the antimonial preparations?

ANTIMONII ET POTASSII TARTRAS (*antimonium and potassium tartrate—tartar emetic—tartarated antimony*); dose, as a *sedative*, gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.01–0.03), increased to gr. j–ij (0.06–0.12), as tolerance to its action is established; as a *diaphoretic* and *expectorant*, gr. $\frac{1}{16}$ – $\frac{1}{4}$ (0.004–0.016), in solution, repeated every 2 or 3 hours until the desired results are produced; as an *emetic*, gr. j–ij (0.06–0.13).

VINUM ANTIMONII (*wine of antimony*) contains 4 parts of tartar emetic in 1000 of the preparation (about gr. ij–f $\overline{3}$ j). Dose as a *diaphoretic* and *expectorant* ℥x–xxx (0.6–2.0); as an *emetic* f $\overline{3}$ j–iv (4.0–15.0).

SYRUPUS SCILLÆ COMPOSITUS (*compound syrup of squill—Coxe's hive syrup*) contains tartar emetic, 2 parts in 1000 of the preparation (about gr. j to the f $\overline{3}$ j). It is used in croup as an *emetic* and also as an *expectorant*; dose ℥v–f $\overline{3}$ j (0.3–4.0).

The following preparations are somewhat uncertain, and are rarely employed:—

ANTIMONII OXIDUM (*antimonious oxide*), dose gr. j–ij (0.06–0.13).

PULVIS ANTIMONIALIS (*antimonial powder—James' powder*) contains antimonious oxide 33 per cent. with precipitated calcium phosphate; dose gr. iij–x (0.2–0.6).

ANTIMONII SULPHIDUM (*antimonious sulphide*) and ANTIMONII SULPHIDUM PURIFICATUM (*purified antimonious sulphide*) are used in making the other preparations.

ANTIMONII SULPHURATUM (*sulphurated antimony—Kermes mineral*), dose gr. j–iij (0.06–0.2); as an emetic gr. v–x (0.3–0.6).

PILULÆ ANTIMONII COMPOSITÆ (*compound pills of antimony—Plummer's pill*). Used as an alterative. Each pill contains of calomel and sulphurated antimony each gr. $\frac{6}{10}$ (0.04).

ACONITUM—ACONITE.

What is aconite, and what is its chemical composition?

Aconite is the *tuber* of *Aconitum Napellus*, aconite, monkshood or wolfsbane (*Nat. Ord. Ranunculaceæ*), a native of the mountains of Europe and Asia. The active principle is an alkaloid called *aconitine*. Other alkaloids have been discovered in it, viz.: *aconine*, *isaconitine*, and *picraconitine*.

What are its physiological effects?

Locally applied, aconite or aconitine benumbs the sensory nerves and causes a sensation of tingling. *Internally* in *small doses* it lessens the force and frequency of the cardiac contractions, diminishes the blood pressure, causes tingling in the lips, tongue and extremities and produces a feeling of weakness. When *large doses* are taken, these symptoms are more marked, the pulse becomes very weak and slow, a tendency to fainting is manifested on assuming the erect posture or on any exertion, and the respirations are reduced in number. After *poisonous* doses these symptoms are much intensified; the countenance is anxious, a cold sweat covers the body, the pulse and respiration are very slow, weak and irregular, general anæsthesia is present, and collapse and death from syncope, sometimes preceded by convulsions, occur.

Aconite affects the *heart* by directly depressing the cardiac muscle, and its contained motor ganglia. It probably also stimulates the cardio-inhibitory apparatus, but does not affect the vaso-motor nerves or centres. It paralyzes the peripheral end organs of both *motor* and *sensory nerves*, the latter being first affected by it, and the nerve trunks and centres being only involved as the poisoning deepens. The loss of reflex function is probably due to interference

with conduction. It depresses and finally paralyzes the *respiratory centre*.

What is the treatment of poisoning by aconite?

Evacuate the contents of the stomach and wash it out by means of the stomach pump; keep the sufferer perfectly quiet, with his head low; give cardiac stimulants, as alcohol, ether or ammonia, and administer digitalis hypodermically.

What are the medicinal uses of aconite?

As a *cardiac sedative* aconite is of great value in the early stages of all *acute inflammations* and all *fevers*, in which the patient is vigorous and young and the pulse tense, full, bounding and strong, and is only *contraindicated* when the heart is weak, when there is gastro-enteric inflammation, or where a typhoid condition is present. Thus it is used to diminish the circulation, arterial tension and temperature, and to promote diaphoresis and diuresis in the sthenic forms of *surgical fever*, in *acute coryza*, *acute broncho-pulmonary catarrhs*, *acute pleuritis* and *acute peritonitis*, in *acute meningitis* either cerebral or spinal, and in the *essential fevers*, when not of the asthenic type. In *scarlet fever* and in *measles* (when the temperature is high), it is of great value, and is often used during the exacerbations of remittent malarial fevers. In *lobar pneumonia veratrum viride* is a safer remedy, but aconite will prove very serviceable if employed with great caution before consolidation has taken place.

In *cardiac hypertrophy* uncomplicated by valvular lesion, or even if the latter is present, provided the hypertrophy be in excess of the requirements of the system, it is an invaluable medicine.

As a *local anæsthetic* it has been advantageously given internally and applied externally in *neuralgia*, particularly of the face and head, and has been used locally in *chronic muscular rheumatism* and in *rheumatic neuralgie*.

What are the preparations of aconite and their doses?

EXTRACTUM ACONITI (*extract of aconite*), dose gr. $\frac{1}{8}$ -ss (0.008-0.03).

EXTRACTUM ACONITI FLUIDUM (*fluid extract of aconite*), dose m_{ss} -ij (0.03-0.123).

TINCTURA ACONITI (*tincture of aconite*), dose m_{j} -v (0.06-0.3), frequently combined with other medicines in fever mixtures.

In fevers and inflammations aconite is preferably given in small doses, repeated every few hours, and carefully watched, until its effects on the pulse are obtained.

Aconitine is not official and should not be used internally, as even in the minutest doses it is a powerful poison. Duquesnel's aconitine has, however, been administered with great care in *tic douloureux* with success. Dose gr. $\frac{1}{300}$ – $\frac{1}{200}$ (0.0002–0.0003). Morson's aconite is probably even more powerful. Aconitine is used *locally* as an anodyne in neuralgia, in the form of liniment (gr. ij (0.13)–f3j (4.0)) or as an addendum to ointments.

VERATRUM VIRIDE.

What is *veratrum viride*, and what does it contain?

It is the *rhizome and roots* of *Veratrum viride*, American or swamp hellebore or Indian poke (*Nat. Ord.* Liliaceæ), a plant indigenous to swampy places in the eastern part of the United States. It contains two alkaloids, *veratroidine* and *jervine*, to which its effects are due; also *rubijervine*, *pseudojervine*, resin, etc.

What are their physiological effects?

Jervine taken *internally* causes great weakness, depresses the spinal cord and lowers reflex action, lessens the force and frequency of the heart by direct action on the cardiac muscle or its contained motor ganglia, greatly reduces the blood pressure, both by its action on the heart and by depressing and finally paralyzing the vaso-motor centres, causes profuse salivation, and death from asphyxia, preceded by convulsions of cerebral origin.

Veratroidine gives rise to the same general effects, but in addition always produces vomiting and often purging. When directly applied to the heart muscle, it paralyzes it. Given internally, it first slows the heart and lowers the blood pressure by stimulating the vagus, then, the pulse remaining slow, the heart-beats increase in force and the blood pressure rises to normal, but suddenly, from peripheral paralysis of the vagi, the cardiac action becomes very rapid and diminishes in force, and, as asphyxia increases, the arterial tension is much elevated. Death is due to paralysis of the respiratory muscles.

Veratrum viride combines the effects of its alkaloids, and is a powerful arterial and spinal depressant, diminishing the frequency

and force of the cardiac contractions, by depressing the heart muscle and stimulating inhibition, and lowering the blood pressure by paralyzing the vaso-motor system. It often causes vomiting. It is a diaphoretic, reduces the temperature, and sometimes proves diuretic. Poisoning rarely occurs, as an overdose causes emesis. Should dangerous symptoms arise, they should be treated with alcoholic and other cardiac stimulants.

What are the medicinal uses of *veratrum viride*?

It is given as a *cardiac sedative* in the same *inflammatory* and *cardiac affections* in which aconite is administered, and its *contra-indications* are identical with those of the latter drug, with the addition that in peritonitis it is not a safe remedy, because of its liability to cause vomiting. In *acute lobar pneumonia* it is considered much safer than aconite. In *acute mania* and in *mania-a-potu* (when the patient is strong and the pulse full and bounding), it is a valuable medicine, but should never be given in delirium tremens. It has been highly recommended in *puerperal eclampsia*, in doses of $\text{m}_{\text{x-xx}}$ (0.6–1.2) of the tincture, given hypodermically, partly as an arterial sedative and partly as a reflex depressant.

What are its preparations and their doses?

EXTRACTUM VERATRI VIRIDIS FLUIDUM (*fluid extract of veratrum viride*), dose $\text{m}_{\text{j-iv}}$ (0.06–0.24).

TINCTURA VERATRI VIRIDIS (*tincture of veratrum viride*), $\text{m}_{\text{ij-x}}$ (0.123–0.6).

Its effects are better obtained by giving small and increasing doses at intervals of an hour, being careful not to cause vomiting, and closely watching the pulse.

VERATRINA—VERATRINE.

What is veratrine?

Veratrine is a *mixture of alkaloids* obtained from the seed of *Asagræa officinalis* (*Nat. Ord. Liliaceæ*), a Mexican plant.

How may its presence be detected?

When warmed in concentrated hydrochloric acid a persistent dark-red color is produced, resembling a strong solution of potassium permanganate. This is *Trapp's test for veratrine*.

What are its physiological actions?

Locally veratrine is irritant, causing heat and pain, followed by numbness, and if long applied, redness of the surface. After *small doses* the pulse is weakened and reduced in frequency. *Large doses* are followed by vomiting and purging and muscular weakness; with cold, clammy skin; weak, rapid, irregular pulse and muscular tremors. Veratrine appears to stimulate the excito-motor *cardiac ganglia and muscle* (temporarily increasing the pulse), but soon depresses, and finally paralyzes them. It excites and then destroys the function of the vagi (thus lessening the pulse-rate, which, however, in poisoning, become rapid again as the vagi are paralyzed). The *vaso-motor centres* are first stimulated, then depressed. It first stimulates and then depresses the nerve trunks (or their peripheral end-organs) and the muscles, without apparently affecting the cerebrum or spinal cord. It depresses and finally paralyzes the respiratory centre, death occurring from this cause. The temperature is also lowered. It is eliminated chiefly by the skin and kidneys, increasing their secretion.

No fatal case of poisoning is recorded; the treatment is the same as that employed in poisoning by *veratrum viride*.

What are its therapeutic uses and preparations?

Although it has been used as a cardiac depressant, yet as it is inferior in this respect to the remedies already considered, it is now but little given *internally*. It is chiefly used as an external application in *rheumatic pains* and in *neuralgia*.

UNGUENTUM VERATRINÆ (*ointment of veratrine*) contains 4 per cent. of veratrine.

OLEATUM VERATRINÆ (*oleate of veratrine*) contains 2 per cent. of veratrine.

GELSEMIUM.

Is gelsemium used as a cardiac sedative?

It has been employed for this purpose in sthenic inflammation and fevers. Its medicinal uses have been described among the depresso-motors.

PULSATILLA.

What is pulsatilla?

It is the *herb* of *Anemone pulsatilla* and *A. pratensis* (*Nat. Ord.* Ranunculaceæ), and contains a volatile oily substance which when kept decomposes, forming *anemonin* and *anemonic acid*.

What are its effects and uses?

Locally pulsatilla is irritant; *internally* it depresses the circulation and lowers the blood-pressure, and in poisonous doses causes vomiting and purging, dilated pupils, convulsions, paralysis, diminished sensation, coma and death. It is *eliminated* by the kidneys. Pulsatilla is not much *used*, as it is decidedly inferior to other cardiac sedatives. It may be given in *doses* of gr. j-v (0.065-0.324) in powder, pill, capsule or infusion, or a tincture or fluid extract may be prepared.

ARNICA.

What is arnica?

It is the *flower-heads* and the *rhizome and roots* of *Arnica montana*, leopard's bane (*Nat. Ord.* Compositæ). It contains *arnicin*, a *volatile oil*, *resins*, etc.

What are its effects and uses?

Locally applied arnica is a stimulant and often a decided irritant. *Internally* it slows the heart, but seems to raise the arterial pressure slightly. In large doses it is emetic and cathartic, causes great muscular weakness, a rapid pulse, and if the dose be excessive, collapse. It seems to be eliminated by the skin and kidneys. Arnica is not much *used* in this country except as an *external* application to sprains and contusions. When thus employed its irritant qualities should be borne in mind. The preparations are:—

EXTRACTUM ARNICÆ RADICIS (*extract of arnica root*), dose gr. v-x (0.324-0.648).

EXTRACTUM ARNICÆ RADICIS FLUIDUM (*fluid extract of arnica root*), dose ℥x-xx (0.6-1.2).

TINCTURA ARNICÆ RADICIS (*tincture of arnica root*) and TINCTURA ARNICÆ FLORUM (*tincture of arnica flowers*), dose ℥x-xxx (0.6-2.0).

EMPLASTRUM ARNICÆ (*plaster of arnica*).

THE VEGETABLE ACIDS.

Name the vegetable acids.

The officinal vegetable acids are : acetic, citric and tartaric acids.

What are their general effects and uses ?

When applied to an abraded surface they are irritant (tartaric acid being most, and citric the least, powerful). Given *internally* they increase the saliva and thus diminish thirst in fever. In *large doses* they are gastro-intestinal irritants and cardiac sedatives. They are probably oxidized in the blood and converted into carbonic acid, which by abstracting part of the base of the alkaline phosphates, converts them into acid phosphates, thus increasing the acidity of the urine. When taken in large quantities they cause flatulence, tormina and diarrhœa. Their long-continued use disorders the digestion, and causes emaciation and an anæmic condition. They are eliminated, chiefly as carbonates, by the intestines and kidneys.

The vegetable acids are *used* as *refrigerants* and perhaps cardiac sedatives in *fevers* (citric acid), as *antiscorbutic agents* (citric or acetic acid), in *acute rheumatism* (lemon juice is, however, inferior to the alkaline treatment), in *indigestion* depending on superacidity of gastric juice (lemon juice before meals), and as *diuretic* to increase the amount of urine, and render it acid in reaction (tartaric or citric acid).

They are also used as *chemical antidotes* to poisoning by the alkalies.

Locally acetic acid is used as a *caustic* to remove corns and warts, as a stimulating liniment in *sprains*, as an injection in *vaginal leucorrhœa*, and diluted with water it is given per rectum to *destroy ascarides*, and as a cooling lotion in fevers. The preparations are—

ACIDUM ACETICUM GLACIALE (*glacial acetic acid*) (nearly or quite absolute acetic acid) and ACIDUM ACETICUM (*acetic acid*) (containing 36 per cent. by weight of absolute acetic acid) are used externally as caustics.

ACIDUM ACETICUM DILUTUM (*diluted acetic acid*) may be used internally; dose f3j-ij (4.0-8.0) diluted. It is a pure form of vinegar.

ACIDUM CITRICUM (*citric acid*), 3j (31.0) to water Oj (480.0), may be used as a substitute for lemonade.

SYRUPUS ACIDI CITRICI (*syrup of citric acid*) contains 10 parts of the acid in the 1000 of the preparation.

LIMONIS SUCCUS (*lemon-juice*), OLEUM LIMONIS (*oil of lemons*), and SPIRITUS LIMONIS (*spirit of lemon—essence of lemon*) are also official.

ACIDUM TARTARICUM (*tartaric acid*) is rarely used in medicine, except as an ingredient of Seidlitz powders.

(C) ON THE SEXUAL ORGANS.

ORDER I.—APHRODISIACS.

What are aphrodisiacs?

Aphrodisiacs are medicines used to excite the functions of the genital organs when they are morbidly depressed.

What medicines are used for this purpose?

When the *nutritive functions* are not properly performed, the tonics or alteratives, especially phosphorus, zinc phosphide, the phosphates and hypophosphites, the salts of iron, gold or arsenic, and cod-liver oil are used. When there is an *absence of sexual desire*, small doses of opium, very minute amounts of alcohol, cannabis Indica, musk, or small doses of camphor may be useful by stimulating the cerebral centres. If impotence be due to *want of erectile power*, the tonics and alteratives are of use, as are also agents acting on the genital centre in the lumbar portion of the spinal cord and agents which either determine a flow of blood to the part by increasing the general or local circulation or which, by contracting the veins, aid in producing and maintaining erection. Thus, nux vomica or its alkaloid, the cardiac stimulants, hamamelis, ergot, cantharides and myrrh are useful in this condition. Good hygiene, cold baths, proper food and sea air and bathing, electricity, and moral means are important. The above medicines have either been considered already or will be found in their appropriate places.

ORDER II.—ANAPHRODISIACS.

What are anaphrodisiacs ?

Anaphrodisiacs are medicines which are used to depress the sexual functions when they are morbidly excited.

What medicines are employed for this purpose ?

Morbid excitation of the sexual organs is treated by medicines which depress the functions of the body generally, as nauseants and purgatives, or those which, by depressing the brain or spinal cord, will prove sedative to centres governing these organs : thus full doses of opium or of camphor will depress the cerebral centre, while potassium or ammonium bromide (very useful), monobromated camphor, iodine and the iodides, potassium nitrate, ammonium chloride, belladonna, tobacco, and conium act more especially on the spinal centre and nerves. Medicines diminishing the circulation of the parts will also prove advantageous, as the cardiac sedatives or full doses of digitalis and the bromides. Cocaine locally applied acts by contracting the vessels and diminishing the secretions of the sensory nerves.

Other agents used for this purpose are the local use of cold, general blood-letting, a low diet, avoidance of stimulation, and the withdrawal of blood to other parts of the body, as by hard mental work or exercise with the upper extremities.

ORDER III.—OXYTICS.

What are oxytics ?

Oxytics are medicines which are employed to increase the power of a contracting uterus, thus aiding in the expulsion of its contents.

Name the medicines belonging to this order.

Ergot ; hydrastis ; hydrastinine hydrochlorate ; gossypii radice cortex, and quinine.

ERGOTA—ERGOT.

What is ergot ?

Ergot is the *sclerotium* of *Claviceps purpurea* (*Nat. Ord. Fungi*), replacing the grain of *Secale cereale* (*Nat. Ord. Graminaceæ*), or, in other words, it is a fungus growing from the diseased ovary of rye. “When more than a year old it is unfit for use.”

What does it contain?

The exact chemical composition of ergot is still a matter of doubt, but the latest researches seem to show that it contains a resinoid substance, *sphacelinic acid*, and a neutral principle, *cornutin*, which together represent the value of the drug.

What are its physiological effects?

The effects of ergot are chiefly manifested on the uterus of the pregnant female and on the circulation. In the *non-pregnant*, *small doses* produce no perceptible effect; *moderate doses* slow the heart, probably by stimulating the peripheral fibres of the vagi, and enormously increase the blood pressure by stimulating the vaso-motor centres. *Large doses* cause nausea and vomiting, increased peristalsis and purging. *Toxic doses* depress the heart by acting directly on its muscle or contained ganglia, and lower the blood-pressure, partly by its action on the heart and partly from depression of the vaso-motor centres.

When *poisonous doses are given to animals*, there is salivation, vomiting and sometimes diarrhœa, rapid pulse and respiration, dilated pupil, trembling, great thirst, prostration, paraplegia, convulsions and death. The nervous symptoms appear to be due to action on the nerve centres.

Ergot possesses the property of causing contraction of all non-striated muscular fibres: thus it contracts the arterioles, increases intestinal peristalsis, and increases the power as well as the duration of the uterine contractions during labor, but does not always originate them during pregnancy or at other times, except immediately prior to and after parturition.

What is ergotism?

When ergot is used continuously in large amounts as an article of diet, it produces a condition of chronic poisoning, which manifests itself either by gangrene (usually dry, but occasionally moist and attended with septicæmia) or by tetanic spasms.

What are the medicinal uses of ergot?

Ergot is used *as an oxytocic in small doses* in *uterine inertia* during the second stage of labor, when there is no resistance either on the part of the bony canal or the soft parts of the mother. When thus administered, it energizes the uterine contractions, but if given in

large doses it renders them tetanic and continuous, thus greatly increasing the danger both to the mother and child.

To prevent *post-partum hemorrhage* by contracting the uterus, and thus compressing the uterine sinuses, it is invaluable, but had better be withheld until the placenta has left the uterine cavity, as prior to this it will tend to imprison the placenta by contracting the cervix. After the uterus is empty, it may be given in full doses, either by the mouth or hypodermically.

It has been used to *induce premature labor*, but is at best an uncertain remedy for this purpose. To arrest hemorrhage in cases of *threatened abortion*, ergot is worse than useless, unless the uterus is emptied. In these cases, by contracting the cervix, it imprisons the ovum, and thus, by preventing its escape, keeps the uterus distended. If given at all in these cases, its administration should be conjoined with the application of vaginal tampons. It is also given to aid the expulsion of *uterine polypi*.

As a *hæmostatic*, it is used in all *hemorrhages* where surgical means of arrest cannot be employed, and is often given as an adjuvant to their local treatment. In *menorrhagia*, *epistaxis*, *hemorrhage from the gums*, and in *broncho-pulmonary*, *intestinal* or *renal hemorrhage* it is very efficient. In *hæmatemesis*, it is not so serviceable. In *purpura*, it acts beneficially. From its influence in contracting blood vessels, it has been used in the early stages of *pneumonia* and other *inflammatory affections*, in *spinal congestions*, in *cerebro-spinal meningitis*, and for the cure of *internal aneurisms*. It has been injected hypodermically near the seat of disease for the cure of *varices*, *hemorrhoids*, and *impotence* due to dilatation of the dorsal vein of the penis. It may be employed with great advantage, either internally or hypodermically, to reduce the size of an *enlarged spleen*. In *glycosuria* and in *polyuria* it does good, and, from its influence on unstriated muscular fibres generally, it is useful in *relaxation of the sphincters*, *diarrhœa*, *dysentery*, *spermatorrhœa*, *uterine subinvolution*, and to diminish the size of *subperitoneal uterine fibroids*.

What are the preparations of ergot and their doses?

EXTRACTUM ERGOTÆ (*extract of ergot*) is five times as strong as the fluid extract; dose gr. v-xv (0.3-1.0). It may be used hypodermically, dissolved in water and filtered.

EXTRACTUM ERGOTÆ FLUIDUM (*fluid extract of ergot*), dose m_v -f3iv (0.3-15.0).

VINUM ERGOTÆ (*wine of ergot*) contains 15 per cent. of powdered ergot; dose f3j-iv (4.0-15.0).

HYDRASTIS.

What is hydrastis?

Hydrastis is the *rhizome and roots* of *H. canadensis*, yellow root or golden seal (*Nat. Ord.* Ranunculaceæ). It has been already considered under the Bitter Tonics (*q. v.*).

HYDRASTINÆ HYDROCHLORAS—HYDRASTINE HYDROCHLORATE.

What is this substance?

It is the hydrochlorate of an artificial alkaloid produced by the oxidation of hydrastine, and has been found to produce powerful uterine contractions, stimulating also the muscular fibres of the heart, arterioles, and intestines. It probably stimulates the voluntary muscles, but in a lesser degree.

In fatal doses it depresses powerfully the entire motor nervous tract and causes death by respiratory failure.

It has been used with success in menorrhagia and metrorrhagia and other uterine hemorrhages.

It is also recommended as a heart tonic. Dose gr. $\frac{1}{4}$ -j (0.016-0.065).

GOSSYPHII RADICIS CORTEX—BARK OF COTTON ROOT.

What is gossypii radicis cortex?

It is the *bark of the root* of *Gossypium herbaceum* and other species of *Gossypium* (*Nat. Ord.* Malvaceæ).

Given to animals, it first slows and then quickens the cardiac action, which is always enfeebled by first stimulating and then paralyzing the cardio-inhibitory apparatus and depressing the cardio-motor ganglia. It stimulates and then depresses the vaso-motor centres, causing a rise, but subsequently a fall, in the blood pressure. It stimulates non-striated muscular fibres, and appears to excite rhythmical contractions of the gravid uterus and to increase their power when present. It depresses the reflex centres of the cord, and causes

death by respiratory paralysis, generally preceded by convulsions, which are not of cerebral origin.

It has been *recommended* as a substitute for ergot in *uterine inertia*, *post-partum hemorrhage*, and to aid the expulsion of *uterine myomata*. It has been used in *chronic metritis*, and although not as efficient as ergot, is useful in *menorrhagia*.

EXTRACTUM GOSSYPH RADICIS FLUIDUM (*fluid extract of cotton root*), dose f3ss-j (2.0–4.0).

QUININA—QUININE.

Is quinine oxytocic?

Although quinine is used for this purpose, it probably possesses no action on the uterus *per se*, but as an antiperiodic or as a tonic, is capable of overcoming the morbid conditions (as malaria) which, by depressing the system at large, may prevent normal, healthy uterine contractions.

ORDER IV.—UTERINE SEDATIVES.

What are uterine sedatives?

They are medicines which are employed to depress the uterine muscular or nervous apparatus and restrain violent, irregular or untimely uterine contractions.

What medicines are used for this purpose, and when are they employed?

They are principally useful during pregnancy, to prevent abortion which is threatening, but not inevitable. They are also of service in arresting the irregular uterine contractions which sometimes occur prior to labor, and are called “false labor pains.” For the former purpose opium, cannabis indica and the motor depressants—as chloral, the bromides, tartar emetic and viburnum prunifolium, especially the first- and last-named remedies—are used, while for the latter opium and chloral are employed. To moderate too violent contractions during labor, ether is the best remedy. Of these medicines all have been described except

VIBURNUM PRUNIFOLIUM.

What is viburnum prunifolium?

It is the *bark* of *V. prunifolium*, sloe or black haw (*Nat. Ord. Caprifoliaceæ*), a small indigenous tree. It contains *valerianic acid*, a bitter principle (*viburnin*), resin, tannin, etc. Its *physiological effects* are not understood. It appears to diminish reflex irritability and to have a sedative effect on involuntary muscular fibre and possibly lowers blood-pressure. Toxic effects are only seen after enormous doses have been administered, and consist of headache, dryness of the mouth and throat, and disordered vision.

It has been employed as a *uterine sedative* in *habitual* and in *threatened abortion*, in congestive and neuralgic *dysmenorrhœa*, in *menorrhagias* and in *nervous disturbances* occurring at the *climacteric* period, in *post-partum dolores* when no clots or fragments of the placenta remain in the uterus, and in *false labor-pains*. It has also been used in *colicky diarrhœa* and *dysentery*, and is even recommended for *cramps* occurring in the *voluntary muscles*.

EXTRACTUM VIBURNI FLUIDUM (*fluid extract of viburnum*), dose fʒss-ij (2.0-8.0) in elixir of orange.

ORDER V.—EMMENAGOGUES.

What are emmenagogues?

Emmenagogues are medicines employed to increase or reëstablish the menstrual flow when it is scanty or suppressed from other causes than pregnancy or the climacteric period.

When the catamenia are arrested from pregnancy, medicines used to bring them on cannot but be injurious to the woman, as the doses necessary are sufficient to cause poisoning, one of the symptoms of which may (or may not) be uterine hemorrhage. This is especially true of cantharides and of the volatile oils.

What medicines are used as emmenagogues?

When amenorrhœa depends upon anæmia or chlorosis the best emmenagogues are iron (combined with aloes or myrrh), arsenic or manganese. When local anæmia or an atonic condition of the uterus is present, aloes (*q. v.*) is very useful. If, on the other hand, pelvic plethora exists with constipation and a scanty flow often of a

thick black discharge accompanied with pelvic pain and fullness, the saline cathartics are employed.

In suppression of the menses from cold, aloes and potassium permanganate are usually more successful. In amenorrhœa due to an atonic condition of the uterus, the more stimulating emmenagogues are given, generally with aloetic purgatives.

STIMULATING EMMENAGOGUES.

Name the stimulating emmenagogues.

Savine, tansy, apiol, cantharides and guaiac.

SABINA—SAVINE.

What is savine ?

Savine is the *tops* of *Juniperus Sabina* (*Nat. Ord. Coniferæ*), a small evergreen shrub of Southern Europe. It contains a *volatile oil*. When applied *locally* savine is irritant. In *medicinal doses* it increases the circulation, is a gastric, uterine and renal irritant, and proves diuretic. In overdoses it is a gastro-intestinal irritant with great depression, unconsciousness, convulsions, hemorrhages from the uterus and kidneys, suppression of urine and, in pregnant females, abortion. Death occurs from collapse. Savine is *used* in relaxed states of the uterus and uterine vessels, and has proved successful in the treatment both of *amenorrhœa* and *menorrhagia* when due to atonic condition of this organ.

EXTRACTUM SABINÆ FLUIDUM (*fluid extract of savine*), dose $\text{m}_{\text{V-X}}$ (0.3-0.6).

OLEUM SABINÆ (*oil of savine*), the best preparation for internal use; dose gtt. v-X (0.3-0.6), in capsules or emulsion, every 3 or 4 hours.

TANACETUM—TANSY.

What is tansy ?

Tansy is the *leaves* and *tops* of *Tanacetum vulgare* (*Nat. Ord. Compositæ*). It contains a volatile oil and a bitter principle (*tanacetin*). It is a gastro-intestinal irritant and diuretic, and in large doses causes death from respiratory paralysis, preceded by epileptiform convulsions of cerebral origin, and unconsciousness. It is used as a stimulating emmenagogue, but is a very dangerous remedy. The minimum fatal quantity is unknown. Dose gr. x-xxv (0.6-1.5), in infusion; of the oil, m_{j} (0.06) may be given.

APIOL.

What is apiol?

Apiol (not official), or parsley-camphor, is obtained from the volatile oil of the root of *Petroselinum sativum* (*Nat. Ord. Umbelliferæ*). It is very efficacious as a stimulating emmenagogue in *functional* and *anæmic amenorrhœa* and in *neuralgic dysmenorrhœa*, but is contraindicated in any condition of plethora. Dose gtt. iij-x (0.18–0.6) in gelatin capsules given morning and evening for 4 or 5 days before the expected menstrual period.

The French capsules or pearls usually contain gr. iij $\frac{9}{16}$ (0.25) of apiol.

CANTHARIDES—GUAIAIC.

What are these substances?

Cantharides will be considered under “Irritants.” The tincture is used as a stimulating emmenagogue, dose m*ij*-v (0.18–0.3), t. i. d., great care being taken to avoid strangury.

Guaiaic has been considered with the “Alteratives.”

(D) ON THE ALIMENTARY CANAL.

ORDER I.—EMETICS.

What are emetics, and how do they act?

Emetics are medicines which are given to induce vomiting. They may act either by directly irritating the stomach, thus causing expulsion of its contents in a reflex manner, when they are called *local emetics*, or by acting, after absorption, on the reflex centre of the medulla which governs emesis, when they are called *systemic emetics*. It does not always follow that a remedy is a systemic emetic because it causes vomiting when given hypodermically or per rectum, for it may be eliminated by, and produce irritation of, the gastric mucous membrane, thus exciting reflex vomiting.

For what purposes are emetics used?

Emetics are used (1) to unload the stomach; (2) to depress the circulatory system by inducing nausea (as in the early stages of bronchitis or of tonsillitis, in which emetics, frequently repeated in

small doses, will sometimes arrest the inflammation); (3) to relax the muscular system and relieve spasm (as in spasmodic croup); (4) to promote secretion and excretion; (5) to expel foreign bodies or secretions from the air passages, and (6) as revulsives.

What are the contraindications to their employment?

They should never be employed in cases of cerebral or gastric inflammation, and should be used cautiously, if at all, in the latter stages of pregnancy, in hernia, or where great depression exists, as in the latter stages of capillary bronchitis or of croup.

Name the medicines employed as emetics.

The vegetable emetics are ipecac, apomorphine and mustard. *The mineral emetics* are zinc sulphate, copper sulphate, alum, turpeth mineral and tartar emetic.

IPECACUANHA—IPECAC.

What is ipecac, and what chemical principles does it contain?

Ipecac is the *root* of *Cephaëlis Ipecacuanha* (*Nat. Ord. Rubiaceæ*), a plant of Brazil. Its effects depend upon an alkaloid, *emetine*, which is combined with *ipecacuanhic acid*.

What are its physiological actions?

Locally applied, ipecac is an irritant, either to the skin or mucous membranes. Given *internally* in *small doses*, repeated, it increases the perspiration, saliva, bile, the secretions from the broncho-pulmonary and gastro-intestinal mucous membranes, and causes nausea. In sufficient amounts it greatly augments the secretions and induces vomiting, without much nausea or depression, no matter how exhibited, emesis being more rapid when it is taken by the mouth.

Toxic doses in animals depress the *respiration* and paralyze the *heart* directly. Post-mortem examination shows the mucous membranes of the stomach, intestines and lungs to be intensely congested. Sometimes, however, the lungs are almost exsanguine.

What are the medicinal uses of ipecac?

Ipecac is employed *as an emetic*, and is the safest remedy which can be given for this purpose, especially to children. It is particu-

larly of service when it is desired to evacuate the contents of an overloaded stomach without producing much depression, as an adjunct to other remedies in *croup* and in the earlier stages of *capillary bronchitis*. In narcotic poisoning it is not as efficient as the mineral emetics.

As an *expectorant* it is added to cough medicines in the dry stage of bronchitis. As a *cholagogue* it is a valuable addition to purgative pills. As a *hæmostatic* it is highly recommended in *hæmoptysis* and *post-partum hemorrhage*.

In the treatment of *dysentery*, especially when epidemic, it is of great service, given in large dose (gr. xx-3ij (1.28-8.0), every few hours). When thus administered a tolerance is soon established, and no nausea or vomiting is produced. It is also useful in various kinds of *diarrhœa*, especially of choleric and tuberculous diarrhœa. *Reflex vomiting*, as the vomiting of pregnancy, has sometimes been arrested by minute doses of the wine of ipecac.

What are the preparations of ipecac and their doses?

Ipecac may be given in powder, dose as an *emetic* gr. x-xx (0.64-1.28); as a *nauseant*, gr. ss-ij (0.03-0.13); as an *expectorant* and *diaphoretic*, gr. $\frac{1}{2}$ - $\frac{1}{4}$ (0.016-0.032); as a *stomachic tonic* and *hepatic stimulant*, gr. $\frac{1}{12}$ - $\frac{1}{8}$ (0.005-0.008).

EXTRACTUM IPECACUANHÆ FLUIDUM (*fluid extract of ipecac*), dose as an emetic m℥xx-xxx (1.23-1.8), repeated until vomiting occurs.

VINUM IPECACUANHÆ (*wine of ipecac*) contains 10 per cent. of the fluid extract; dose as emetic f3j-iv (4.0-16.0).

SYRUPUS IPECACUANHÆ (*syrup of ipecac*) contains 7 per cent. of the fluid extract; as an emetic f3j-ij (4.0-8.0) may be given to a child, or m℥v-x (0.3-0.6) as an expectorant.

TROCHISCI IPECACUANHÆ (*troches of ipecac*), each contains about gr. $\frac{1}{3}$ (0.02) of ipecac.

Powder of ipecac and opium (Dover's powder), *tincture of ipecac and opium*, and *troches of ipecac and morphine* have been considered under "Opium."

APOMORPHINÆ HYDROCHLORAS—APOMORPHINE HYDROCHLORATE.

What is apomorphine hydrochlorate?

It is the *hydrochlorate* of an *artificial alkaloid* obtained from morphine or codeine.

What are its physiological actions?

Given to warm-blooded animals in *small doses* it causes vomiting; after *large doses* there is salivation, violent vomiting and muscular tremors; *toxic doses* induce great restlessness, paresis, convulsions, hurried respiration and death from asphyxia, without the occurrence of emesis.

It is a systemic emetic, probably in medicinal doses stimulating the reflex centre for *vomiting*, but in large doses depressing and paralyzing it. No matter how administered, emesis takes place promptly, with but little nausea and rarely accompanied by much depression. Large doses appear to stimulate and finally paralyze the respiratory centre. It increases the *pulse-rate* by stimulating the accelerator fibres (?), and in large doses by stimulating the vasomotor centres, it raises the arterial pressure. In toxic doses it depresses the cardiac muscle and thus reduces the blood pressure. In large doses it is at first a *cerebral* stimulant, but later depresses and paralyzes the brain. The cause of the convulsion is not understood. It is a *muscle* poison.

What are the uses and doses of apomorphine hydrochlorate?

It may be employed *as an emetic* whenever prompt evacuation of the stomach is required, and may be administered either hypodermically or by the mouth. In *narcotic poisoning* it is highly recommended.

As an expectorant in the early stages of *acute bronchitis* or in *chronic bronchitis* without much expectoration, it is very serviceable, and can be advantageously used in *capillary bronchitis* of children both as an expectorant and emetic. The adult dose as an *emetic* is gr. $\frac{1}{16}$ (0.004) hypodermically, or gr. $\frac{1}{16}$ – $\frac{1}{8}$ (0.005–0.008) by the mouth. As *an expectorant* smaller doses are required.

SINAPIS—MUSTARD.

What is mustard, and how is it used as an emetic?

Mustard is the *seed* of *Brassica nigra* and *B. alba* (*Nat. Ord.* Cruciferae). Mustard flour is irritant to the mucous membranes as well as to the skin, and is a prompt and efficient local emetic, especially useful when torpor of the stomach is present. It will be more fully considered hereafter. Dose as an emetic from a teaspoonful to

a tablespoonful (4.0–16.0) of the flour, in a glass of tepid water, repeated, if necessary, in 15 or 20 minutes.

What mineral emetics are employed?

ZINCI SULPHAS (*zinc sulphate*) is a prompt and efficient local emetic. Dose gr. x–xx (0.6–1.2), or even gr. xxx (1.9) in narcotic poisoning.

CUPRI SULPHAS (*copper sulphate*) is a prompt local emetic, but more irritating than the zinc salt. Dose gr. iij–v (0.10–0.3) or even gr. x (0.6) in narcotic poisoning. It is the best emetic in poisoning by phosphorus (*q. v.*).

ALUMEN (*alum*) is also a local emetic of value, especially useful in *membranous croup*, and useful also in *capillary bronchitis*; dose a teaspoonful (4.0) of the powdered drug in molasses, or, better, in syrup of ipecac.

HYDRARGYRI SUBSULPHAS FLAVUS (*yellow mercurial subsulphate—Turpeth mineral*), used as an emetic in *croup*; dose gr. j–v (0.06–0.3) repeated every 10 or 15 minutes until free vomiting occurs.

ANTIMONII ET POTASSII TARTRAS (*antimonium and potassium tartrate—tartar emetic*) is rarely used as an emetic because of the great depression which it induces; dose gr. j–ij (0.06–0.13).

Of these substances the three first will be found under the head of astringents, while the two last have already been considered.

ORDER II.—GASTRIC SEDATIVES.

What are gastric sedatives?

They are medicines employed to diminish gastric irritability, and thus allay nausea and vomiting.

What medicines are used for this purpose?

When vomiting is due to *acute irritation*, as from an overloaded stomach, the ingestion of acrid substances, or cholera morbus, it is best treated, after the contents have been evacuated, by minute doses of calomel (gr. $\frac{1}{16}$ (0.005)) with sodium bicarbonate (gr. j (0.06)); pellets of ice dissolved in the mouth; lime-water and rest. A small dose of brandy will sometimes “settle the stomach.” If *irritability persists*, small doses of morphine, or of diluted hydrocyanic acid or

creasote, or bismuth and sodium bicarbonate answer better. When due to the formation of gases from *fermentation*, creasote, carbolie acid, thymol, sulphurous acid, the sulphites or charcoal are given.

When vomiting occurs on taking food after *long abstinence*, and perhaps *preceding irritation*, calomel and sodium bicarbonate in minute amounts, followed by small doses of brandy, and milk and lime-water by the teaspoonful (4.0) at intervals, should be employed. *Ether-vomiting* is best met with teaspoonful (4.0) doses of very hot water given every 10 or 15 minutes. *Reflex vomiting*, as of seasickness or of pregnancy, usually requires a medicine which will depress the centre in the medulla presiding over the act, as opium, the bromides, or diluted hydrocyanic acid. Among other remedies used for the latter variety of vomiting may be mentioned, tincture of nux vomica (gtt. $\frac{1}{4}$ -j (0.015-0.06), cerium oxalate (gr. $\frac{1}{4}$ -v (0.016-0.3)), wine of ipecac (gtt. $\frac{1}{4}$ -ij (0.015-0.12)), and Fowler's solution (m $\frac{1}{4}$ -j (0.015-0.06)). Counter-irritation over the epigastrium often aids the action of gastric sedatives. The above remedies have already been noticed or will be discussed in their appropriate places.

ORDER III.—CARMINATIVES.

What are carminatives ?

Carminatives are medicines employed to aid the expulsion of gases from the stomach and intestines.

What medicines are used as carminatives ?

The *aromatics*, which owe their virtues principally to the volatile oil which they contain, are especially used for this purpose. Other drugs containing volatile oils, as asafetida, eucalyptus, etc., are also carminative ; and ether is sometimes employed for the same object.

What are the general effects of the aromatics ?

Locally they are irritant and rubefacient. The oils are also local anæsthetics. *Internally* in *medicinal doses* they are gastric and intestinal stimulants, increasing peristalsis and temporarily increasing to a slight extent the force and frequency of the pulse. In *larger doses* they stimulate the brain and cause a species of intoxication. In *very large doses* they are irritant and narcotic poisons.

What are their medicinal uses?

The aromatics are used internally as *carminatives*; to prevent the *gripping* of purgatives; to *correct* the unpleasant *taste* or *smell* of other medicines, and as *gastric stimulants*.

Locally they are employed as *rubefacients*, *antiseptics* and *anodynes*. They should *never be used* when inflammation of the stomach or bowels is present, but in diarrhoea due to relaxation they are often of service.

Name the aromatics, their chemical constituents, preparations and doses.

CAPSICUM, or red pepper, is the *fruit* of *C. fastigiatum* (*Nat. Ord. Solanaceæ*). It contains a *resin*, a *fixed* and a *volatile oil*. It is *used* as a *condiment*, as a *gastric stimulant*, particularly in the case of drunkards, as an addendum to tonic pills, as a *carminative* in cases of feeble digestion, as a gargle in tonsillitis, and externally as a *rubefacient*.

EXTRACTUM CAPSICI FLUIDUM (*fluid extract of capsicum*), dose $\text{m}_{\text{v-x}}$ (0.3-0.6).

TINCTURA CAPSICI (*tincture of capsicum*), dose $\text{m}_{\text{v-f3j}}$ (0.3-4.0).

OLEORESINA CAPSICI (*oleoresin of capsicum*), dose gr. ss-j (0.03-0.06).

EMPLASTRUM CAPSICI (*plaster of capsicum*), for external use.

PIPER (*pepper*) is the *unripe fruit* of *Piper nigrum* (*Nat. Ord. Piperaceæ*). It contains a *volatile oil*, an *acid resin* and a neutral principle, *piperin*, which increases the respiration and circulation, dilates the pupil, and finally paralyzes the heart in systole. It has been used as an *antiperiodic*. Pepper is used as a *condiment* and sometimes as a *stomachic stimulant*, combined with other medicines, in atonic dyspepsia, and as a *carminative*.

OLEORESINA PIPERIS (*oleoresin of pepper*), dose gr. ss-ij (0.03-0.13).

PIPERINUM (*piperin*), dose gr. j-v (0.06-0.3).

CINNAMOMUM CASSIA (*cassia cinnamon*), the *bark of the shoots* of one or more undetermined species of *cinnamomum*, grown in China (*Nat. Ord. Laurinææ*); CINNAMOMUM SAIGONICUM (*Saigon cinnamon*), the *bark* of an undetermined species of *cinnamomum*

(*Nat. Ord. Laurineæ*) (which comes from Cochin-China); and CINNAMOMUM ZEYLANICUM (*Ceylon cinnamon*), the inner bark of the shoots of *cinnamomum Zeylanicum* (*Nat. Ord. Laurineæ*), are all recognized by the Pharmacopœia. Of these varieties the latter is considered the finest and the former the poorest in quality. They contain *volatile oil*, *tannic acid*, etc., are slightly astringent, and are used as *carminatives*, *flavoring ingredients*, and *vehicles*.

TINCTURA CINNAMOMI (*tincture of cinnamon*) contains 10 per cent. of powdered Ceylon cinnamon; dose fʒj-ij (4.0-8.0).

AQUA CINNAMOMI (*cinnamon water*) contains 0.2 per cent. of the oil of cinnamon, and is used as a vehicle.

OLEUM CINNAMOMI (*oil of cinnamon*) is a volatile oil distilled from cassia cinnamon; dose gtt. j-ij (0.06-0.18).

SPIRITUS CINNAMOMI (*spirit of cinnamon*) contains 10 per cent. of the oil; dose ℥x-xxx (0.6-2.0).

MYRISTICA (*nutmeg*); MACIS (*mace*). These are parts of the fruit of *Myristica fragrans* (*Nat. Ord. Myristicaceæ*), the *arillode of the seed* being mace, and the *seed* nutmegs. Both contain volatile oils. They are aromatic stimulants, carminatives, and in large doses narcotics.

OLEUM MYRISTICÆ (*oil of nutmeg*), dose gtt. j-v (0.06-0.3).

SPIRITUS MYRISTICÆ (*spirit of nutmeg*) contains 5 per cent. of the oil; dose fʒss-ij (1.8-7.3).

CARYOPHYLLUS (*cloves*) is the *unexpanded flowers* of *Eugenia aromatica* (*Nat. Ord. Myrtaceæ*). They contain *volatile oil*, *tannic acid*, *caryophyllin* and *eugenin*. They are gastric stimulants and carminatives, but are principally employed as a condiment. The oil is used as a local anodyne to carious teeth.

OLEUM CARYOPHYLLI (*oil of cloves*), dose gtt. iij-v (0.18-0.3).

PIMENTA, or allspice, is the *nearly ripe fruit* of *Pimenta officinalis* (*Nat. Ord. Myrtaceæ*). It contains a *volatile oil*, and is carminative.

OLEUM PIMENTÆ (*oil of pimenta*), dose gtt. iij-v (0.18-0.3).

ZINGIBER (*ginger*) is the *rhizome* of *Z. officinale* (*Nat. Ord. Scitamineæ*). It contains a *volatile oil* and an *acrid resin*. It is a

gastric stimulant and *carminative*, much employed in flatulency and colic and as an addition to other stomachics.

EXTRACTUM ZINGIBERIS FLUIDUM (*fluid extract of ginger*), dose $\text{m}_{\text{x-xxx}}$ (0.6-2.0).

TINCTURA ZINGIBERIS (*tincture of ginger*), dose $\text{f}\text{3ss-j}$ (2.0-4.0).

SYRUPUS ZINGIBERIS (*syrup of ginger*), used as a vehicle for other medicines.

OLEORESINA ZINGIBERIS (*oleoresin of ginger*), used as a corrective to purgative pills to prevent griping; dose $\text{m}_{\text{ss-ij}}$ (0.03-0.12).

TROCHISCI ZINGIBERIS (*troches of ginger*), dose 2 or 3 lozenges.

CARDAMOMUM (*cardamom*) is the *fruit* of *Elettaria repens* (*Nat. Ord. Scitamineæ*), and contains a *volatile oil*. It is used as a gastric stimulant and carminative, and as an adjuvant and corrective of other medicines.

TINCTURA CARDAMOMI (*tincture of cardamom*), dose $\text{f}\text{3j-ij}$ (4.0-8.0).

TINCTURA CARDAMOMI COMPOSITA (*compound tincture of cardamom*) contains also caraway, cassia, cinnamon, and cochineal; dose $\text{f}\text{3j-ij}$ (4.0-8.0).

PULVIS AROMATICUS (*aromatic powder*) consists of ceylon cinnamon and ginger (35 per cent. each), with cardamom and nutmeg (15 per cent. each). An excellent carminative preparation. Dose gr. x-xxx (0.6-2.0).

EXTRACTUM AROMATICUM FLUIDUM (*aromatic fluid extract*) is a fluid extract of aromatic powder. Dose $\text{m}_{\text{xv-xxx}}$ (1.0-2.0).

OLEUM CAJUPUTI (*oil of cajuput*) is the *volatile oil* distilled from the *leaves* of *Melaleuca Leucadendron* (*Nat. Ord. Myrtaceæ*), and is very destructive to low organisms. It has been employed as a remedy against ascarides and, locally, in parasitic skin diseases, but is chiefly used as a gastric stimulant and carminative in flatulent colic and to prevent the griping of cathartics. Dose $\text{m}_{\text{j-v}}$ (0.06-0.3).

CALAMUS is the *rhizome* of *Acorus Calamus* (*Nat. Ord. Aroideæ*). It contains a *volatile oil* and a glucoside, *acorin*. It is an aromatic stimulant and is somewhat tonic.

EXTRACTUM CALAMI FLUIDUM (*fluid extract of calamus*), dose $\text{m}_{\text{x-xxx}}$ (0.6-2.0).

OLEUM GAULTHERIÆ (*oil of gaultheria*) is a *volatile oil* distilled from the *leaves* of *G. procumbens* or wintergreen (*Nat. Ord. Ericaceæ*), and has been considered under the head of "Salicylic Acid". It consists almost entirely of methyl salicylate and is nearly identical with volatile oil of *Betula* (*vid. Salicylates*); dose in rheumatism $\text{m}_{\text{V-XV}}$ (0.31–0.93 c. c.).

SPIRITUS GAULTHERIÆ (*spirit of gaultheria*) contains 5 per cent. of the oil; dose $\text{m}_{\text{X-XXX}}$ (0.62–1.85 c. c.).

The following herbs, belonging to the *Nat. Ord. Labiataë*, contain volatile oils, and are used as carminatives, gastric stimulants and flavoring ingredients:—

OLEUM LAVENDULÆ FLORUM (*oil of lavender flowers*) is a *volatile oil* distilled from the *flowers* of *L. vera*. Dose gtt. j–v (0.06–0.3). *Spiritus Lavendulæ* (*spirit of lavender*) and *Tinctura Lavendulæ Composita* (*compound tincture of lavender*) are also official; dose of either $\text{f}\text{3}_{\text{ss-j}}$ (2.0–4.0).

MENTHA PIPERITA (*peppermint*) and MENTHA VIRIDIS (*spear-mint*). The *leaves* and *tops* are official. The oils (*oleum menthæ piperitæ* and *oleum menthæ viridis*, dose gtt. j–v (0.06–0.3)) are usually given in the form of *spirit* (*spiritus menthæ piperitæ* and *spiritus menthæ viridis*), dose $\text{m}_{\text{V-XXX}}$ (0.3–2.0). The *waters* (*aqua menthæ piperitæ* and *aqua menthæ viridis*) are used as vehicles. *Troches of peppermint* (*trochisci menthæ piperitæ*) are sometimes used in flatulent colic.

OLEUM ROSMARINI (*oil of rosemary*) is a *volatile oil* distilled from the *leaves* of *Rosmarinus officinalis*. It is carminative, but is chiefly used as an ingredient of rubefacient liniments.

HEDEOMA (*pennyroyal*). The *leaves* and *tops* of *H. pulegioides* are official. The *volatile oil* (*oleum hedeomæ*) may be given in doses of $\text{m}_{\text{j-v}}$ (0.06–0.3).

MARRUBIUM (*horehound*). The *leaves* and *tops* of *M. vulgare* are used as a gastric stimulant and expectorant.

SALVIA (*sage*). The *leaves* of *S. officinalis* are slightly astringent as well as aromatic. An infusion is sometimes used as a gargle in sore throat.

OLEUM THYMI (*oil of thyme*), a *volatile oil* distilled from the *leaves* and *flowering tops* of *Thymus vulgaris*, contains much *thymol*

(*q. v.*), and is therefore not only aromatic and carminative, but antiseptic and antifermentative.

The following *seed* are derived from plants belonging to the *Nat. Ord. Umbelliferæ*:—

FÆNICULUM (*fennel*), from *F. capillaceum*. Of the oil (*oleum fœniculi*), the dose is gtt. v-x (0.3-0.6). A water (*aqua fœniculi*) is also official.

CARUM (*caraway*), from *C. Carvi*. Dose of the oil (*oleum cari*) gtt. j-x (0.06-0.6).

ANISUM (*anise*), from *Pimpinella anisum*. Dose of the oil (*oleum anisi*) gtt. v-xv (0.3-0.9). A spirit (*spiritus anisi*) and a water (*aqua anisi*) are also official.

CORIANDRUM (*coriander*), from *C. sativum*. The oil (*oleum coriandri*) is official.

ILLICIIUM (*star-anise*) is the *fruit* of *I. verum* (*Nat. Ord. Magnoliaceæ*). It contains a volatile oil, and is used as a substitute for anise.

The following preparations are only used as flavors and vehicles:—

AURANTII AMARI CORTEX (*bitter orange peel*); AURANTII DULCIS CORTEX (*sweet orange peel*). These are respectively the *rind* of the *fruits* of *Citrus vulgaris* (bitter orange) and of *C. Aurantium* (sweet orange) (*Nat. Ord. Rutaceæ*). The following preparations are official:—

Oil of orange flowers, or oil of neroli (*oleum aurantii florûm*), a volatile oil distilled from the fresh flowers of the bitter orange; *stronger orange flower water* (*aqua aurantii florûm fortior*—*aqua aurantii florûm*, Pharm. 1880—*triple orange flower water*), water saturated with the volatile oil of fresh orange flowers, obtained as a by-product in the distillation of the oil of orange flowers; *orange flower water* (*aqua aurantii florûm*), equal volumes of stronger orange flower water and distilled water freshly mixed; *syrup of orange flowers* (*syrupus aurantii florûm*); *syrup of orange peel* (*syrupus aurantii*); *oil of orange peel* (*oleum aurantii corticis*); *spirit of orange* (*spiritus aurantii*); *compound spirit of orange* (*spiritus aurantii compositus*) contains oil of orange peel (200), oil

of lemon (50), oil of coriander (20), and oil of anise (5) in 1000 parts of the preparation. It is used in making *elixir aromaticum* (*aromatic elixir*), which contains 12 parts of the compound spirit in 1000 parts of the preparation; *fluid extract of bitter orange peel* (*extractum aurantii amari fluidum*); *tinctures of bitter and of sweet orange peel* (*tinctura aurantii amari*, *tinctura aurantii dulcis*)—dose of either tincture, fʒj-ij (4.0–8.0).

VANILLA, the *fruit* of *V. planifolia* (*Nat. Ord. Orchidacææ*), contains *vanillin*. The *tincture* (*tinctura vanillæ*) is official.

The *oil of turpentine* is an excellent carminative and will be considered hereafter.

ORDER IV.—CATHARTICS.

What are cathartics?

Cathartics are medicines employed to produce an evacuation from the bowels either by increasing intestinal peristalsis, or by increasing the secretions of the canal and its appendages.

For what purposes are they used?

They are used (1) to unload the bowels; (2) to deplete the blood vessels by abstracting water from the blood; (3) to promote absorption; (4) to stimulate secretion; (5) to eliminate noxious material from the blood (as the poison of uræmia); (6) to act as revulsives, and (7) to influence the pelvic circulation.

How may cathartics be classified?

Cathartics are divided into laxatives, simple purgatives, salines and drastics.

LAXATIVES.

What are laxatives?

Laxatives are such substances as unload the bowels, acting without irritation and not causing purgation.

Name the laxatives.

Tamarind, manna, cassia fistula, frangula, cascara sagrada, castor oil, sulphur and sulphurated potassa.

TAMARINDUS—TAMARIND. MANNA. VIOLA TRICOLOR.

What are these substances?

TAMARINDUS (*tamarind*) is the *preserved pulp of the fruit* of *T. indica* (*Nat. Ord. Leguminosæ*), a tree of India. Eaten as a preserve (℥ss-j (15.0-31.0)) it is laxative. An infusion makes a pleasant refrigerant drink. It enters into *confection of senna*.

MANNA is the *concrete saccharine exudation* of *Fraxinus Ornus* (*Nat. Ord. Oleaceæ*). It contains *mannit*, *fraxin*, and *resin* which is probably the laxative principle. Manna is usually combined with other cathartics. Dose for an adult ℥j-ij (31.0-62.0); for children, ℥j-iv (4.0-15.0).

CASSIA FISTULA. FRANGULA. RHAMNUS PURSHIANA.

What are these substances?

CASSIA FISTULA is the *fruit* of *C. Fistula* or purging cassia (*Nat. Ord. Leguminosæ*). It is an ingredient of the *confection of senna*. Dose ℥j-℥j (4.0-31.0).

FRANGULA is the *bark* of *Rhamnus Frangula* or alder buckthorn (*Nat. Ord. Rhamnaceæ*). It contains *frangulin* and *emodin*, and as it is an active emetic and hydragogue when fresh, it should be kept a year at least before being used; dose of the *fluid extract* (*extractum frangulæ fluidum*) f℥ss-j (2.0-4.0).

RHAMNUS PURSHIANA (*cascara sagrada* or *chittem bark*) is the *bark* of *R. Purshiana* (*Nat. Ord. Rhamnaceæ*), a tree of California. It contains several resins and is an efficient laxative, very useful in habitual constipation; dose of the *fluid extract* (*extractum rhamni purshianæ fluidum*) m℥x-f℥j (0.6-4.0).

OLEUM RICINI—CASTOR OIL.

What is castor oil?

It is a *fixed oil* expressed from the *seed* of *Ricinus communis*, or *Palmi Christi* (*Nat. Ord. Euphorbiaceæ*).

What are its effects and uses?

In *small doses* it is a mild laxative, in *larger amounts* a simple

purgative. Applied with friction to the abdomen of children it frequently purges. It is very useful whenever it is desired to evacuate the bowels and remove any irritant materials, as undigested food. For this purpose it is often used in acute dysenteries and diarrhœas, especially of children; dose $\text{f}\text{3j}-\text{iv}$ (4.0–16.0), best given in whiskey and water, in such a manner that the entire inside of the glass will be wetted to its margins with the vehicle and the oil form a globule surrounded by the fluid; or it may be given beaten up with double the quantity of sherry; or mixed with equal parts of glycerin and flavored with some aromatic oil.

SULPHUR.

What is sulphur?

Sulphur is a chemical element obtained by purifying native sulphur or by decomposing native sulphides.

What are its effects, uses and preparations?

It acts as a mild laxative and is eliminated in all the excretions. Its continued use causes flatulence from the formation of sulphuretted hydrogen. It is used as a *laxative*, especially in cases of *hemorrhoids*, and as an *alterative* both internally and externally in *chronic rheumatism*, *muscular rheumatism*, *sciatica*, and in various forms of *skin disease*, especially those of parasitic origin. As sulphur is excreted largely by the hair, it has been used locally and internally in the treatment of *alopecia*. It is also used as an *expectorant*. It is official as SULPHUR SUBLIMATUM (*sublimed sulphur*, sometimes called “flowers of sulphur”), SULPHUR LOTUM (*washed sulphur*), and SULPHUR PRÆCIPITATUM (*precipitated sulphur*); dose of either preparation, $\text{3j}-\text{iv}$ (4.0–16.0) in syrup. UNGUENTUM SULPHURIS (*sulphur ointment*, containing 30 per cent. of washed sulphur) is used externally.

POTASSA SULPHURATA—SULPHURATED POTASSA.

What is sulphurated potassa?

Sulphurated potassa, or liver of sulphur, is an uncertain compound containing about 50 per cent. of potassium sulphide.

In *small doses* it is decomposed in the stomach and acts like sulphur. In *large doses* it is a gastro-intestinal irritant and has

produced death. It may be *used* to fulfil the indications of sulphur, but is chiefly employed as an *alterative* where a tendency to the formation of abscesses exists, in chronic rheumatism and in scaly skin affections. It is used *externally* in the form of bath or ointment; dose gr. ij-x (0.1-0.6).

SIMPLE PURGATIVES.

What are the simple purgatives?

They are medicines employed to produce active purgation, but which do not produce sufficient irritation to cause inflammation or depression, even when large doses are given.

Name the agents of this group.

Castor oil, rhubarb, juglans, aloes, senna, leptandra, euonymus, and mercurials.

Castor oil has been described with the *laxatives*.

RHEUM—RHUBARB. JUGLANS.

What are these medicines?

RHUBARB is the *root* of *Rheum officinale* (*Nat. Ord.* Polygonaceæ). It contains *chrysophan*, *chrysophanic acid*, 4 resins (*erythroretin*, *emodin*, *phæroetin* and *aporetin*), *rheotannic* and *rheumic acids*. It is a tonic, mild purgative, and an astringent owing to the rheotannic acid, and hence after purging it has a tendency to cause constipation. It tinges the milk and urine yellow, and stimulates the flow of bile. It is much used in *diarrhœa* from relaxation, in the early stages of *summer complaint* of children and in *dyspepsia* with constipation. The official preparations and their doses are: the *extract* (*extractum rhei*), gr. v-xx (0.3-1.3); *pills* (*pilulæ rhei*) each contain gr. iij (0.2) of rhubarb; *compound pills* (*pilulæ rhei compositæ*) each contain gr. ij (0.13) of rhubarb, gr. jss (0.1) of aloes, with myrrh and oil of peppermint; dose No. j-iv; *compound powder* (*pulvis rhei compositus*) contains rhubarb and magnesia, dose 3ss-j (2.0-4.0); *fluid extract* (*extractum rhei fluidum*), ℥xx-xxx (1.2-2.0); *mixture of rhubarb and soda* (*mistura rhei et sodæ*), for a child f3ss-j (2.0-4.0), for an adult f3j-iv (4.0-16.0); *syrup* (*syrupus rhei*), ℥x-xx (0.6-1.2) for an infant; *aromatic* or *spiced syrup* (*syrupus rhei aromaticus*), for an infant f3ss-j (2.0-4.0); *tincture* (*tinctura rhei*), f3j-ij (4.0-8.0); *aromatic tincture* (*tinctura*

rhei aromaticus), f3ss-j (2.0-4.0); *sweet tincture* (*tinctura rhei dulcis*), f3j-ij (4.0-8.0).

JUGLANS is the *bark* of the *root* of *Juglans cinerea*, butternut or white walnut (*Nat. Ord.* Juglandaceæ), collected in autumn. It is used as a substitute for rhubarb, which it resembles in action. The *extract* (*extractum juglandis*) may be given in doses of gr. x-xxx (0.6-2.0).

ALOE—ALOES.

What is aloes?

Aloes is the *inspissated juice* of the *leaves* of *Aloe Perryi*, socotrine aloes (*aloe socotrina*) and of *Aloe vera*, Barbadoes aloes (*aloe Barbadoensis*) (*Nat. Ord.* Liliaceæ). Each variety contains a neutral crystalline principle, *aloin*. It is a tonic to the mucous membrane and a purgative, being slow in its action and affecting principally the large intestine, and apparently causing a determination of blood to the pelvic viscera. It is also a hepatic stimulant. It is *used* in habitual constipation, especially when torpor of the large bowel exists; in *hemorrhoids* from relaxation of the veins and congestion of the portal circle; in *amenorrhœa* and in *menorrhagia* due to want of tone of the uterine vessels. The official preparations are made of *purified* (socotrine) *aloes* (*aloe purificata*), and are, *pills* (*pilulæ aloes*), each containing gr. ij (0.13) of aloes; *pills of aloes and mastich* (*pilulæ aloes et mastiches*), called also *Lady Webster's dinner-pill*, each containing gr. ij (0.13) of aloes; *pills of aloes and myrrh* (*pilulæ aloes et myrrhæ*) or *Rufus' pill*, each containing gr. ij (0.13) of aloes; *pill of aloes and iron* (*pilulæ aloes et ferri*) each containing gr. j (0.07) of aloes, of dried ferrous sulphate and of aromatic powder (the two last named pills are useful in amenorrhœa); *pills of aloes and asafetida* (*pilulæ aloes et asafetidæ*), each containing gr. j½ (0.09) of aloes, asafetida and soap (useful in flatulent dyspepsia with constipation); *extract* (*extractum aloes*), dose gr. j-v (0.06-0.3); *tincture* (*tinctura aloes*), dose f3j-iv (4.0-16.0); *tincture of aloes and myrrh* (*tinctura aloes et myrrhæ*), dose f3j-ij (4.0-8.0).

ALOINUM (*aloin*) is official, dose gr. 10-j (0.006-0.06).

SENNA. LEPTANDRA. EUONYMUS.

What are these medicines?

SENNA is the *leaflets* of *Cassia acutifolia* (Alexandria Senna) and of *C. angustifolia* (Indian Senna) (*Nat. Ord.* Leguminosæ). It contains *cathartic acid*, *chrysophanic acid*, *sennit* and *sennacrol*. It is a prompt and efficient purgative, and probably acts on the entire length of the intestine, causing some griping. It is a mild hepatic stimulant. It is used where a prompt and decided cathartic effect is desired, usually in combination to prevent griping. Its preparations are, the *fluid extract* (*extractum sennæ fluidum*) dose f3j-iv (4.0-16.0); *confection* (*confectio sennæ*), an excellent purgative during pregnancy, 3j-ij (4.0-8.0); *compound infusion* (*infusum sennæ compositum*), also called *black draught*, f3ss-iv (15.0-120.0); *syrup* (*syrupus sennæ*) f3j-ij (4.0-8.0); *compound powder of glycyrrhiza* (or of *liquorice*) (*pulvis glycyrrhizæ compositus*), an excellent purgative, dose a teaspoonful (4.0) at bedtime.

LEPTANDRA is the *rhizome* and *roots* of *Veronica virginica*, Culver's root or physic (*Nat. Ord.* Scrophularinæ). It contains *leptandrin*, and is a feeble stimulant to the liver and intestinal glands. Dose of the *extract* (*extractum leptandræ*) gr. j-iv (0.06-0.25); of the *fluid extract* (*extractum leptandræ fluidum*) f3ss-j (2.0-4.0).

EUONYMUS, or wahoo, the *bark* of the root of *E. atropurpureus* (*Nat. Ord.* Celastrinæ), contains a bitter principle, *euonymin*, *resins*, *euonic acid*, etc.; is a powerful hepatic stimulant, and increases the intestinal secretion. It is used as a purgative in torpidity of the liver and intestines. Dose of the *extract* (*extractum euonymi*) gr. ij-iv (0.13-0.26) in pill.

MERCURIAL CATHARTICS.

What mercurials are used as simple purgatives?

HYDRARGYRI CHLORIDUM MITE (*mercurous chloride* or *calomel*), dose gr. ss-j to x (0.03-0.6), at bedtime; and MASSA HYDRARGYRI (*mercurial* or *blue mass*), dose gr. v-xv (0.3-1.0). They will be found under the head of "*Alteratives*."

SALINE CATHARTICS.

What are saline cathartics?

The saline cathartics are certain *salts* which produce an evacuation of the bowel not only by increasing peristalsis from their irritant effect, but also by promoting outward osmosis from the blood, thus rendering the stool liquid, depleting the bloodvessels, and consequently favoring absorption of effused fluids.

Name the medicines of this group.

The magnesium salts, some of the sodium and some of the potassium salts, and manganese sulphate.

Describe the effects, uses, preparations, and doses of the magnesium salts.

MAGNESIA (*calcined magnesia*) and MAGNESIA PONDEROSA (*heavy magnesia*—is similar to *Henry's*, *Husband's*, and *Ellis' magnesia*) are antacid and cathartic, are mild in their action and are excellent preparations for children; dose as an antacid gr. xx (1.2), as a cathartic ʒj (4.0) or more.

MAGNESII CARBONAS (*magnesium carbonate—magnesia alba*), not much used; dose as an antacid gr. x-xx (0.6-1.2), as a cathartic ʒj-ij (4.0-8.0).

MAGNESII SULPHAS (*magnesium sulphate—Epsom salt*) is much more powerful than the preceding preparations, and produces free, watery purgation with very little irritation. It is used as a *purgative* to remove *fecal accumulations*; as a *depletant* in *entero-colitis* and *acute dysentery* (with opium); to *aid absorption* (given in very concentrated solutions) in *dropsical effusions*, especially hydrothorax; and for the relief of *colica pictonum* and other forms of lead-poisoning; dose ʒss-j (15.0-31.0).

MAGNESII CITRAS EFFERVESCENS (*effervescent magnesium citrate*), dose ʒj-iv (4.0-16.0) taken in water while effervescing.

LIQUOR MAGNESII CITRATIS (*solution of magnesium citrate*). The citrates are similar in effect to the sulphate, but are more apt to irritate; dose half to a whole bottle full (180.0-360.0 c.c.).

What salts of sodium are used as cathartics?

SODII SULPHAS (*sodium sulphate—Glauber's salt*) is similar in its

effects to Epsom salt, but more irritant and disagreeable. It is a mild hepatic stimulant. It is not much used; dose $\mathfrak{z}\text{j}$ -iv (4.0-16.0).

SODII PHOSPHAS (*sodium phosphate*) is a mild cathartic and hepatic stimulant. It is an excellent purgative in *catarrhal conditions of the bowel*, in *catarrhal jaundice*, and in *chronic infantile diarrhœa* with pasty stools; dose $\mathfrak{z}\text{j}$ - $\mathfrak{z}\text{j}$ (4.0-31.0).

SODII PYROPHOSPHAS (*sodium pyrophosphate*) resembles the preceding salt in action and uses; dose $\mathfrak{z}\text{ss}$ -iv (2.0-16.0).

What potassium salts are used as cathartics?

POTASSII SULPHAS (*potassium sulphate*) was formerly, but is not at present, much used; dose gr. xv- $\mathfrak{z}\text{j}$ (1.0-4.0).

POTASSII BITARTRAS (*potassium bitartrate—cream of tartar—cremor tartar*) is a diuretic, refrigerant and cathartic. In overdoses it causes gastro-enteritis. Dose as *diuretic* gr. x- $\mathfrak{z}\text{j}$ (0.6-4.0), as a *mild laxative* $\mathfrak{z}\text{ss}$ -j (2.0-4.0), as a saline cathartic $\mathfrak{z}\text{ss}$ (16.0).

POTASSII ET SODII TARTRAS (*potassium and sodium tartrate—Rochelle salt*) resembles Epsom salt, but is not so powerful; dose $\mathfrak{z}\text{ss}$ -j (16.0-31.0).

PULVIS EFFERVESCENS COMPOSITUS (*compound effervescing powder—seidlitz powder*) consists of Rochelle salt $\mathfrak{z}\text{ij}$ (7.75) with sodium bicarbonate gr. xl (2.59) in a blue, and tartaric acid gr. xxxv (2.25) in a white paper, their contents to be dissolved separately, the solutions mixed and taken while effervescing, on an empty stomach.

What is manganese sulphate?

MANGANI SULPHAS (*manganese sulphate*) has been discussed with the tonics. It is a hepatic stimulant and cathartic also, resembling Epsom salt, but is a dangerous remedy, and is not much employed; dose gr. j-v (0.06-0.3).

DRASTIC CATHARTICS.

What are drastic cathartics?

Cathartics which cause much irritation and act very powerfully are so called. Most of them in overdoses cause irritant poisoning. When they produce very free, watery evacuations they are sometimes spoken of as "*hydragogues*." The more powerful drastics are used as revulsives in cerebral affections.

Describe the medicines employed as drastics.

JALAPA (*jalap*), the *tuberous root* of *Ipomœa Jalapa* (*Nat. Ord.* Convolvulaceæ), a Mexican plant, contains a *resin*, to which its purgative properties are due.

It is a prompt and powerful hydragogue and cholagogue in overdoses, causing gastro-intestinal irritation and even death. It is *used* as a hydragogue in *dropsies* of cardiac or renal origin, as in ascites or anasarca, and sometimes as a cerebral revulsant. Its preparations are *extract* (*extractum jalapæ*), dose gr. j-x (0.06-0.6); *compound powder* (*pulvis jalapæ compositus*), containing 35 per cent. of jalap with cream of tartar, dose gr. x-3j (0.6-4.0); and the *resin* (*resina jalapæ*), dose gr. j-v (0.06-0.3).

BRYONIA (*bryony*), the *root* of *B. alba* and *B. dioica* (*Nat. Ord.* Cucurbitaceæ), European climbing vines, contains a glucoside, *bryonin*, probably the active principle. It is a violent hydragogue cathartic resembling jalap in its *effects*, and also proving *diuretic*. It is *used* as a hydragogue in dropsies, especially in hydrothorax and hydropericardium. The *tincture* (*tinctura bryoniæ*) may be used in doses of f3ss-j (2.0-4.0).

PODOPHYLLUM, the *rhizome* and *roots* of *P. peltatum*, May-apple or mandrake (*Nat. Ord.* Berberideæ), a common indigenous plant, containing a resinous substance *podophyllotoxin*, consisting of picropodophyllin, the purgative principle, and podophyllinic acid, which is inert.

It is an active hydragogue and cholagogue, acting slowly and especially on the upper part of the small intestine and causing much irritation. In overdoses it is a gastro-intestinal irritant capable of causing death.

It is *used* as a purgative and cholagogue in acute and chronic *constipation*, *bilious attacks* and in *torpor of the liver*. The dose of the *extract* (*extractum podophylli*) is gr. v-xv (0.3-1.0); of the *fluid extract* (*extractum podophylli fluidum*) m̄x-xx (0.6-1.2); of the *resin* gr. $\frac{1}{10}$ - $\frac{1}{2}$ (0.006-0.03); the latter preparation is the only one in common use.

Podophyllotoxin (not official) may be used in the form of a 1 per cent. alcoholic solution, the dose of which is gtt. x-xxx (1.0-3.0). It is used hypodermically, dose gr. $\frac{1}{4}$ (0.016).

IRIS, the *rhizome* and *roots* of *I. versicolor*, or blue-flag (*Nat. Ord. Irideæ*); contains a purgative *resin*. It is a powerful stimulant to the liver and intestinal glands, *acting* like podophyllum, but with less irritation. It is *diuretic*, and in overdoses irritant and depressant. It is *used* in *dropsies*, *jaundice of malarial origin*, and *torpor of the liver*. Dose of the *extract* (*extractum iridis*) gr. ij-iv (0.1-0.25); of the *fluid extract* (*extractum iridis fluidum*) ℥x-3j (0.6-4.0).

CHELIDONIUM, celandine or tetterwort, is the *herb* *C. majus* (*Nat. Ord. Papaveraceæ*); contains *chelerythrine* and *chelidonine* combined with *chelidoninic acid*. It is *used* as a *hydragogue cathartic*, and is supposed to possess *cholagogue effects* and some *narcotic properties*. It may be given in doses of gr. x-3j (0.6-4.0), in a fluid extract or infusion.

SCAMMONIUM (*scammony*) is a *resinous exudation* from the living *root* of *Convolvulus Scammonia* (*Nat. Ord. Convolvulaceæ*), a Syrian plant. The resin *scammonin* is the active portion. It is a *violent hydragogue*, resembling jalap in its action, but much more irritating. It is never given except in combination with other cathartics. The dose of the *resin* (*resina scammonii*) is gr. ij-vij (0.13-0.5). It is one of the ingredients of *compound extract of colocynth*.

COLOCYNTHIS (*colocynth*), the *fruit*, deprived of the rind, of *Citrullus colocynthis*, or bitter cucumber (*Nat. Ord. Cucurbitaceæ*), contains a glucoside (the purgative principle) called *colocynthin*, also a *resin*, *colocynthitin*, etc.

It is an *hepatic stimulant* and *violent hydragogue*, and in overdoses causes death from gastro-enteritis. It is never given alone, but is *used* in small doses to increase the effects of milder cathartics. The *extract* (*extractum colocynthidis*) is used in combination with other purgatives, dose gr. ij-v (0.13-0.3); *compound extract* (*extractum colocynthidis compositum*) containing also aloes, resin of scammony, cardamom and soap, dose gr. iij-x (0.2-0.64).

CAMBOGIA (*gamboge*) is a *gum-resin* obtained from *Garcinia*

Hanburii (*Nat. Ord. Guttiferæ*), a tree of Siam. The resin is the purgative principle and is called *cambogic acid*. It is a very powerful *hydragogue cathartic*, and in overdoses has caused death. It is rarely used alone. Dose gr. ij-v (0.13-0.3).

PILULÆ CATHARTICÆ COMPOSITÆ (*compound cathartic pills*). Each pill contains of compound extract of colocynth gr. $j\frac{1}{2}$ (0.08), of calomel and of extract of jalap each gr. j (0.06) and of gamboge gr. $\frac{1}{4}$ (0.015).

PILULÆ CATHARTICÆ VEGETABILES (*vegetable cathartic pills*). Each pill contains of compound extract of colocynth gr. j (0.06), of extract of hyoseyamus and of jalap each gr. ss (0.03), of extract of leptandrin and of resin of podophyllum each gr. $\frac{1}{4}$ (0.015) with oil of peppermint.

ELATERINUM (*elaterin*) is a *neutral principle* extracted from *Elaterium*, a substance deposited by the juice of the fruit of *Ecballium Elaterium*, or squirting cucumber (*Nat. Ord. Cucurbitaceæ*). It is one of the most powerful of the *hydragogues*, also proving *diuretic*, and in excessive doses often causing death. It is much used in *dropsies*, in *uræmia* and as a cerebral *revulsive*. The dose is gr. $\frac{1}{10}$ (0.003); of the *trituration* (*trituration elaterini*) gr. $\frac{1}{4}$ -ss (0.016-0.03).

OLEUM TIGLII (*croton oil*) is a *fixed oil* expressed from the seeds of *Croton Tiglium* (*Nat. Ord. Euphorbiaceæ*), a tree of India. The active principle has not been isolated.

When applied to the skin croton oil is an intense irritant, causing a papular eruption which soon becomes pustular. Given internally it is a powerful and rapid *hydragogue cathartic*, causing severe griping pains and to some extent depressing the circulation. In overdoses it produces great depression and often proves fatal. In *obstinate constipation* it is very serviceable, and as a *revulsive* in cerebral affections, as apoplexy, it is the best *hydragogue*. It has been used externally as a *counter-irritant* (*q. v.*). Dose gtt. j (0.016-0.12), which may be given in pill or emulsion, or mixed with a few drops of glycerin or olive oil and placed on the patient's tongue when he cannot or will not swallow.

ENEMATA.

What are enemata?

Enemata are liquids injected into the rectum. They may be *nutritive*, *cathartic*, or *forced enemata*.

Nutritive enemata are used to nourish the patient when the stomach cannot retain or digest food. They should be small in amount, not exceeding fʒiij-iv (90.0-120.0) at a time, nor should they be too frequently repeated, else they will cause irritability of the rectum. They should be combined with some digestant, as pepsin or pancreatin.

Cathartic enemata are used to unload the bowels when irritation or inflammation of the stomach is present; to aid the action of cathartics taken by the mouth; to remove fæcal accumulations or retained and irritating secretions from the large intestine; to relieve tympanites; to remove ascarides; or to act as revulsives. The fluid used may be cold or warm water, either simple or medicated, and the quantity will depend on the case. The injection should be made slowly and carefully, and the introduction of air is to be avoided.

Forced enemata are made slowly and gradually, the fluid, either warm water or warmed oil, being poured into a large funnel or rubber bag (fountain syringe) to which a long rubber tube is attached, armed with an injection pipe. By elevating the funnel or bag the fluid flows in a steady stream, the force being regulated by the height at which the bag is held. Forced enemata are employed principally to reduce intussusception of the bowels.

ORDER V.—ANTHELMINTICS.

What are anthelmintics?

Anthelmintics are medicines used to kill (*vermicides*) or cause the expulsion of (*vermifuges*) intestinal worms.

How are they administered?

They should be given fasting, and should be followed in 4 to 6 hours by a purgative, as calomel or castor oil.

Describe the medicines used as anthelmintics.

SPIGELIA, or pinkroot, the *rhizome* and *roots* of *S. marilandica* or Carolina pink (*Nat. Ord.* Loganiaceæ), contains a *bitter principle*,

volatile oil, resin, tannic acid, etc. It slows the heart by stimulating the inhibiting centres, reduces arterial pressure by depressing the heart, and in overdoses quickens the pulse, increases the blood-pressure, dilates the pupil, causes delirium, and, in fatal cases, death from respiratory paralysis. It is very efficient in causing the expulsion of *lumbrici*, but should be followed by a purge. Dose of the *fluid extract* (*extractum spigeliæ fluidum*) f3j-ij (4.0-8.0), or to a child gtt. x (0.5) on sugar.

CHENOPODIUM or American wormseed, the *fruit* of *C. ambrosioides* or Jerusalem oak (*Nat. Ord.* Chenopodiaceæ), contains a *volatile oil*, to which its effects are due. It is a very efficient remedy for the expulsion of *lumbrici*, and should be followed by a brisk cathartic. The *oil* (*oleum chenopodii*) is given in doses of gtt. v-x (0.25-0.5) on sugar, to a child 3 years old, before meals.

SANTONICA or Levant wormseed is the *unexpanded flower-heads* of *Artemisia pauciflora* (*Nat. Ord.* Compositæ). The active principle is *santonin*. *Large doses* cause salivation, rapid breathing, slow, labored pulse, trembling, yellow vision and yellow urine, unconsciousness, convulsions, dilated pupils and death. It is used almost exclusively as a remedy for *lumbrici*. Dose of *santonin* (*santoninum*) for an adult gr. ss-v (0.03-0.3); for a child 2 years old gr. $\frac{1}{4}$ (0.016). *Troches of santonin* (*trochisci santonini*) each contain gr. ss (0.03) of santonin, dose 1 to 6 troches.

ASPIDIUM, the *rhizome* of *Dryopteris filix-mas*, or male fern, and *D. marginale* (*Nat. Ord.* Filices), contains *filicic acid*, *volatile* and *fixed oils*, *resin*, etc., and is used as a *teniacide*. Like all remedies employed for this purpose, its administration should be preceded by a fast of 24 hours and followed by a brisk cathartic. In overdoses it has caused death from enteritis. Dose of the *oleoresin* (*oleoresina aspidii*) f3ss-j (2.0-4.0).

GRANATUM (*pomegranate*), the *bark* of the *stem* and *root* of *Punica Granatum* (*Nat. Ord.* Lythrarieæ), contains four alkaloids, *pelletierine* and three allied alkaloids, which are powerful *teniacides*, as is also granatum itself. In overdoses they are said to act on the muscles like woorara. Dose of pomegranate in decoction (3ij (59.147) to water Oij (946.358) boiled to Oj (473.179)) f3ij (59.147) repeated 3 times at an hour's interval, before breakfast; of *pelletierine tannate* gr. v-x (0.3-0.6).

CUSO (*Kousso*; *Brayera*, Pharm. 1880), the *female inflorescence* of *Hagenia abyssinica* (Nat. Ord. Rosaceæ), contains *protokosin* (a crystalline, inactive resin) and *kosotoxin* (an amorphous, highly active resin), and is used as a *teniacide*; dose of the *fluid extract* (*extractum cusso fluidum*), f3ij-iv (8.0-15.0).

KAMALA, the *glands and hairs* from the capsules of *Mallotus philippinensis* (Nat. Ord. Euphorbiaceæ), contains *rottlerin* and *isorottlerin*, resinoid substances, and is used as a *teniacide*; dose 3j-ij (4.0-8.0) of the powder in syrup, or given in the form of tincture.

PEPO (*pumpkin-seed*), the *seed* of the *Cucurbita pepo* or pumpkin (Nat. Ord. Cucurbitaceæ), contains a *resin*, and is one of the most powerful *teniacides* in the materia medica and perfectly harmless; dose of the fresh seed (powdered with sugar) 3j-ij (30.0-65.0).

OLEUM TEREBINTHINÆ (*oil of turpentine*) is an efficient but not a harmless remedy for *tænia* and *lumbrici*. It will be found under the head of "*Diuretics*." Dose f3ss-j (15.0-30.0) combined with castor oil.

HYDRARGYRI CHLORIDUM MITE (*mercurous chloride or calomel*) is used alone or combined with other anthelmintics for the expulsion of *lumbrici* and (rarely) *tænia*. It should be given in full purgative doses (*v. Alteratives*).

What remedies are used for the destruction of ascarides?

Mild purgatives, as *calomel* or *sulphur*, may be used to dislodge them from the small intestines, but *injections of quassia-infusion*, *common salt*, *naphthol*, *tannin*, *alum* or *lime water* must be employed to get rid of those in the colon. *Injections of sweet oil* at bedtime are useful to kill any which may be in the rectum, and prevent the excessive itching which they cause. The child should be prevented from scratching the anus, as the ova are very tenacious, and adhering to the finger nails may be conveyed to the mouth and reproduce the trouble.

What are useful adjuncts to the anthelmintics?

Iron and the *bitter tonics*, by restoring tone to the mucous membranes, prevent the excessive secretion of mucus which forms a suitable nidus for the development of parasites.

Class III.—Medicines acting on the body by their effect on the various secretions and excretions.

ORDER I.—DIURETICS.

What are diuretics?

Diuretics are medicines used to increase the excretion of urine, acting either *directly* on the secreting cells of the kidney, or *indirectly* by their influence on the blood-pressure in the glomeruli.

For what purposes are they used?

They are used (1) to maintain the *function of the kidneys* in suppression of urine independent of the cause; (2) to promote the excretion of *urinary solids* (injurious waste products) or of poisons from the blood, as in uræmia, gout, rheumatism, fevers, and in cases of various metallic and vegetable poisons; (3) to promote the excretion of *water*, thus tending to dilute highly concentrated urine or to aid in the removal of effused (dropsical) fluids; and (4) to alter the *chemical reaction* of the urine and render that fluid more bland, thus lessening vesical and urethral irritation and any tendency which may exist to the formation of calculi.

A certain relation exists between the functions of the skin, kidneys and bowels, so that free catharsis or free diaphoresis will diminish the urinary secretion and *vicê versâ*.

How may diuretics be classified?

They may be divided into alkaline, hydragogue and alterative diuretics.

ALKALINE DIURETICS.

What are the alkaline diuretics?

They are those salts which, without materially increasing the amount of water, usually increase the elimination of its solid ingredients and modify or alter its chemical reaction.

Name the alkaline diuretics.

The potassium and lithium salts, and strontium lactate.

POTASSIUM.

What is the source of potassium?

Potassium is obtained from the ashes of plants, from native *nitre*, and from *tartar* or *argol*, deposited from wine during fermentation.

What are the general physiological actions of its salts?

Locally applied *caustic potash* destroys tissues by abstracting water, dissolving albumin and saponifying fats. *Internally* in *small doses* before meals, they increase the acidity of the gastric juice; in *large doses*, or after meals, they neutralize the acid present in the stomach. The *bicarbonates*, when taken on an empty stomach, enter the blood readily, and abstracting the base from basic and neutral potassium phosphate convert it into the acid phosphate, thus *increasing the acidity of the urine*; but if taken after or during meals, they are decomposed in the stomach, and entering the blood as a carbonate, *diminish the acidity of the urine*. The *salts formed with the vegetable acids* are converted into carbonates in the system and render the urine less acid. Potassium and its salts promote oxidation (and hence cause emaciation when used continuously), decreasing the amount of uric acid and increasing the urea excreted by the urine. The *nitrate* and *chlorate* are not decomposed in the system, and having effects peculiar to themselves, will be separately considered. All the potassium salts are protoplasmic poisons, in very large doses paralyzing the heart in diastole by direct action on its muscle and contained ganglia. They also depress and paralyze the functions of the brain and spinal cord. They all prove diuretic, increasing the amount of water passed and, by promoting oxidation, the elimination of urea. *Poisoning by caustic potash* is due to its corrosive properties, and if recovery takes place from the acute effects, various stenoses are apt to follow. The *carbonate* in overdoses causes violent gastro-enteritis with great cardiac depression and sometimes paralysis. The *treatment* of potassium poisoning consists in neutralizing the alkali by a vegetable acid, giving demulcents and fixed oils as protectives, and stimulating if requisite.

What are the therapeutical uses of the potassium salts?

The potassium salts are used as *diuretics* (increasing as they do both the water and salts of the urine, and converting uric acid into

the more soluble urea as well as rendering the urine alkaline) in *acute rheumatism*, *uric acid diathesis*, in *desquamative nephritis*, and to alkalinize the urine in *cystic* and *urethral irritation*.

What are the potassium preparations, their uses and doses ?

LIQUOR POTASSÆ (*solution of potassa*) contains 5 per cent. of potassium hydrate, and is *antacid* and *diuretic*. It is used to render the urine alkaline in cystic irritability and in gonorrhœa; dose ℥x-xx (0.6-1.2) freely diluted.

POTASSII CARBONAS (*potassium carbonate*) is *antacid* and *diuretic*, but is more irritating to the stomach than the bicarbonate; dose gr. x-xx (0.6-1.2).

POTASSII BICARBONAS (*potassium bicarbonate*) is *antacid* and *diuretic*; dose gr. xx-℥j (1.2-4.0). It is much used in the treatment of *acute rheumatism*, especially in plethoric persons (℥jss (45.0) are given in the first twenty-four hours, half as much during the second day, and afterward enough to keep the urine alkaline).

POTASSII CITRAS (*potassium citrate*) is used as a *diuretic* in acute rheumatism and as a refrigerant diaphoretic in fevers and acute inflammations; dose gr. xx (1.2) in solution every hour or two.

LIQUOR POTASSII CITRATIS (*solution of potassium citrate*) contains 8 per cent. of the bicarbonate and 6 per cent. of citric acid. It is often prescribed with sweet spirits of nitre, in fever mixtures; dose f℥ss (15.0).

POTASSII CITRAS EFFERVESCENS (*effervescent potassium citrate*) is made by mixing thoroughly, drying and powdering citric acid (63.0), potassium bicarbonate (90.0), and sugar (47.0). Dose ℥ss-j (2.0-4.0).

Neutral mixture (not official) is made by neutralizing fresh lemon-juice, with the bicarbonate. The official *solution of potassium citrate* has superseded it, as being the more exact and more elegant method of prescribing the same medicine; dose f℥ss-j (15.0-30.0).

Effervescing draught (not official) is made by adding f℥j (29.57) of a solution of the bicarbonate (℥j (3.88)-f℥ij (88.7)) to f℥j (29.57) of lemon-juice and water (equal parts). It is particularly acceptable to febrile patients with irritable stomachs.

POTASSII ACETAS (*potassium acetate*) is *antacid*, *diuretic*, and in large doses a *mild cathartic*. It is chiefly used as a *diuretic* in acute rheumatism and to some extent in dropsies, and in uric acid diathesis; dose gr. xx-℥j (1.2-4.0).

POTASSII BITARTRAS (*potassium bitartrate—cream of tartar*) is a *diuretic* and *saline cathartic* (*q. v.*) much employed in dropsies.

POTASSII NITRAS (*potassium nitrate—saltpetre—nitre*) locally applied is a powerful irritant to the mucous membranes and to raw surfaces. It passes through the system unchanged, and is eliminated chiefly by the urine, increasing the amount passed but not affecting its reaction to the same extent as the other salts. *Poisoning* occurs more frequently from the nitrate than from any other potassium salt, the symptoms being burning pain in the throat and epigastrium, hæmatemesis and bloody stools, muscular weakness, syncope, collapse sometimes preceded by convulsions or paralysis, suppression of urine and death. The *treatment* consists in evacuating the contents of the stomach and administering opiates, demulcents and stimulants.

It is *used in acute rheumatism* as a *diuretic*, but is probably inferior to the salts formed with the vegetable acids. It is used in *fevers* for the same purpose and has been given in small doses combined with Dover's powder as a *diaphoretic* in *acute muscular rheumatism*. Dose of potassium nitrate gr. x-xxx (0.6-1.9). *Papers of potassium nitrate* (*charta potassii nitratis*), consisting of strips of unsized paper immersed in a 20 per cent. solution of the salt, are burned and the fumes inhaled with success in spasmodic asthma. It is probable that the *nitrite* is formed by the combustion and is the active agent in these cases.

POTASSII CHLORAS (*potassium chlorate*) is a *local irritant*. *Internally* in large doses it causes gastro-enteritis, affects the circulation as do the other potassium salts, damages the blood corpuscles and interferes with oxygenation by the formation of methæmoglobin, and produces a condition of acute tubal nephritis. Cases of poisoning are not infrequent, and death is usually preceded by profuse salivation, great dyspnoea, cyanosis, delirium, coma, tonic and clonic spasms, and sometimes ecchymoses and jaundice. It increases the excretion of urine, but does not affect its chemical reaction.

It has been *used* as a diuretic in dropsies, but is dangerous and inferior to other preparations. As an *alterative* in the exanthemata it is now little employed. In *mercurial, follicular* or *aphthous stomatitis* it is of unquestionable value, given internally with sugar or used as a gargle. In ordinary *sore throat* and in *tonsillitis* it is

useful as a local application; dose gr. v-xxx (0.3-2.0). It must be given freely diluted. It should not be prescribed in powder with readily oxidizable or combustible substances, as when triturated with them it forms explosive compounds. As a gargle ℥ss-iv (2.0-15.0) in water Oj (473.1) may be used.

The *troches (trochisci potassii chloratis)* each contain gr. v (0.3) of the salt.

The other potassium preparations have been or will be considered in their appropriate place.

LITHIUM.

What are the sources of lithium?

Lithium is a chemical element, found native in the form of silicates, phosphates, etc.

What are the physiological effects and uses of the lithium salts?

The lithium salts appear to resemble those of potassium. They are eliminated by the urine as urates, increasing the urine and rendering it much more strongly alkaline than a corresponding amount of a potassium salt. As the urate of lithium is much more soluble than either the urate of sodium or of potassium, the lithium preparations are advantageously *used* in the treatment of *chronic gout* and *lithæmia*, and as a *local application* to gouty deposits either before or after the skin has broken. It has also been employed internally in *glycosuria*.

What are the preparations of lithium and their doses?

LITHII CARBONAS (*lithium carbonate*), dose gr. v-x (0.3-0.6) freely diluted. As a local sorbefacient for gouty deposits a solution (gr. v (0.3)-f℥j (30.0)) should be kept constantly applied.

LITHII CITRAS (*lithium citrate*), dose gr. v-xx (0.3-1.2).

LITHII CITRAS EFFERVESCENS (*effervescent lithium citrate*) is made by triturating and thoroughly drying citric acid (370.0) with sugar (about 200.0); then triturating the mixture with lithium carbonate (70.0), sodium bicarbonate (280.0) and sugar (enough to make the product weigh 1000.0). Dose ℥j-ij (4.0-8.0).

LITHII BENZOAS (*lithium benzoate*), dose gr. v-xx (0.3-1.2).

Lithium bromide and *L. salicylate* have been *previously mentioned*.

What sodium salt has been used as a diuretic?

SODII ACETAS (*sodium acetate*) is supposed to possess properties similar to those of potassium acetate, and has been used for the same purposes and in the same doses.

What preparation of strontium is used as a diuretic?

STRONTII LACTAS (*strontium lactate*) has been used as a diuretic, especially in *acute desquamative nephritis*, with scanty urine, where it has been found to increase the amount of urine and diminish the albumin present in that excretion. It has also been used in *other forms of kidney disease*, but is much less efficient in chronic than in acute cases. It has been used as an *intestinal antiseptic*, but is of little value for this purpose. Dose gr. v–xv (0.3–1.0) in solution.

HYDRAGOGUE DIURETICS.

What are hydragogue diuretics?

The medicines of this group greatly increase the flow of the *watery* element of the urine, and are hence used in the treatment of dropsies.

Name the hydragogue diuretics?

Squill, digitalis, caffeine, sodio-theobromine salicylate, apocynum, broom, dandelion, sweet spirit of nitre, and mercurials.

SCILLA—SQUILL.

What is squill?

Squill is the *bulb* of *Urginea maritima* (*Nat. Ord. Liliaceæ*) deprived of its dry, membranous outer scales, and cut into thin slices, the central portions being rejected. It contains *scillipicrin*, *scillitoxin*, and *scillin*. In *small doses* squill increases the mucous secretions and is a stimulating expectorant, while it markedly increases the urinary secretion. In *large doses* it causes nausea, vomiting, and sometimes purging, and in *excessive doses* produces in addition great abdominal pain, with scanty, bloody, or even suppressed urine, great depression, collapse, convulsions and death.

Scillipicrin appears to be the diuretic principle. *Scillitoxin* slows the pulse, and in overdoses causes diastolic cardiac arrest by stimulating the peripheral vagi; after large doses the vagi are finally paralyzed and the pulse becomes rapid. *Scillin* is devoid of diuretic effects, but appears to be the emetic principle.

The *treatment of poisoning* consists of evacuating the contents of the stomach and the use of opiates and demulcents. If collapse occurs alcohol must be employed.

Squill is used as a *diuretic* in *dropsy*, but should not be employed if kidney disease or cystitis is present. It is frequently combined with digitalis in cardiac dropsies, and in chronic pleurisy or pericarditis is often given with calomel. As an *expectorant* it is much used in *bronchial* affections. As an *emetic* it is too dangerous a remedy to employ.

As a *diuretic* or *expectorant* squill may be given in doses of gr. j-ij (0.06-0.13) repeated every 2 hours and gradually increased until nausea occurs. Doses of gr. vj-xij (0.389-0.778) generally prove emetic. The dose of the *fluid extract* (*extractum scillæ fluidum*) is ℥j-ij (0.06-0.12); of the *tincture* (*tinctura scillæ*) ℥x-xxx (0.6-1.8); of the *vinegar* (*acetum scillæ*) ℥x-xxx (0.6-1.8); of the *syrup* (*syrupus scillæ*) fʒss-j (2.0-4.0); of the *compound syrup* (*syrupus scillæ compositus*—*Coxe's hive syrup*), containing squill and senega and gr. j (0.059) of tartar emetic in each ounce (29.5), ℥v-fʒj (0.3-4.0).

DIGITALIS—CAFFEINE.

Are these drugs used as diuretics?

The uses of these remedies as diuretics have already been discussed under the head of “*Cardiac Stimulants*.” For this purpose the best preparations of digitalis are the *infusion*, the *powdered leaves* or the *tincture*.

SODIO-THEOBROMINE SALICYLATE.

What is sodio-theobromine salicylate?

Sodio-theobromine salicylate, or *diuretin* (the *proprietary name*), is a non-official preparation “obtained by mixing aqueous solutions of equal molecules of sodium theobromine and salicylic acid, and evaporating to dryness; a definite compound appears to be formed containing theoretically 49.7 per cent. of theobromine and 38.1 per cent. of salicylic acid. It occurs as a white powder, odorless, of a saline, alkaline taste, and soluble in one-half its weight of warm water, the solution remaining perfect on cooling.”*

* National Dispensatory.

Its physiological action is not fully understood. It is a powerful *diuretic* in certain diseased conditions, as when *anasarca* is present, due to disease of the heart or kidneys, but appears to have no diuretic effect when given to healthy persons.

It has little if any effect upon the *circulation* except in the diseased conditions above mentioned, when it appears to increase the force and render more regular the beat of the heart, probably by increasing diuresis and removing the fluid which has been acting as a mechanical obstruction to the circulation.

The action on the *kidneys* is not understood, but is non-irritating in character. Under its use the amount of *solid* as well as the *fluid* of the urine is much increased, the latter sometimes enormously, as much as three gallons of urine having been reported as passed in 24 hours while using the drug. As the dropsical accumulation diminishes, the amount of urine lessens.

Sometimes sodio-theobromine salicylate causes "nervousness," headache or vertigo, tinnitus aurium, and palpitation. Occasionally a rise in temperature is seen after its use. In rare cases a cutaneous eruption attended with more or less intense itching appears. More frequently, however, it acts as a gastro-intestinal irritant, causing loss of appetite, nausea, and sometimes severe vomiting and purging.

The *therapeutical uses* of this drug are indicated by what has been already said. It is of great value in all *dropsical effusions* depending on disease of the kidney or heart, and should always be tried in these conditions when remedies like digitalis, strophanthus, etc., have not succeeded in reducing the dropsy. In pleuritic effusions it is not so certain in its action. In ascites due to portal obstruction it usually appears to produce no effect. In dropsies dependent on tubercular inflammation it is absolutely valueless. Dose gr. xv (1.0) repeated every 2 or 3 hours until some effect is produced. The author has never found it necessary to give more than 3jss (6.0) in 24 hours, although some observers have prescribed double that quantity in the same period of time without ill effect. It is administered in solution in plain water flavored with peppermint or other aromatic. It may be given in capsules.

APOCYNUM.

What is apocynum?

Apocynum, Canadian hemp, or milk-weed, is the *root* of *A. cannabinum* (*Nat. Ord.* Apocynaceæ) and contains *apocynin* and *apocynëin*, a glucoside. These substances are supposed to be cardiac tonics, acting on the heart similarly to digitalis. They are also diuretic. Apocynum is laxative in *small*, but emetic and cathartic in large doses. It is chiefly *used* as a diuretic in dropsies, and should not be given in quantities sufficient to purge when the former effect is desired; *extractum apocyni fluidum* (*fluid extract of apocynum*), dose $\mathfrak{m}_{\text{V-xxx}}$ (0.3–2.0).

SCOPARIUS.

What is scoparius?

Scoparius is the *tops* of *Cytisus Scoparius* or broom (*Nat. Ord.* Leguminosæ), and contains a volatile liquid alkaloid, *sparteine* (*v.* “*Cardiac Stimulants*”), and a neutral principle, *scoparin*, in which the diuretic properties are supposed to reside. Broom is an efficient hydragogue diuretic, and in large doses produces free purging and sometimes vomiting. It is much *used* in the treatment of *dropsies* and is best given in decoction (\mathfrak{J}_{ss} (14.7) to water Oj (473.1) boiled down to Oss (236.5), dose $\mathfrak{f}\mathfrak{J}_{\text{j}}$ (30.0) every 2 or 3 hours until an effect is produced) or the *fluid extract* (*extractum scoparii fluidum*) may be given in doses of $\mathfrak{f}\mathfrak{J}_{\text{ss-j}}$ (2.0–4.0).

TARAXACUM.

What is taraxacum?

It is the *root* of *Taraxacum officinale* or dandelion (*Nat. Ord.* Compositæ) gathered in the autumn, and contains a bitter principle, *taraxacin*; also *taraxacerin*, *inulin*, resin, etc.

It is *diuretic*, laxative, and slightly cholagogue, and is *used* in dropsies of hepatic origin and dyspepsia with torpor of the liver. Dose of the *extract* (*extractum taraxaci*) gr. xx– \mathfrak{J}_{j} (1.28–4.0), of the *fluid extract* (*extractum taraxaci fluidum*) $\mathfrak{f}\mathfrak{J}_{\text{j-ij}}$ (4.0–8.0).

SPIRITUS ÆTHERIS NITROSI—SPIRIT OF NITROUS ETHER.

Is this medicine a diuretic?

Sweet spirit of nitre is diuretic in large doses if the surface of the body is kept cool. It is usually combined with other diuretics, but

may be given alone where the urinary secretion is scanty from simple congestion or from purely functional causes. It will be considered more fully under "*Diaphoretics*." Dose f3j-iv (4.0-15.0) freely diluted.

MERCURIALS.

Are the mercurial preparations ever used as diuretics?

Calomel (*q. v.*) has been employed as a diuretic in cardiac dropsy, dose gr. iss 4 or 5 times a day. It is a useful adjunct to squill in hydrothorax and hydropericardium.

Corrosive sublimate will probably prove to be a more powerful diuretic, as by its action on the liver it increases the formation of urea, which of itself always increases the excretion of water by the urine.

ALTERATIVE DIURETICS.

What are alterative diuretics?

They are medicines which are employed not only as diuretics but also for their effect on the mucous membrane of the urinary tract, by which they are possibly to some extent excreted, or on which they exercise an alterant and sedative effect during their passage in the urine.

Name the medicines belonging to this group.

Buchu, pareira, uva ursi, chimaphila, triticum, juniper, oil of erigeron, oil of santal, stigmata maydis, turpentine, copaiba, cubeb, matico, cantharides.

BUCHU. PAREIRA.

What are these medicines?

BUCHU, the *leaves* of *Barosma betulina* and other species (*Nat. Ord. Rutaceæ*), natives of S. Africa, contains a *volatile oil* and a *bitter principle*. It is a mild tonic and diuretic, and slightly stimulates the genito-urinary mucous membranes, having also an alterative effect upon them.

It is *used* in chronic catarrhal conditions of the genito-urinary mucous tract, as *chronic pyelitis*, *chronic cystitis* and especially in irritation of the neck of the bladder; dose of the *fluid extract* (*extractum buchu fluidum*) f3ss-j (2.0-4.0).

PAREIRA, the *root* of *Chondodendron tomentosum* (*Nat. Ord. Menispermaceæ*), a Brazilian plant, contains a bitter alkaloid, *cissampeline*, and has been used with success in the treatment of *chronic cystitis*, *gleet* and "*irritable bladder*." It is said to be also tonic and laxative. Dose of the *fluid extract* (*extractum pareiræ fluidum*) f3ss-j (2.0-4.0).

UVA URSI. CHIMAPHILA.

What are these medicines?

UVA URSI is the *leaves* of *Arctostaphylos Uva ursi* or bearberry (*Nat. Ord. Ericaceæ*), and contains *tannic* and *gallic acid*, a glucoside called *arbutin*, *ericolin* (a bitter substance), etc. It is used as a mild astringent tonic, and diuretic in *chronic pyelitis* and *cystitis*, and has also been employed in *chronic bronchitis*.

Arbutin is the diuretic principle, and as it is, partially at least, eliminated as hydroquinone (*vid. "Antipyretics"*), it is supposed to act as an antiseptic to the urogenital tract. It has been employed in the above diseases and internally in *gonorrhœa*.

Dose of the *fluid extract* (*extractum uva ursi fluidum*) f3ss-ij (2.0-8.0), t. i. d.; of the *extract* (*extractum uva ursi*) gr. v-x (0.3-0.9).

CHIMAPHILA, the *leaves* of *C. umbellata*, pipsissewa, wintergreen, or ground holly (*Nat. Ord. Ericaceæ*), contains *arbutin*, *tannic acid*, *chimaphilin*, etc. It is tonic, astringent, and diuretic, resembling buchu and uva-ursi in its effects and uses. Special alterative qualities have been claimed for it in scrofulous affections.

Dose of the *fluid extract* (*extractum chimaphilæ fluidum*) f3ss-j (2.0-4.0), t. i. d.

TRITICUM.

What is triticum?

Triticum, the *rhizome* of *Agropyrum repens* or couch grass (*Nat. Ord. Gramineæ*), contains *triticin* and several *sugars*, and is feebly diuretic. It is useful in *cystitis* and in irritation of the neck of the bladder.

Dose of the *fluid extract* (*extractum tritici fluidum*) f3ss-ij (2.0-8.0).

JUNIPERUS—JUNIPER.

What is juniper?

Juniperus (juniper) (not official) is the *fruit* (or berries) of *J. communis* (*Nat. Ord. Coniferæ*), and contains a *volatile oil* (*oleum juniperi*, dose gtt. v–xv (0.3–0.9)), on which its diuretic effects depend. It is a stimulant to the kidneys, capable in overdoses of causing strangury and bloody urine, and is sometimes used as an emmenagogue. It is chiefly employed as an adjuvant to other diuretics, particularly the alkaline group. As an *alterative diuretic* it is used in *chronic pyelitis* and *chronic cystitis*. It is sometimes given in *subacute renal congestion*, but is contraindicated where active inflammation is present. It is occasionally used as a *carminative* and *stomachic*. The preparations (official) are the *spirit* (*spiritus juniperi*), containing 5 per cent. of the oil, dose ʒj–ij (4.0–8.0); the *compound spirit* (*spiritus juniperi compositus*) contains also the oils of caraway and fennel in alcohol and water, and is about the same strength as *gin*, which is often used as a substitute for it, dose fʒj–jv (4.0–15.0).

OLEUM ERIGERONTIS—OIL OF ERIGERON.

What is the oil of erigeron?

It is the *oil distilled* from the fresh flowering *herb* of *Erigeron canadense*, or Canada fleabane (*Nat. Ord. Compositæ*), and yields on distillation a *resin* and a *terpene*. It is *diuretic* and *hemostatic*, and is useful in hemorrhages due to relaxation of the genito-urinary mucous membrane. It has been used with success in *gonorrhœa*. Dose ʒv–xxx (0.3–2.0) on sugar or in capsule, every two or three hours.

OLEUM SANTALI—OIL OF SANTAL.

What is the oil of santal?

It is a *volatile oil* distilled from the *wood* of *Santalum album* (*Nat. Ord. Santalacæ*). It is an *alterative*, *diuretic*, and *blennorrhetic*, sometimes causing vertigo, when its use should be discontinued or the dose decreased. Occasionally it produces gastrointestinal irritation. It is used in the second stage of *gonorrhœa*, and in *chronic bronchitis* it is often beneficial; dose ʒv–xx (0.3–1.2) in emulsion or capsule, t. i. d.

ZEA.

What is this substance?

Zea, or corn silk, the *styles* and *stigmas* of Zea Mays or Indian corn (*Nat. Ord.* Gramineæ), is a mild, stimulating diuretic, and is strongly recommended in *cystitis*, both acute and chronic, in *gonorrhœa*, in *irritability of the bladder* depending upon a highly concentrated urine, in *nocturnal incontinence* in children, and even in *cardiac dropsy*. It is given in infusion ad libitum, or in the form of *fluid extract* (*extractum zee fluidum*), of which the dose is f3j-ij (4.0–8.0) every hour or two.

TEREBINTHINA—TURPENTINE.

What are the officinal varieties of turpentine?

1. TEREBINTHINA (*turpentine* or *white turpentine*) is a *concrete oleoresin* obtained from *Pinus palustris*, yellow or pitch pine, and other varieties of *Pinus* (*Nat. Ord.* Coniferæ) trees indigenous to the Southern States.

2. TEREBINTHINA CANADENSIS (*Canada turpentine*, *Canada balsam* or *balsam of fir*) is a *liquid oleoresin* obtained from *Abies balsamea*, the American silver fir, or balm of gilead tree (*Nat. Ord.* Coniferæ), found in the northern portions of N. America.

3. Unofficinal varieties known to commerce are *Chian turpentine* derived from *Pistacea Terebinthinus* (*Nat. Ord.* Anacardaceæ), formerly used as a specific for internal (especially uterine) cancer, but possessing no curative power; *Bordeaux turpentine*, *Venice turpentine*, etc.

What do the turpentines contain?

They contain a *resin* or *rosin* (*resina*) united to a *volatile oil* (*oleum terebinthinæ*).

What are the effects and uses of turpentine?

The *effects* of turpentine are those of the oil which it contains. They are rarely *used* internally, but have been given as an alternative to the broncho-pulmonary and genito-urinary mucous membranes in chronic catarrhal conditions, and have been used as an anthelmintic, but for all these purposes the oil is preferred.

Dose as a *diuretic* and *blennorrhetic* gr. xx-3j (1.3–4.0), in pill or emulsion; as an *anthelmintic* ʒss-j (15.5–31.0).

RESINA (*resin*) is only used externally as a basis for plasters and ointments. Its preparations are a *cerate* (*ceratum resinæ*), sometimes called basilicon ointment, and a *plaster* (*emplastrum resinæ*), adhesive or sticking plaster.

OLEUM TEREBINTHINÆ—OIL OF TURPENTINE.

What is the oil of turpentine?

Oil of turpentine or spirit of turpentine is a *volatile oil distilled* from turpentine.

What are its effects and uses?

Locally it is rubefacient. *Internally* it is carminative, in large doses causing mental exhilaration and increasing the force and frequency of the pulse and proving cathartic. It is eliminated by the lungs and kidneys, imparting a violet-like odor to the urine. In *excessive doses* it is capable of causing death (rare). The symptoms of poisoning are unconsciousness, bloody, often suppressed urine, and rapid, feeble pulse. Vomiting and strangury or purging sometimes occur and occasionally the pupils are dilated. Small doses seem to increase the pulse rate by direct action on the *heart*, while large doses stimulate the vagi and slow the heart. Moderate doses are diuretic, but when large amounts are given renal irritation results. It is *used* as an *alterative diuretic* in *chronic cystitis*, *gleet* and *chronic pyelitis*. In *impotence* and *spermatorrhœa* depending on relaxation it is sometimes employed. In *typhoid pneumonia* and *bronchitis*, and in *hypostatic pulmonary congestion*, as in *typhoid fever*, it is often used externally and internally with the happiest effect. It is very useful in *ulceration of the stomach* or *intestines*, and in *typhoid fever* when the tongue becomes brown and dry or cracked, red and glazed, or when the ulcers are very slow to heal; it acts not only locally but also as a cardiac stimulant. In *hemorrhages* from the lungs, stomach or bowels, in *post-partum hemorrhages* and in *purpura hemorrhagica* it is sometimes useful. As an addition to stimulating enemata to cause the expulsion of flatus or to rouse the system to shock it is much resorted to. It is also used as an *anthelmintic*.

Locally as a counterirritant, either as a liniment or fomentation, it is much used in *bronchitis* and in *muscular rheumatism*, and in the

latter manner in *peritonitis*, *hysterical abdominal pains* and in *flatulency*. Dose $\mathfrak{m}\mathfrak{v}$ -xv (0.3-0.9) 4 or 5 times daily in emulsion or capsule; as a cathartic enema or anthelmintic $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$ -j (15.0-30.0) may be given. Externally a *liniment* (*linimentum terebinthinæ*) is employed, or the oil may be used either full strength or diluted with sweet oil. A saturated solution of camphor in oil of turpentine is a good rubefacient liniment.

OLEUM TEREBINTHINÆ RECTIFICATUM (*rectified oil of turpentine*), dose $\mathfrak{m}\mathfrak{v}$ -xxx (0.3-1.8).

COPAIBA.

What is copaiba?

Copaiba is the *oleoresin* of *Copaiba Langsdorffii* and other species (*Nat. Ord.* Leguminosæ), trees of S. America. It is cathartic, and sometimes emetic, in large doses, slightly stimulates the circulation and acts as an alterative blennorrhetic on the mucous membranes by which it is excreted. After large doses a resinous precipitate will occur if nitric acid be added to the urine, which, however, is soluble in alcohol. When the *resin* is taken in considerable quantities the urine responds to the copper tests for sugar. *Copaiba* is chiefly eliminated by the genito-urinary mucous membranes and to a slight extent by the lungs. Its use is sometimes followed by urticaria. It is *used* in *chronic genito-urinary inflammations*, especially in *gonorrhœa* after the subsidence of the acute stage. In *bronchorrhœa* it is sometimes employed. The *oil* has been lauded as a hydragogue diuretic in dropsies when not of renal origin. The *dose* of *copaiba* is $\mathfrak{m}\mathfrak{x}$ -3j (0.6-3.6) or more in emulsion, pill, or capsule. The *mass* (*massa copaibæ*—*solidified copaiba*) contains 94 per cent. with magnesia, dose gr. v-xxx (0.3-2.0); of the *oil* (*oleum copaibæ*) the dose is $\mathfrak{m}\mathfrak{v}$ -xv (0.3-0.9) in emulsion or capsule.

CUBEBA—CUBEÆ.

What is cubeb?

Cubeb is the *unripe fruit* of *Piper Cubeba* (*Nat. Ord.* Piperacæ), an East Indian plant. It contains a *volatile oil*, a neutral principle (*cubebin*) and *cubebic acid*. Cubeb is diuretic and carminative, slightly accelerating the pulse. In large doses it often causes vomiting and purging. It is eliminated by the mucous membranes and

urine, increasing the excretion of uric acid. Its use is sometimes followed by urticaria. The addition of nitric acid to the urine after the use of cubeb produces a resinous precipitate.

It is *used* in *gonorrhœa* and is less apt than *copaiba* to disorder the digestion. It is also used in *bronchitis*, and has been smoked in cigarettes to relieve *sore throat* and *hoarseness*. The powder, used as a snuff, is employed in *coryza*. The *dose* of powdered cubeb is \mathfrak{z}_{ss} - \mathfrak{ij} (2.0–11.6); of the *oil* (*oleum cubebæ*) \mathfrak{m}_{x-xv} (0.6–0.9) gradually increased. It is not as efficacious as the *oleoresin* (*oleoresina cubebæ*), dose \mathfrak{m}_{v-xxx} (0.3–2.0) given in emulsion. Of the *tincture* (*tinctura cubebæ*) the dose is $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ - \mathfrak{ij} (4.0–8.0); of the *fluid extract* (*extractum cubebæ fluidum*) $\mathfrak{f}\mathfrak{z}_{ss}$ - \mathfrak{j} (2.0–4.0). The troches (*trochisci cubebæ*) each contain gr. ss (0.04) of the oleoresin.

MATICO.

What is matico?

Matico is the *leaves* of *Piper angustifolium* (*Nat. Ord. Piperaceæ*), a shrub of Peru. It contains a *volatile oil*, *resin*, *artanthic* and *tannic acids*, etc. It is *used* as an *alterative blennorrhetic* in *gonorrhœa*, as a *hemostatic* in *internal hemorrhages*, and locally as a *styptic*. Dose of the *fluid extract* (*extractum matico*) $\mathfrak{f}\mathfrak{z}_{ss}$ - \mathfrak{j} (2.0–4.0); of the *tincture* (*tinctura matico*) $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ - \mathfrak{ij} (4.0–8.0).

CANTHARIS—CANTHARIDES.

Is cantharides used as an alterative diuretic?

Cantharides is an intense stimulant and irritant to the kidneys, being diuretic in small doses, and frequently causing strangury, priapism and bloody urine. It is sometimes used in *gleet*, and in *spermatorrhœa* and impotence from weakness. It will be noticed at length under "*Epispastics*." Dose of the *tincture* (*tinctura cantharidis*) gtt. v-x (0.3–0.6) t. i. d.

ORDER II.—DIAPHORETICS.

What are diaphoretics?

Diaphoretics are medicines which increase the secretion of sweat, either by (1), stimulating the secretory nerve fibres in the glands; (2) stimulating the nerve centre governing the secretion of sweat;

or (3) relaxing the skin, dilating the blood vessels and regulating the force of the circulation.

For what purposes are they used and how are they classified?

They are used (1) to abort an oncoming disease (as a catarrh or inflammation of mucous or serous membranes after exposure to cold) or aid in its subsidence (as in the paroxysm of a malarial fever); (2) to aid absorption, as in dropsies, and (3) to eliminate poisons from the blood.

They are divided into Nauseating, Sedative, Saline and Special Diaphoretics.

NAUSEATING DIAPHORETICS.

What are the nauseating diaphoretics?

The act of emesis is accompanied by general relaxation and diaphoresis, and even in nausea a more or less pronounced perspiration occurs. Any of the emetics in doses just sufficient to nauseate will produce this effect, but the only one which is in general use for this purpose is IPECAC (v. "*Emetics*"), which is used either alone (in doses of gr. ss-j every two hours) or combined with opium in the form of DOVER'S POWDER (v. "*Opium*") in the early stages of *inflammations*, in "*cold*," in *muscular rheumatism* and in acute *suppression of the menses* to break up the train of morbid symptoms.

SEDATIVE DIAPHORETICS.

What are sedative diaphoretics?

When the circulation is much excited, as in fevers and inflammations generally, medicines which tend to reduce the force and rapidity of the circulation and to lower the temperature, act as diaphoretics. Many of these agents (particularly *antimonials*, *aconite* and *veratrum viride*) tend to nauseate or even to cause vomiting when given in full doses. In *sthenic inflammations* the ANTIMONIAL PREPARATIONS (q. v.) especially if combined with minute doses of morphine are often very efficacious, and may be used in the *early stages* of *laryngitis* and *bronchitis* and in combination with other remedies in fever. ACONITE and VERATRUM VIRIDE are also sometimes resorted to in the early stages of sthenic inflammations for their com-

bined sedative and sudorific effect. In this group also should be placed SALICYLIC ACID and other ANTIPYRETICS (*q. v.*) which produce more or less profuse perspiration.

SALINE DIAPHORETICS.

What are the saline diaphoretics ?

While some of the saline diaphoretics have a sedative effect upon the circulation, which is not, however, as marked as the remedies of the preceding group, they have no tendency to nauseate even when given in large doses, on the contrary, they are acceptable to the stomach, and form the basis of most fever mixtures.

POTASSIUM CITRATE and its preparations have been noticed among the "*Diuretics.*"

LIQUOR AMMONII ACETATIS—SOLUTION OF AMMONIUM ACETATE.

What is the solution of ammonium acetate ?

The *solution of ammonium acetate* or *spirit of mindererus* is a mild stimulant, diaphoretic and diuretic, often employed in febrile and inflammatory affections combined with other remedies. Dose $\text{f}\overline{3}\text{ss-j}$ (15.0–30.0)

SPIRITUS ÆTHERIS NITROSI—SPIRIT OF NITROUS ETHER.

What is this substance ?

Spirit of nitrous ether, or *sweet spirit of nitre*, is a solution of about 4 per cent. of nitrous ether (ethyl nitrite) in alcohol. In *small doses* it is a diaphoretic and stimulant, in *larger doses* an antispasmodic, and (if the skin be kept cool) diuretic; in *excessive doses*, either internally or when inhaled, it is a depresso-motor, acting similarly to the other nitrites (*q. v.*) and capable of causing death. It is much *used* as a diaphoretic in *fevers*, and occasionally as a diuretic (*q. v.*) and antispasmodic in nervous children. Dose for an *adult* $\text{f}\overline{3}\text{j-iv}$ (4.0–15.0). It is given as a diaphoretic in minute doses, frequently repeated and largely diluted, the body being kept warm.

SPECIAL DIAPHORETICS.

PILOCARPUS.

What is pilocarpus?

Pilocarpus or *Jaborandi* is the *leaflets* of *P. Selloanus* (Rio de Janeiro *Jaborandi*) and of *P. Jaborandi* (Pernambuco *Jaborandi*) (*Nat. Ord.* Rutaceæ), a Brazilian shrub.

What are its chief chemical constituents?

It contains an alkaloid, *pilocarpine*, and a *volatile oil*; a second alkaloid, *jaborine*, probably a derivative of pilocarpine, has been isolated, and lately two other derivative alkaloids *pilocarpidine* and *jaborandine* have been described.

What are the physiological effects of these substances?

In *medicinal doses* *pilocarpus* or *pilocarpine* produces profuse diaphoresis and salivation, sometimes copious, with flushed face and quickened circulation and respiration. Sometimes the mucous secretions are increased also, and nausea, even vomiting, may occur. It increases both the watery portion and the solids (especially urea) of the *perspiration* by a direct influence on the nerve endings governing its secretion, and probably increases the secretion of the *gastric juices* and *bronchial mucus* in the same manner. It appears to act directly as a stimulant upon the cells of the salivary glands. There seems to be some relation between its influence over the secretions of perspiration and saliva, the one being more profuse when the other is scanty, and *vice versâ*. In small doses frequently repeated it proves diuretic. It seems to stimulate the nutrition of the hair. The cardiac action is at first quickened, but soon slowed, and the arterial tension is elevated. Applied to the eye it contracts the pupil (by peripheral action) and increases the tension of accommodation. *Pilocarpine* and *atropine* are antagonists as regards their action on the heart, intestines and pupil, and on the secretions of perspiration and saliva. *Jaborine* acts upon the heart, pupil, intestines and salivary glands like atropine. *Pilocarpidine* causes profuse diaphoresis, salivation, vomiting and purging; while *jaborandine* resembles atropine in its cardiac effects.

What are the medicinal uses of pilocarpus and its alkaloid?

Pilocarpus or *pilocarpine* is used as a *diaphoretic* to abort disease

in incipient catarrh and at the onset of a paroxysm of malarial fever; in various forms of *dropsy*, to aid in removal of the fluid, especially *hydrothorax*, and in *pulmonary œdema*; in *uræmia*, to eliminate the poison from the blood, it is of great value, and similarly in *acute* or *chronic Bright's disease*, to relieve the kidneys of some of their work, it is very useful. It is also employed in *subacute* and *muscular rheumatism*.

As a *diuretic*, minute doses (pilocarpine gr. $\frac{1}{12}$ (0.005), are given in *acute suppression of urine*, but when thus used in the early stages of acute desquamative nephritis it must be employed with great caution. As an *expectorant* it has been given in *bronchitis* and *asthma* and, with varying success, in *croup* and *diphtheria*. To stimulate the growth of hair in *alopecia*, pilocarpine may be administered internally or hypodermically, and applied locally as well, and it has been employed hypodermically with success to arrest severe and prolonged hiccough.

It should *not be given* in diseases of the gastro-intestinal mucous membrane, nor in weak heart due to cardiac disease.

What are the officinal preparations of pilocarpus and their doses?

EXTRACTUM PILOCARPI FLUIDUM (*fluid extract of pilocarpus*),
Jōse f3ss-j (2.0-4.0).

PILOCARPINÆ HYDROCHLORAS (*pilocarpine hydrochlorate*), dose
gr. $\frac{1}{8}$ -ss (0.008-0.032). It is well adapted to hypodermic use.

ORDER III.—EXPECTORANTS.

What are expectorants?

Expectorants are remedies which are given to modify the secretions of, and promote their expulsion from, the air passages.

How may they be classified?

They are divided into sedative and stimulating expectorants?

SEDATIVE EXPECTORANTS.

What are sedative expectorants?

They are remedies which increase the secretion and at the same

time tend to lower the blood pressure, cause nausea and, if frequently repeated, vomiting.

Name the sedative expectorants.

Ipecac, apomorphine, tartar emetic, pilocarpine, lobelia and grindelia. Of these ipecac and apomorphine have been discussed with the "*emetics*," tartar emetic with the "*sedatives*," pilocarpine with the "*diaphoretics*," and lobelia with the "*depresso-motors*."

GRINDELIA.

What is grindelia?

Grindelia is the leaves and flowering tops of *G. robusta* and *G. squarrosa* (*Nat. Ord. Compositæ*). It appears to depress the motor nerves and muscles, and in large doses to slow the heart and reduce the blood-pressure. Toxic doses produce narcosis with dilated pupils. It is eliminated by the kidneys, increasing the flow of urine, and to some extent by the broncho-pulmonary mucous membrane. It has been highly recommended in *spasmodic* and *cardiac asthma*, in *hay asthma* and *hay fever*, in *whooping-cough* and in *acute* and *chronic bronchitis*. In *chronic cystitis* it has been advantageously employed, and as a *local* application in *vaginitis* and in *poisoning* by rhus toxicodendron has met with some success. Dose of the *fluid extract* (*extractum grindeliæ fluidum*) m_x -f3j (0.6-4.0).

STIMULATING EXPECTORANTS.

What are stimulating expectorants?

They are remedies which probably modify the secretion by toning up the broncho-pulmonary mucous membrane through which they are eliminated. Most of them seem to exert an alterative effect in passing through the membrane, and many of them elevate the arterial tension.

Name the stimulating expectorants.

Ammonium chloride, senega, quillaja, quebracho, asclepias, eriodictyon, garlic, sanguinaria, ammoniac, benzoin, Peruvian balsam, balsam of Tolu, squill, eucalyptus, serpentaria, turpentine, tar, terpin hydrate, and terebene.

AMMONII CHLORIDUM—AMMONIUM CHLORIDE.

What is ammonium chloride?

Ammonium chloride, called also *ammonium muriate* or *sal ammoniac*, is a crystalline salt obtained from gas liquor.

What are its effects and uses?

Locally it is irritant. In medicinal doses it is believed to influence especially the mucous membranes, promoting very decidedly nutritive changes and hastening epithelial exfoliation. When given continuously for a length of time it diminishes the solids of the blood and stimulates the secretions. It is supposed to stimulate the hepatic functions, for the amount of urea excreted is greatly increased, as are all the other solids of the urine except uric acid. Large doses are purgative. It is used as a *stimulating* and *alterative expectorant* in the moist stage of *acute* and in *chronic bronchitis*, in *pneumonia* and the various *catarrhs* of children. Abroad it is employed also in *gastric* and *intestinal catarrhs*, in *chronic hepatitis* and in the early stages of *hepatic cirrhosis*. In cases of *albuminoid liver*, *spleen* or *kidneys* it is very useful. In subacute *muscular rheumatism* and in *neuralgia*, especially ovarian, it is advantageous, and it has been used internally as an alterative and employed locally as a sorbefacient in scrofulous *adenitis* of the *cervical glands*, and *uterine fibroid tumors*.

Dose gr. x-xxx (0.6-2.0). The unpleasant taste may be partially disguised by liquorice. The *troches* (*trochisci ammonii chloridi*) each contain about gr. ij (0.1) of the salt.

SENEGA.

What is senega?

Senega is the *root* of *Polygala Senega* or *senega snake-root* (*Nat. Ord.* Polygalææ) and contains *polygalic acid* or *senegin*, a glucoside identical with *saponin*.

What are its effects and uses?

Locally, it is an irritant. *Internally*, it stimulates the mucous membranes, and in large doses is emetic and cathartic. It is given as a *stimulating expectorant* in the latter stages of *acute* and in subacute and *chronic bronchitis*, especially when the mucous membranes are wanting in tone. It has also been used as an *emmenagogue* in

amenorrhœa and as a *diuretic* in dropsies. Dose of the *fluid extract* (*extractum senegæ fluidum*) $\text{m}_{\text{x-xx}}$ (0.6–1.2); of the *syrup* (*syrupus senegæ*) $\text{f}\text{3j-ij}$ (4.0–8.0). It is an ingredient of the *compound syrup of squill*.

QUILLAJA.

What is quillaja?

Quillaja, or soap-bark, is the *inner bark* of *Q. Saponaria* (*Nat. Ord. Rosaceæ*), a S. American tree, containing *saponin*, which foams like soap when in infusion. *Locally*, *saponin* is an irritant and protoplasmic poison. Given *internally* in sufficient doses it is a nerve and muscle poison, and completely antagonizes the effects of *digitalis* on the heart, which it paralyzes in diastole. It increases the secretion of bronchial mucus. *Quillaja* has been used as an *expectorant* in *chronic bronchitis*, *emphysema*, and *dilated bronchi*. It is also used to aid in emulsifying oils. Dose of the *tincture* (*tinctura quillajæ*) $\text{m}_{\text{xv-xxx}}$ (0.9–2.0).

ASPIDOSPERMA.

What is aspidosperma?

Aspidosperma, or *Quebracho*, is the *bark* of *A. Quebracho-blanchi* (*Nat. Ord. Apocynaceæ*). It contains 5 alkaloids, *aspidospermatine* (which is believed to represent the entire virtues of the bark), and *aspidospermine*, *quebrachine*, *hypoquebrachine*, and *quebrachamine*. It has been used in *dyspnœa* independent of cause, but as it appears to have no action on the heart or kidney it probably is of service only in *bronchial asthma* and *emphysema*. It has also been used as a *bitter tonic*. Dose of the *fluid extract* (*extractum aspidospermatis fluidum*) $\text{m}_{\text{xv-f3j}}$ (1.0–4.0).

ASCLEPIAS.

What is asclepias?

Asclepias, or *pleurisy root*, is the *root* of *A. tuberosa* (*Nat. Ord. Asclepiadeæ*). It is an *expectorant*, *diaphoretic*, and *diuretic*, and causes considerable cardiac depression. In large doses it is a *gastro-intestinal irritant*. It is but little used. Dose of the *fluid extract* (*extractum asclepiadis fluidum*) $\text{m}_{\text{xv-xxx}}$ (1.0–2.0).

ERIODICTYON.

What is eriodictyon?

It is the *leaves* of *E. glutinosum*, or California tar-bush (*Nat. Ord.* Hydrophyllaceæ), and is known also as *yerba santa*. It is used as an *expectorant* in *chronic pulmonary disorders*, and as a *vehicle* (in the form of elixir) to disguise the taste of quinine and other bitter drugs. Dose of the *fluid extract* (*extractum eriodictyi fluidum*) \mathfrak{M}_{xx-f3j} (1.2-4.0).

ALLIUM—GARLIC.

What is garlic?

Garlic is the *bulb* of *Allium sativum* (*Nat. Ord.* Liliaceæ) and contains a very volatile *oil*. *Locally* applied it is a rubefacient. Taken *internally* it is a condiment, stimulating digestion, increasing the secretions generally, especially the broncho-pulmonary, and quickening the circulation. It is *used* as a *stimulating expectorant* in subacute and chronic *bronchitis* and in *acute bronchitis of infants*, especially when the disease begins to assume the capillary form. Externally it is used as a counter-irritant in the above diseases, and in infantile convulsions, applied to the spine, legs and feet in the form of a poultice. Dose of the *syrup* (*syrupus allii*) $f3j$ (4.0) for a child.

SANGUINARIA.

What is sanguinaria?

Sanguinaria or blood-root, the *rhizome* of *S. canadensis* (*Nat. Ord.* Papaveraceæ), contains an alkaloid, *sanguinarine*. It is a *local irritant*. *Large doses* cause vomiting with great depression, a slow, irregular pulse (from direct action on the heart) and lowered blood-pressure, dilated pupil, collapse and death from paralysis of the respiratory centre, sometimes preceded by convulsions of spinal origin and lowered reflex activity. *Moderate doses* increase the arterial pressure from stimulation of the vaso-motor centres, which are paralyzed by larger doses. It stimulates the liver, the mucous membranes and the salivary and intestinal glands. It is used chiefly as a *stimulating expectorant* in lingering cases of *bronchitis*. Dose of the *tincture* (*tinctura sanguinaria*) \mathfrak{M}_{v-xxx} (0.3-2.0); of the *fluid extract* (*extractum sanguinarie fluidum*) \mathfrak{M}_{ij-v} (0.12-0.3). *Sanguinarine* (not official) has also been used as an expectorant in the dose of gr. $\frac{1}{12}-\frac{1}{8}$ (0.005-0.008).

AMMONIACUM—AMMONIAC.

What is ammoniac?

Ammoniac is a *gum-resin* obtained from *Dorema Ammoniacum* (*Nat. Ord. Umbelliferae*). It is used internally as a *stimulating expectorant* in *chronic bronchitis* and in *bronchorrhœa*, and locally as a *counterirritant*. Dose gr. x-xxx (0.6-2.0); of the *emulsion* (*emulsum ammoniaci*—formerly called *the mixture*, Pharm. 1880) fʒss (15.0). A *plaster* is official (*emplastrum ammoniaci cum hydrargyrô*—*plaster of ammoniac with mercury*).

BENZOINUM—BENZOIN.

What is benzoïn?

Benzoïn, a *balsamic resin* obtained from *Styrax Benzoïn*, or *Benjamin tree* (*Nat. Ord. Styraceae*) contains *benzoic acid*, *resin*, *volatile oil*, and often *cinnamic acid*.

The effects of benzoïn are due to *benzoic acid* (v. “*Antiseptics*”). *Locally* it is an irritant. Internally it stimulates the broncho-pulmonary and genito-urinary mucous membrane. It is eliminated chiefly by the urine as hippuric acid, increasing the acidity of that excretion.

It is chiefly *used* by inhalation (compound tincture ʒj(4.0), in a cupful (120.0) of boiling water,) in *laryngitis*, or to quiet spasmodic laryngeal cough, as in influenza. It has also been employed internally in *chronic bronchitis* and in *chronic cystitis with alkaline urine and phosphatic deposits*, but for the latter purpose benzoic acid or its salts are preferred. It is added to ointments to prevent rancidity, and is much employed in the form of fumigating pastilles. It is also used locally as an *antiseptic and protective* (*compound tincture*) in *fissured nipple*, *chapped hands*, and *fissure of the anus*, and has even been employed as a substitute for carbolic and salicylic acids, for its antiseptic effects, in the *treatment of wounds*. Dose of the *tincture* (*tinctura benzoïn*) fʒss-j (2.0-4.0); of the *compound tincture* (*tinctura benzoïn*i *composita*) fʒss-ij (2.0-8.0).

Benzoïnated lard (*adeps benzoïnatus*) contains 2 per cent. of benzoïn.

BALSAMUM PERUVIANUM—BALSAM OF PERU. BALSAMUM TOLUTANUM—BALSAM OF TOLU.

What are these substances?

BALSAM OF PERU is a *balsam* obtained from *Toluifera Pereiræ* (*Nat. Ord. Leguminosæ*), and contains *cinnamein*, *resin*, *benzalcohol*, *benzylic benzoate*, *stilbene*, and *cinnamic* and *benzoic acids*.

It is a stimulant and tonic to the mucous membranes, and has been used in various catarrhs. Dose f3ss (2.0) in emulsion.

BALSAM OF TOLU is a *balsam* obtained from *Toluifera Balsamum* (*Nat. Ord. Leguminosæ*) and contains *resins*, *benzylic benzoate* and *cinnamate*, *tolene*, and *cinnamic* and *benzoic acids*. Its effects and uses are similar to those of balsam of Peru, to which it is preferred because of its more pleasant taste. Dose gr. x-xxx (0.6-2.0) in emulsion, every few hours. It is, however, principally used as a vehicle. Dose of the *tincture* (*tinctura toluiani*) is f3j-ij (4.0-8.0); of the *syrup* (*syrupus toluianus*) f3j-iv (4.0-15.0). It is also an ingredient of the *compound tincture of benzoin*.

PIX LIQUIDA—TAR.

What is tar?

Tar is an *empyreumatic oleoresin* obtained by destructive distillation of the wood of *Pinus palustris* and other species of *P.* (*Nat. Ord. Coniferæ*). When distilled it yields *pyroligneous acid*, *oil of tar*, and a black residuum known as *pitch*. The *oil of tar* contains *creasote*. Tar resembles turpentine in its effects on the mucous membranes, but is more *antiseptic*. *Externally* it is an *irritant*. It is used as a *stimulating expectorant* in the later stages of *acute bronchitis*, particularly where the disease does not yield readily to other treatment, and in *chronic bronchitis*. *Externally* it is employed in *chronic skin diseases*, especially in squamous affections.

Tar-water (not official) may be given in doses of f3ss-j (15.0-30.0) or the *syrup* (*syrupus picis liquidæ*), which is better, in doses of f3ss (15.0). *Locally*, the *oil* (*oleum picis liquidæ*) is used as a substitute for "oil of cade" (*oleum cadinum*), which is also official, or the *ointment* (*unguentum picis liquidæ*) may be applied.

TERPINI HYDRAS—TERPIN HYDRATE.

What is terpin hydrate?

It is the *hydrate* of the diatomic alcohol *terpin*. It occurs in colorless, rhombic, lustrous prisms; is odorless, but has a slightly aromatic, rather bitter taste.

It *acts* on the *nervous system* and on *mucous membranes* in a manner similar to turpentine. It is eliminated by the urine, to which it imparts a turpentine odor. When given in large doses or for a long time, it may cause strangury, albuminuria, or even hæmaturia.

From its stimulating influence on mucous membranes it has been much *used* as an *expectorant* in *bronchitis*, either acute or chronic, especially when the secretion is very free.

It has also been used in *cystitis*, in *gonorrhœa*, and in *chronic kidney disease*. Dose gr. j-ijj (0.06-0.2) every 4 hours, in pill or capsule.

TEREBENUM—TEREBENE.

What is terebene?

Terebene is a liquid consisting chiefly of *pinene*, and containing not more than very small proportions of terpinene and dipentene, and is prepared by gradually adding sulphuric acid to the oil of turpentine. In its general effects it resembles turpentine. It is a good *stimulating expectorant*, very useful in the declining stages of *acute* and *chronic bronchitis*. It is also useful in *emphysema*, *phthisis*, and *pleurisy*. In *fetid bronchitis* and *bronchiectasis* it is very efficacious. It has also been employed in *flatulent dyspepsia* and *genito-urinary inflammations*. Dose ℥v-xx (0.3-1.2) in capsule or emulsion t. i. d., or its vapor may be inhaled diluted with steam.

What substances are often used as expectorants which have been previously considered in other medicinal groups?

Squill and turpentine (v. "*Diuretics*"), eucalyptus, serpentaria, and Prunus Virginiana (v. "*Aromatic Bitters*").

ORDER IV.—ASTRINGENTS.

What are astringents?

Astringents are remedies employed to cause contraction of the tissues with which they come in contact and thus lessen secretion.

For what purposes are they used?

They are used to check profuse secretion (*e. g.*, ptyalism, etc.) or morbid discharges (*e. g.*, diarrhoea), to overcome relaxation of the tissues, and as hæmostatics and styptics.

How may the astringents be classified?

They are divided into vegetable and mineral astringents.

VEGETABLE ASTRINGENTS.

What are the vegetable astringents?

They are tannic and gallic acids, and the various plants which are used as astringents on account of containing these principles.

Name the vegetable astringents.

Tannic acid, gallic acid, nutgalls, catechu, kino, krameria, hæmatoxylon, white-oak, geranium, witch-hazel, red and white rose, sumach, chestnut leaves.

ACIDUM TANNICUM—TANNIC ACID.

What is tannic acid?

Tannic acid (gallo-tannic acid—digallic acid) is an *organic acid* obtained from nutgall. It is widely distributed throughout the vegetable kingdom and exists under two forms, *gallo-tannic acid* (which is the official variety and strikes a blue-black color with the ferric salts) and *kino-tannic acid* (which produces a greenish-black color with the same salts). Gallo-tannic acid is a light yellowish, amorphous powder, usually cohering in the form of glistening scales or spongy masses, without smell, but having a strongly astringent taste.

It is freely soluble in water and in alcohol, and may readily be dissolved in glycerin by the aid of moderate heat.

What are its physiological effects?

Tannic acid is a powerful astringent, coagulating albumins, contracting blood-vessels and constricting tissues with which it comes in contact. It checks the secretions of the alimentary canal when taken internally, and restrains peristalsis, thus tending to constipate. It is converted into an alkaline tannate and into gallic acid in the *prima via* and slowly absorbed. It is eliminated by the fæces and urine under its own form and as gallic and pyrogallic acids.

For what purposes is it used in medicine?

When *remote* astringent effects are desired gallic acid is preferable, but tannic acid is employed whenever the part to be acted on is so situated that the drug can come in direct contact with it. Thus, it is given to *arrest morbid discharges* from the bowels in *diarrhœa* and *dysentery*, both internally and by enema, and is used externally for the same purpose in *leucorrhœa*, *gleet*, *colliquative sweating*, and as an application to *chronic ulcers*. To *counteract relaxation of tissues* it is employed in the treatment of *spongy gums*, *chronic pharyngitis* and *laryngitis*, *mercurial stomatitis* and hemorrhoids. It is used also to *arrest hemorrhages* from the nose, gums, pharynx, stomach and bowels, and locally as a styptic; to *harden tissues* subjected to pressure, as in threatened bed-sores and tender feet, and as an *antidote to antimony* and other metallic poisons.

Dose, in diarrhœa, gr. j-xv (0.06-1.0) in pill; in hemorrhage, gr. v-xx (0.3-1.3) in powder. The *troches* (*trochisci acidi tannici*) each contain gr. j (0.06), and may be used in relaxed palate or sore throat.

The *ointment* (*unguentum acidi tannici*) or the *glycerite* (*glyceritum acidi tannici*), a solution in glycerin, may be used locally.

ACIDUM GALLICUM—GALLIC ACID.

What is gallic acid?

Gallic acid is an *organic acid*, usually prepared from *tannic acid*, and is tannic acid + a molecule of water; thus, $\text{HC}_{14}\text{H}_9\text{O}_9$ (tannic acid) + $3\text{H}_2\text{O} = 2(\text{HC}_7\text{H}_5\text{O}_5 + \text{H}_2\text{O})$ (gallic acid). It occurs as white or pale fawn-colored, silky, interlaced needles or triclinic prisms; odorless, having an astringent or slightly acidulous taste. It is very soluble in boiling alcohol and boiling water, less so in cold alcohol, glycerin, ether, or cold water. It does not coagulate albumin, and hence is not as efficacious as a local astringent, but is preferred to tannic acid when it is desirable that the astringent should be absorbed and act through the blood on some remote part. It is used as a *hemostatic* in *hemoptysis*, *hematuria*, and *uterine hemorrhage*. It is also employed to *lessen secretion*, as in *colliquative sweats* and *bronchorrhœa*, and to *check morbid discharges*, as in *chronic albuminuria*. Dose gr. v-xxx (0.3-2.0), in powder, pill, or solution.

Describe the preparations, doses, and uses of the vegetable astringents which owe their virtues to tannic and gallic acids.

GALLA (*nutgall*) is an *excrecence* found on *Quercus lusitanica*, the gall oak (*Nat. Ord.* Cupuliferæ) caused by the punctures and deposited ova of a fly (*Cynips Quercus folii* or *Diplolepis Gallæ tinctoriæ*). Before the maturity of the larvæ and their escape from the gall, they contain *tannic acid* (50–60 per cent.) and *gallic acid* (3 per cent.); afterward they are much inferior and contain comparatively little of these substances.

They are but little used except as the source of tannic and gallic acids. The *tincture* (*tinctura gallæ*) is chiefly used as a chemical test. The *ointment* (*unguentum gallæ*) combined with powdered opium is sometimes applied to hemorrhoids.

CATECHU, an *extract* prepared from the *wood* of *Acacia Catechu* (*Nat. Ord.* Leguminosæ), contains *kino-tannic* and *catechuic acids* (*catechin*). It is a powerful astringent, much used in *diarrhæas* and to some extent in *hemorrhages*, and locally as an *astringent* in sore throat, as an injection in *gonorrhœa* and *leucorrhœa*, and as a *styptic*. Dose gr. x–xxx (0.6–2.0) in emulsion or capsules. Of the *compound tincture* (*tinctura catechu composita*) f3j–ij (4.0–12.0) may be given. The *troches* (*trochisci catechu*) each contain gr. j (0.06).

KINO, the *inspissated juice* of *Pterocarpus Marsupium* (*Nat. Ord.* Leguminosæ) and other trees; contains *kino-tannic acid*, *kinoïn*, *pyrocatechin*, etc., and has similar properties and uses to catechu. Dose of the powder gr. x–xxx (0.6–2.0); of the *tincture* (*tinctura kino*) f3j–ij (4.0–8.0).

KRAMERIA, the *root* of *K. triandra* and of *K. Ixina* (*Nat. Ord.* Polygalæ), S. American shrubs; contains *kramero-tannic acid*, and is a powerful astringent, similar to kino and catechu. Dose of the *extract* (*extractum krameriæ*) gr. v–xv (0.3–1.0); of the *fluid extract* (*extractum krameriæ fluidum*) f3ss–j (2.0–4.0); of the *tincture* (*tinctura krameriæ*) f3ss–ij (2.0–8.0); of the *syrup* (*syrupus krameriæ*) f3j–jv (4.0–15.0). The *troches* (*trochisci krameriæ*) each contain gr. j (0.06).

HÆMATOXYLON, logwood or Campeachy wood, is the *heart wood* of *H. campeachianum* (*Nat. Ord. Leguminosæ*), containing *tannic acid*, *hæmatoxylin*, etc. It is a good astringent in diarrhœas due to relaxation, especially in children, but the discharges will stain the linen blood-red. Of the *extract* (*extractum hæmatoxyli*) the dose is gr. x-xxx (0.6-2.0).

QUERCUS ALBA, or white oak (*Nat. Ord. Cupuliferæ*). The *bark* is official, and contains *querci-tannic acid*. It is occasionally used in *diarrhœas*, but more often as an enema in *hemorrhoids* and *prolapsus ani*, or as a vaginal wash in *leucorrhœa* or relaxed conditions with tendency to cystocele or rectocele. It makes an indelible stain on linen.

GERANIUM, the *rhizome* of *G. maculatum*, crow's foot or crane's bill (*Nat. Ord. Geraniacæ*); contains *tannic* and *gallic acids*, and is an efficient indigenous astringent. Dose of the *fluid extract* (*extractum geranii fluidum*) f ʒss-j (2.0-4.0).

HAMAMELIS, the *leaves* of *H. virginica* or witch hazel (*Nat. Ord. Hamamelacæ*); contains *tannic acid* and a *bitter principle*. It is an astringent and seems to contract the veins of the skin and mucous membranes. It has been used in passive hemorrhages as a hemostatic, as an astringent enema in hemorrhoids, and locally in congestions and inflammations. Dose of the *fluid extract* (*extractum hamamelidis fluidum*) f ʒss-j (2.0-4.0).

ROSA GALLICA (*red rose*); ROSA CENTIFOLIA (*pale rose*), (*Nat. Ord. Rosacæ*). Their *petals* are official. The former is a very mild astringent, but they are principally used as flavoring ingredients and vehicles. The preparations of red rose are a *fluid extract* (*extractum rosæ fluidum*), a *honey* (*mel rosæ*), a *syrup* (*syrupus rosæ*) and a *confection* (*confectio rosæ*); those of pale rose are a *triple* or *stronger rose water* (*aque rosæ fortior*), and another milder *water* (*aque rosæ*) from which is made an *ointment* (*unguentum aque rosæ—cold cream*).

RHUS GLABRA (*sumach*) (*Nat. Ord. Anacardiæ*). The *fruit*

is official. It is a good astringent in sore mouth and pharyngitis. The *fluid extract* (*extractum rhois glabræ fluidum*) may be diluted with two or more parts of water and used as a gargle.

CASTANEA, the *leaves* of *C. dentata* or chestnut (*Nat. Ord.* Cupuliferæ), is astringent and supposed to possess antispasmodic properties. It is chiefly used in whooping-cough. Dose of the *fluid extract* (*extractum castaneæ fluidum*) f3ss-ij (2.0-8.0).

MINERAL ASTRINGENTS.

Name the mineral astringents.

Most of the metallic salts possess astringent properties, but those especially used for this purpose are the lead salts ; alum ; aluminium sulphate ; the salts of copper, of zinc, of silver, and of bismuth ; and cerium oxalate.

PLUMBUM—LEAD.

What is the source of lead ?

Lead is obtained by roasting *galena*, the native sulphide.

What are its physiological actions ?

Metallic lead is inert, as are probably also the *sulphide* and *sulphate*. *All other salts of lead*, when *applied locally* in solution, are astringent, or if the solution be strong, irritant. In medicinal doses they are astringent to the alimentary canal, decreasing its secretions and also the secretion of bile. They are but slowly absorbed. When taken for some time they reduce the volume and frequency of the pulse by stimulation of the cardio-inhibitory centre and the peripheral filaments of the vagi. They are *eliminated* by the liver and kidneys.

What are its toxicological actions ?

In large doses several of the lead salts are capable of causing *acute poisoning*, the symptoms being intense gastro-intestinal irritation, vomiting, coma, paralysis and collapse.

Chronic poisoning is caused by saturation of the system with lead, either by drinking pure water conveyed in lead pipes (the contained air forming a soluble oxide), constantly eating substances adulterated with lead salts, exposure in lead factories, etc. *Lead colic* (or

painters' colic), the most common form, consists of sharp pains about the umbilicus, obstinate constipation, hard, contracted abdomen, and often nausea and vomiting. This is probably due to intense stimulation of the intestinal ganglia by the lead, causing arrest of peristalsis and spasm of the muscular coat of the bowel. It is quickly relieved by belladonna, atropine or pilocarpine. Sometimes the poisoning takes the form of *lead-arthritis*, manifesting itself by severe pain in the joints and limbs, and cramps.

Lead paralysis is loss of voluntary motion in groups of muscles, particularly the flexors of the upper extremity, causing *wrist-drop*. Electro-muscular contractility is affected, the muscles waste, and often anæsthesia of the affected or opposite side is present. The poison may also affect the brain, kidney, or other organs. An important symptom of lead poisoning is a *blue line upon the gums where they join the teeth*, and in case of doubt lead may be detected in the urine by the chemical tests.

What are the tests for lead, and how should saturnine poisoning be treated?

Sulphuretted hydrogen gives a *black*, and a solution of *potassium iodide* a *yellow precipitate* with soluble lead salts. The *treatment of acute poisoning* consists in evacuating the contents of the stomach, and administering diluted sulphuric acid, or sodium or magnesium sulphate as a chemical antidote. In *chronic poisoning*, discover its source and remove it (in the case of painters, insist on scrupulous cleanliness of the hands and nails, and the use of very diluted sulphuric acid as a drink); give purgatives to overcome the resulting constipation; give diluted sulphuric acid or the soluble sulphates, with baths of potassium sulphide as antidotes, and aid elimination by large quantities of diluents and potassium iodide. In *lead colic* atropine is of service, and in *lead palsies* electricity (the faradaic current when the muscles respond) and strychnine are advantageously employed.

For what medicinal purposes are the lead salts used?

They are used *internally* as *astringents* in *diarrhœas* and *dysenteries*, as *hemostatics*, and in the treatment of *internal aneurism*. Locally, as *astringent injections* in *leucorrhœa* and *gonorrhœa*, and

as *sedative* and *astringent* lotions in inflammations. The *acetate* is the only lead salt which is administered internally.

What are the preparations of lead and their uses ?

PLUMBI OXIDUM (*lead oxide*), or *litharge*, is occasionally used as a dusting powder for ulcers, but is chiefly employed to make a *plaster* (*emplastrum plumbi*), which is the basis for most official plasters; *diachylon ointment* (*unguentum diachylon*), useful in skin diseases, and *soap-plaster* (*emplastrum saponis*).

PLUMBI ACETAS (*lead acetate*), or *sugar of lead*, is used *internally* as an *astringent* in *diarrhœa*, often combined with opium, and is also useful in *dysentery*. To lessen the expectoration in *chronic bronchitis* and to check the *night sweats of phthisis* it is sometimes given. As a *hemostatic* in yellow fever it does some good, and perhaps also in *hemoptysis*. It has been employed as a *sedative* in the treatment of internal *aneurism*. *Locally* as a *sedative* and *astringent* lotion it is much used in inflammations (not applicable to ocular therapeutics, as it may cause corneal opacity), and to check morbid discharges. Dose gr. j-vij (0.06-0.5) two or three times a day. As an injection in gonorrhœa gr. ss-ij (0.03-0.12)-f℥j (30.0); as a lotion in inflammation gr. v-x (0.3-0.6)-f℥j (30.0).

LIQUOR PLUMBI SUBACETATIS (*solution of lead subacetate*—*Goulard's extract*) contains about 25 per cent. of lead subacetate in solution, and is used externally and in the preparation of *lead water* (*liquor plumbi subacetatis dilutus*—*diluted solution of lead subacetate*), which contains 3 per cent. of the stronger solution; and of *Goulard's cerate* (*ceratum plumbi subacetatis*—*cerate of lead subacetate*).

PLUMBI IODIDUM (*lead iodide*). It may be used internally, dose gr. ij-ijj (0.13-0.19), but is chiefly employed externally in the form of *ointment* (*unguentum plumbi iodidi*), which contains 10 per cent. of the salt.

PLUMBI NITRAS (*lead nitrate*) is used as a disinfectant (*Ledoyen's disinfecting fluid* contains ℥j to the f℥j—about 4.0 to 30.0), *astringent* and *sedative*, in wounds, ulcers, cracked nipples, *onychia maligna*, etc.

PLUMBI CARBONAS (*lead carbonate*—*white lead*) is used chiefly as an application to *fresh burns*. If applied to a large surface, care

should be taken to guard against poisoning. An *ointment (unguentum plumbi carbonatis)* is official, and contains 10 per cent. of the carbonate.

ALUMEN—ALUM.

What is alum ?

Official *alum* is a double sulphate of aluminum and potassium. There are other varieties in which potassium is replaced by some other base, as ammonium or sodium.

What are its effects and uses ?

Alum is a powerful astringent, and in large quantity an irritant, whether used locally or internally. In large doses it is emetic, and in excessive doses has caused death from gastro-intestinal irritation. It is *used* in solution as an *astringent*, by atomization in *bronchorrhœa*, by injection in *leucorrhœa*, *gonorrhœa* and *gleet*, as a gargle in *sore-throat*, *relaxed palate* and *uvula*, as a lotion in *conjunctivitis* and on cotton as a tampon in *cystocele*, *rectocele* and *uterine prolapse*. As a *hemostatic* in *hemoptysis*, a saturated solution may be used by atomization ; and as a *styptic* (*aqua pagliari* consists of alum, compound tincture of benzoin and water boiled together) it is employed locally to arrest hemorrhage. As an *emetic* it is used in *croup* in *capillary bronchitis* and as an adjuvant to other emetics. It is also efficient in the treatment of *lead colic*. Dose as an *astringent* gr. x-xxx (0.6-2.0) ; as an *emetic* ʒj-iv (4.0-16.0). In lead-poisoning a teaspoonful (4.0) may be given every 3 or 4 hours.

Dried alum (alumen exsiccatum) is alum deprived of its water of crystallization by heat. In excessive doses it has proved fatal. Dose gr. v-x (0.3-0.6). Externally it is applied as a mild escharotic to fungous granulations.

ALUMINI SULPHAS—ALUMINUM SULPHATE.

What is aluminum sulphate ?

It is a powerful *astringent* and *antiseptic*, and is employed for these purposes *externally* and as an *injection in gonorrhœa*. Its solution has been used to preserve cadavers for dissection.

CUPRUM—COPPER.

What are the sources and the effects of copper?

Copper is obtained from copper pyrites, a double sulphide of iron and copper. *The metal* itself is inert. *The salts* are mild *astringents* and *stimulants* when in weak solution, and when strong, act as *irritants*; in powder they are *mild caustics*. Taken *internally in small doses* they are *astringent*, and are supposed to act as tonics to the nervous system. In *larger doses* they are local *emetics*. In *excessive amounts* they act as *irritant poisons*, causing gastro-intestinal inflammation, hæmaturia, sometimes jaundice, convulsions, paralysis, delirium, coma and death. The *post-mortem* reveals fatty degeneration of the liver and decomposition of the blood corpuscles. When taken continuously for a *very long time* a condition of chronic poisoning results, with giddiness, occasional diarrhœa, cramps, wasting, and sometimes progressive palsies, respiratory failure and death. A *green line* is sometimes observed on the gums.

The *tests* for the soluble copper salts are *potassium ferrocyanide*, which gives a mahogany-colored precipitate; *ammonia*, which gives a beautiful blue color, and *metallic iron* on which metallic copper will be deposited from a solution. The treatment of *acute poisoning* is albumen (milk or eggs); opium, demulcents and treating symptoms as they arise. In *chronic poisoning*, endeavor to eliminate it as if the poison were lead.

For what purposes is copper used in medicine?

It is used *internally* as an *astringent* in *chronic diarrhœa*, especially when ulceration exists, as an *emetic* (the sulphate), especially in narcotic poisoning, and occasionally as a *tonic* in nervous debility and functional nervous diseases. *Locally* it is efficacious as a *stimulant* to ulcers, as an *escharotic* in granular conjunctivitis and flabby granulations, and as an *astringent* injection in gonorrhœa.

What are the preparations of copper and their doses?

CUPRI SULPHAS (*copper sulphate*) called also *blue vitriol*, or *blue stone*. Dose as an *astringent* gr. $\frac{1}{4}$ -j (0.01-0.06), in pill; as an *emetic* gr. iiij-v (0.2-0.3); as an injection in gonorrhœa gr. $\frac{1}{4}$ -j (0.01-0.06) to f $\frac{3}{4}$ j (30.0).

ZINCUM—ZINC.

What are the sources, action, uses and preparations of zinc?

Zinc is obtained principally from the native carbonate and sulphide. The *effects* of the *zinc salts* are similar to those of copper, but are less powerful. In *excessive doses* they cause an *acute poisoning* very similar to that produced by copper, the *antidotes* being milk and eggs, alkalies and their carbonates. *Chronic poisoning* is so rare as to be disputed. The *test* for zinc in solution is the addition of *ammonium sulphide*, which precipitates *white zinc sulphide*, the only white sulphide met with.

They are used as *astringents* in diarrhœas; as *emetics*; as *nerve tonics* in epilepsy and chorea, and locally as *astringents* in conjunctivitis, gonorrhœa, etc.; as *sedatives* and *astringents* in eczema and erysipelas; as *dusting powders*, and as mild *stimulants* to ulcers.

ZINCI SULPHAS (*zinc sulphate—white vitriol*), dose as a *tonic* or *astringent* gr. ij–v (0.13–0.3); as an *emetic* (safer than copper sulphate) gr. x–xx (0.6–1.3). Locally, as a collyrium or as an injection in gonorrhœa, gr. j–ij (0.06–0.13) to fʒj (30.0).

ZINCI OXIDUM (*zinc oxide*) used internally in *catarrhal diarrhœa*, *membranous enteritis*, *epilepsy*, *chorea*, and occasionally to check *colliquative sweats*. Dose gr. j–v (0.06–0.3) in pill. It is used also as a *dusting powder*.

Oleatum zinci (*oleate of zinc*) contains 5 per cent. of zinc oxide, and is used externally.

The *ointment* (*unguentum zinci oxidi*) contains 20 per cent. of the salt and is much employed.

Impure or *commercial zinc oxide*, sometimes called "*tutty*," is used in pharmacy.

ZINCI ACETAS (*zinc acetate*) is used as a *collyrium* (gr. ij (0.13) to fʒj (30.0)) as an injection in the third stage of *gonorrhœa* (gr. j–xx (0.06–1.3) to fʒj (30.0)) and in *leucorrhœa* (ʒj to Oj—about 4.0 to 473.0).

ZINCI CARBONAS PRÆCIPITATUS (*precipitated zinc carbonate*) is used as a *dusting powder* and a mild *astringent ointment* (ʒj–ʒj—about 4.0 to 30.0).

ZINCI CHLORIDUM (*zinc chloride*) is a powerful *caustic* and *anti-septic*, employed to destroy lupus and indolent granulations. It is used also to preserve cadavers for dissection. *Burnett's disinfecting fluid* contains gr. 200 to fʒj (about 13.0 to 30.0).

Liquor zinci chloridi (solution of zinc chloride) may be used to disinfect water-closets and sinks.

ZINCI IODIDUM (zinc iodide) is a local stimulant and caustic.

ZINCI VALERIANAS (zinc valerianate) is employed as a nerve tonic in epilepsy and chorea; dose gr. j-ij (0.06-0.13) t. i. d.

ZINCI BROMIDUM (zinc bromide) is also official.

ZINCI PHOSPHIDUM (zinc phosphide) has been mentioned under phosphorus (*vid. Mineral Tonics*).

ARGENTUM—SILVER.

What are the physiological effects of silver?

Metallic silver is inert. Of its soluble salts, the oxide and nitrate are chiefly employed. *Locally* applied the nitrate coagulates albumen, is an astringent and superficial caustic, contracting the cutaneous vessels and producing a white stain on the skin which becomes black on exposure to light. Taken *internally* they are astringent, and after absorption exert a tonic influence upon the nervous system. In *large doses* they are gastro-intestinal irritants, often producing also violent convulsions, disturbed respiration, coma, dilated pupils, paralysis and death from respiratory failure. If *too long continued*, even in minute doses, *chronic argyria* results, consisting in emaciation, impaired memory and a peculiar indelible *blue discoloration*, first manifesting itself on the mucous membranes, and extending to the skin and internal organs.

What are the tests for silver, and how should argyria be treated?

The addition of *hydrochloric acid* or *sodium chloride* to a solution of silver produces a white precipitate of silver chloride, soluble in ammonia. *Acute argyria* is treated by giving immediately, as an *antidote*, a large amount of common salt, and treating symptoms p. r. n. In *chronic argyria* nothing does much good as a rule; potassium iodide may be tried as an eliminative agent.

What are the medicinal uses of the silver salts?

They are used as *astringents* in *chronic diarrhœas* and in *dysenteries* (internally or by enema) especially when ulceration is present, and are very useful in organic dyspepsias especially in *gastric ulcer* and in *chronic gastritis*. As a *nerve tonic* they are employed in

locomotor ataxia and other *spinal sclerosis*. *Externally* as a *stimulant* and *mild caustic* the nitrate is applied to slowly *granulating surfaces* or to unhealthy *ulcers*. As an *antiphlogistic* and *astringent* it is used in solution in various inflammations of the mucous membranes, as *conjunctivitis*, *pharyngitis*, *laryngitis*, etc., and may be used as an injection in the later stages of gonorrhœa (gr. j-ij (0.06-0.13) to f3j (30.0)) in the male, or in *acute vaginitis* (painting the inflamed vagina with strong solutions). It is employed also in superficial inflammations of the skin, in erysipelas, and to prevent pitting in smallpox.

What are the official silver salts and their doses ?

ARGENTI NITRAS (*silver nitrate*), dose gr. $\frac{1}{8}$ - $\frac{1}{4}$ (0.008-0.016) gradually increased to gr. ss-j (0.032-0.06). Great care should be taken to prevent staining the skin. For external use solutions are employed varying in strength from gr. j-3ij (0.06-8.0) to f3j (30.0) of distilled water.

Moulded silver nitrate or *lunar caustic* (*argenti nitras fusus*) is silver nitrate melted and poured into small moulds, for external use.

Diluted silver nitrate (*argenti nitras dilutus*) is 30 parts of silver nitrate and potassium nitrate 60 parts, moulded as above.

ARGENTI OXIDUM (*silver oxide*), dose gr. ss-j (0.03-0.06) in pill.

ARGENTI IODIDUM (*silver iodide*), dose as an alterative gr. j (0.06).

ARGENTI CYANIDUM (*silver cyanide*), used for preparing diluted hydrocyanic acid.

BISMUTHUM—BISMUTH.

What are the actions, uses and preparations of bismuth ?

Metallic bismuth is inert. The *less soluble salts* are not actively toxic when pure, as they are absorbed with difficulty. *Bismuth and ammonium citrate* is much more irritant, and may in large doses cause gastro-enteritis. In medicinal doses, they are astringent and sedative to the mucous membranes. A portion is absorbed and is eliminated by the kidneys chiefly, while that which is not absorbed is converted into a sulphide and colors the stools black. *Locally* applied they are antiseptic and sedative.

They are *used* internally as *sedatives* and *astringents* to check

nausea and vomiting in *gastric irritation*, and to relieve pain in *gastralgia*, *gastric ulcer* and *cancer*. In *acute diarrhœas* from irritation and in *chronic diarrhœas*, especially those occurring in children during the summer months, they are very valuable. *Externally* as mild astringents and antiseptics they are useful in the treatment of *ulcers*, and as an injection (suspended in mucilage) in *acute gonorrhœa* and in *vaginitis*. They are also used as dusting powders in *intertrigo* and as snuff in *acute coryza*.

BISMUTH SUBNITRAS (*bismuth subnitrate*) called also *pearl white* or *magistery of bismuth*, dose gr. v- ʒj (0.3-4.0).

BISMUTHI SUBCARBONAS (*bismuth subcarbonate*), dose gr. v-xxx (0.3-2.0).

BISMUTHI ET AMMONII CITRAS (*bismuth and ammonium citrate*), dose gr. j-v (0.06-0.3).

CERII OXALAS—CERIUM OXALATE.

What is cerium oxalate?

It is supposed to resemble the salts of bismuth in its effects, but has not been investigated. It is *used* with success in *reflex vomiting*, particularly the *vomiting of pregnancy*. Dose gr. j- ij (0.06-0.19), t. i. d. in pill or capsule.

ORDER V.—ANTACIDS.

What are antacids?

Antacids are remedies capable of neutralizing acids. The term, however, is here used in a more limited sense, and embraces only those medicines which are used in medicine to correct excessive acidity in the alimentary canal, as manifested by heartburn, acid eructations, gastric uneasiness, and diarrhœa of infants with "chopped spinach stools." Excessive acidity may be due to an excessive secretion of acid by the stomach, and here the antacids may be given after meals to neutralize the acid present, or the same effect may be produced by administering an acid before meals, which will decrease the acid secretion. On the other hand, it may be due to a deficient secretion of acid and consequent imperfect digestion and fermentation of the food. In these cases an antacid given before meals is useful by increasing the outward osmosis of acid, or a mixture of pepsin and acid after meals may supply the place of the deficient secretion. In acidity due to the development of the *sarcina*

ventriculi; sulphurous acid, the sulphites, carbolic acid or other antizymotics give the best results.

Name the antacids employed for the above purposes.

While all of the alkalies, alkaline earths and their carbonates are really antacids, yet in medical practice the sodium and calcium salts are chiefly used. *Solution of potassa* (q. v.) is sometimes, but not often, employed. The *spirit* and *aromatic spirit of ammonia* (q. v.) are frequently used, especially in sick headache or where a stimulating antacid is required.

SODIUM.

What are the effects and uses of the sodium salts?

Caustic soda and its solution are escharotics. The *sodium salts* taken on an empty stomach increase the secretion of acid gastric juice, but if given after a meal in sufficient quantity they neutralize the acid present. They diffuse readily into the blood and act somewhat as do the potassium salts, being, however, less irritant, more feebly diuretic, and possessing little if any poisonous influence over the (mammalian) heart, nerves or muscles.

The medicinal uses of the sodium preparations have been considered in discussing the general uses of the antacids.

What are the preparations of sodium?

LIQUOR SODÆ (*solution of soda*), dose mij-x (0.12-0.3) freely diluted.

SODII CARBONAS (*sodium carbonate*) is much used as an antacid in dyspepsia. In overdoses it is a corrosive poison. The antidotes are the vegetable acids. Dose gr. x-xxx (0.6-2.0). It is sometimes deprived of its water of crystallization and prescribed in pill form as the *dried carbonate* (*sodii carbonas exsiccatus*), in doses of gr. v-xv (0.3-1.0).

SODII BICARBONAS (*sodium bicarbonate*), is preferred to the carbonate because it is a more permanent salt and is less irritating to the stomach. Dose gr. x-xxx (0.6-2.0). It is an ingredient of *seidlitz powders* (q. v.) and other preparations. The *troches* (*trochisci sodii bicarbonatis*) each contain gr. iij (0.2) of the bicarbonate.

The other sodium salts are rarely used as antacids, and most of them have been sufficiently described previously.

CALCIUM.

What are the effects, uses and preparations of calcium?

Unslaked lime is a caustic. The *calcium salts* are antacid, slightly astringent and sedative to the mucous membranes, and are consequently much used in vomiting and in diarrhœa. The vapor arising from slaking lime is inhaled with advantage in membranous croup and in diphtheria. The preparations are LIQUOR CALCIS (*lime-solution—lime-water*), used as an antacid and gastric sedative in nausea and vomiting from irritability of the stomach, and often added to milk to increase its digestibility and render it more readily tolerated by the stomach. As an antacid and astringent it is useful in diarrhœa, especially in teething children. Locally it is used as an alkaline wash in various skin affections and as a dressing to burns, scalds, ulcers, etc., in the form of the *liniment* (*linimentum calcis*) often called *carron oil* (equal parts of lime-water and linseed oil).

SYRUPUS CALCIS (*syrup of lime*), a saturated, aqueous solution of calcium hydrate, the percentage of which varies with the temperature, but is about 0.17 at 59° F., is used as an astringent in diarrhœas and as an antidote to poisoning by carbolic acid. Dose f3ss-ij (2.0-8.0), freely diluted.

CALCI CARBONAS PRÆCIPITATUS (*precipitated calcium carbonate*) (dose gr. xx-3j (1.3-4.0)) possesses no especial advantage over *prepared chalk* (*creta præparata*), which is used in diarrhœas. Dose gr. x-xxx (0.6-2.0). They are used externally as dusting-powders in intertrigo. *Compound chalk powder* (*pulvis cretæ compositus*) contains 30 per cent. of prepared chalk. *Chalk mixture* (*mistura cretæ*) contains 20 per cent. of the compound powder in water and cinnamon water, and is much used, combined with astringents and opium, in diarrhœas, especially those of teething children. Dose f3ss-j (15.0-30.0). *Troches of chalk* (*trochisci cretæ*); each contains gr. jv (0.25) of prepared chalk.

Other preparations of calcium have been previously considered.

GROUP II.—TOPICAL MEDICINES.

Class I.—Antiseptics.

What are antiseptics?

Antiseptics are medicines which prevent fermentation and decomposition by destroying, or preventing the development of, the germs on which these processes depend. They are not always *germicides* (remedies which kill the germs of disease), for some of them hinder the growth and prevent the development of fresh germs, without destroying those already present; nor are they always *deodorants* (remedies which destroy disagreeable odors), for the microbes of disease may be destroyed, and foul chemical emanations continue, or the latter may exist without the presence of the former.

Name the chief antiseptics.

Antiseptics may be divided, according to their chemical composition, into *the inorganic or elementary*, and *the organic antiseptics* (or those belonging to the group of carbon compounds, including antiseptic oils and their camphors). To the first group belong hydrogen dioxide, potassium permanganate, chlorine water, chlorinated lime, Labarraque's solution, mercuric chloride, bromine, iodine, mercuric iodide, iodoform, iodol, sulphurous acid, sulphites, sulphides, boric acid, borates, and many of the mineral salts previously mentioned, as iron sulphate, lead nitrate, etc. In the second division are included carbolic acid and the carbonates, creosote, benzoic acid and the benzoates, naphthalin, naphthol, formaldehyd, the antipyretics (*q. v.*), the antiseptic oils (*v.* "*Carminatives*"), and thymol.

AQUA HYDROGENII DIOXIDI—SOLUTION OF HYDROGEN DIOXIDE.

What is solution of hydrogen dioxide?

It is a slightly acid, aqueous solution of hydrogen dioxide (also known as *hydrogen peroxide*), containing, when freshly prepared, about 3 per cent., by weight, of the pure dioxide, corresponding

to about 10 volumes of available oxygen. It occurs as a colorless liquid, without odor, but having a slightly acidulous taste.

It parts readily with its oxygen, and thus acts as an *oxidizing disinfectant*.

It is of course greatly inferior to mercuric chloride as a disinfectant, antiseptic, and germicide, but has the advantage of being non-poisonous. When brought in contact with pus, as on a suppurating surface, gas is evolved more or less abundantly, depending on the amount of pus present; a slight, superficial coagulation forms on the surface and the pus corpuscles are disintegrated.

It has been used as an *antiseptic injection* into *abscess cavities* after the evacuation of their contents, and into *sinuses*; and also as a local application to *superficial wounds* and *ulcers*, whether of common or specific origin. It is also used as an injection in *gonorrhœa*. In *scarlatinal* or *diphtheritic sore throat*, and in *tonsillitis*, it is often used as a gargle or spray.

It is a good *buccal disinfectant* and may advantageously be used for this purpose in *typhoid* and *other low fevers*, or in any case where the secretions tend to collect in the mouth and become offensive.

On account of the oxygen evolved it is an active *bleaching agent*, and is used in solutions of 10 to 15 per cent. strength to bleach black hair.

POTASSII PERMANGANAS—POTASSIUM PERMANGANATE.

What is potassium permanganate?

Potassium permanganate has been considered with the preparations of manganese (v. "*Mineral Tonics*"). Its disinfectant powers are great, but it is decomposed immediately on contact with organic matter, and hence is chiefly used in solution as a gargle in scarlatina and diphtheria, as a lotion to cancerous and other ulcers, to correct fœtor in otorrhœa, etc., and as an injection in leucorrhœa and in gonorrhœa. It is given internally as an emmenagogue (*q. v.*).

AQUA CHLORI—CHLORINE WATER.

What is chlorine-water?

It is an aqueous solution, containing at least 0.4 per cent. of *chlorine*. It is a powerful irritant and antiseptic. Albumin is the best *antidote*.

It is chiefly used, diluted, as a gargle in *diphtheria*, and as a wash for foul *ulcers* and *wounds*.

CALX CHLORATA—CHLORINATED LIME.

What is chlorinated lime?

It is composed of calcium hypochlorite and chloride, and when exposed to the air is decomposed, yielding about 35 per cent. of chlorine. It is an antiseptic, and is used to disinfect the air of sick rooms, urine, sputum and fecal discharges and privies.

LIQUOR SODÆ CHLORATÆ—SOLUTION OF CHLORINATED SODA.

What is this substance?

Solution of chlorinated soda, or *Labarraque's solution*, contains at least 2.6 per cent., by weight, of available chlorine, and may be used for the same purposes as the foregoing preparations.

HYDRARGYRI CHLORIDUM CORROSIVUM—MERCURIC CHLORIDE.

What is mercuric chloride?

This preparation has been considered with the mercurial salts (*v.* "*Alteratives*"). It is a valuable germicide, and is used in aqueous solutions of various strengths (1–2000 to 1–16,000) in all manner of surgical operations and affections where an antiseptic is required and can be employed. As several cases of poisoning have been reported, especially from its use in gynæcological and obstetrical practice, it should be employed with some caution.

BROMUM—BROMINE. IODUM—IODINE.

Are these substances antiseptic?

Bromine (*v.* "*Escharotics*") and *iodine* are powerful antiseptics, but are rarely used for this purpose, as their management requires great care and experience.

HYDRARGYRI IODIDUM RUBRUM—MERCURIC IODIDE.

Is mercuric iodide antiseptic?

Mercuric iodide is a powerful germicide, and has been employed as a substitute for corrosive sublimate, but, so far, has not superseded the latter.

IODOFORMUM—IODOFORM.

What is iodoform?

Iodoform is formyl teriodide, and occurs in minute yellowish crystals, with a disagreeable odor. It is insoluble in water, but freely soluble in ether and alcohol.

What are its effects and uses?

Locally, it is an antiseptic and local anæsthetic. It prevents the multiplication of germs, but will not destroy them if present before its use. It seems to be more readily absorbed by abraded than by mucous surfaces. *Internally*, it is tonic and alterative. *Moderate doses* stimulate the cardio-inhibitory centre, slowing and strengthening the cardiac beat and elevating the blood pressure, while *large doses* depress this centre, and finally paralyze the cardiac muscle. The *temperature* is generally reduced. It increases the salivary, biliary and intestinal secretions, and is eliminated chiefly by the kidneys (as sodium iodate), but to a slight extent by the lungs also. When excessive amounts are absorbed, either from internal or external sources, glomerulo-nephritis and fatty degenerations of various organs occur. Cases of poisoning may be divided into six classes, the danger increasing progressively from the first to the last: 1. High fever; 2. Fever, gastro-entero irritation, rapid pulse; 3. Very rapid, compressible pulse, no fever; 4. Very rapid pulse and high fever; 5. Great depression and collapse, rapidly fatal; 6. Symptoms like meningitis. Should poisoning occur, remove every particle of the dressing if it has been used locally, give opium and stimulants and treat symptoms. Iodoform has been *used* internally as a *cardiac tonic* in valvular disease with insufficient hypertrophy, and as an *alterative* in syphilitic rheumatism, neuralgiæ and the early stages of phthisis. In *diabetes mellitus* it has been advantageously given. *Externally*, it is employed as an anæsthetic and antiseptic dressing to irritable ulcers of all kinds, burns and wounds. In *etheral solution* it has been injected into the cavities of strumous abscesses, after the evacuation of their contents, to destroy the giant cells of tubercle and promote healthy granulation. In *suppositories* it is used in inflammation and ulceration of the vagina, uterus and rectum. *Bougies* containing iodoform are sometimes used in septic affections of the uterus and in gonorrhœa.

Dose gr. $\frac{1}{4}$ -v (0.01-0.3), in pill or capsule.

The *ointment* (*unguentum iodoformi*) contains 10 per cent. of iodoform.

IODOL.

What is iodol?

Iodol (not (official) is made by the action of iodine on pyrol, and contains 88.9 per cent. of the former. It is recommended as a substitute for iodoform, being tasteless, odorless, and, as it is more slowly absorbed, less likely to produce dangerous effects. It is especially recommended in nasal, pharyngeal and laryngeal cases, particularly in tubercular laryngitis, and may be used either in powder (by insufflation) or in solution with alcohol and glycerin. It has also been used internally in those cases to which iodoform is applicable. Dose gr. ss-ijj (0.03-0.19), in pill or capsule.

ACIDUM SULPHUROSUM—SULPHUROUS ACID.

What is sulphurous acid?

Sulphurous acid is an aqueous solution of about 6.4 per cent., by weight, of sulphurous acid gas.

What are its effects and uses?

It is a powerful deoxydizing agent, and deodorant. When sufficiently concentrated the gas is probably a germicide. When inhaled it is irritant, and, if in sufficient amount, causes inflammation and even acts as a caustic from the formation of sulphuric acid. It is eliminated by the urine and feces as sulphates. It is *used internally* (diluted) in *yeast-vomiting* to prevent the formation of *sarcina ventriculi* in *flatulency* depending on fermentation of the food, and *locally* as a wash in various parasitic skin diseases. In the form of burning sulphur it is employed to disinfect rooms. Dose $\mathfrak{m}\mathfrak{v}$ -f3j (0.3-4.0), freely diluted.

THE SULPHITES AND SULPHIDES.*

What are the effects and uses of these salts?

The *effects* of the *sulphites* depend upon the sulphurous acid which they contain, and which is readily evolved on the addition of almost

* See footnote, p. 71.

any acid. On exposure to the air they are converted into sulphates. Taken internally they are decomposed by the acid of the gastric juice, evolve sulphurous acid, are finally converted into sulphates, and thus absorbed and eliminated. They are *used* to fulfill the indications for sulphurous acid. The dose of *sodium sulphite* (*sodii sulphis*) is gr. v-xxx (0.3-2.0) t. i. d.; *sodium bisulphite* (*sodii bisulphis*) gr. v-xx (0.3-1.3); *sodium hyposulphite* (*sodii hyposulphis*) gr. v-xx (0.3-1.3).

The *sulphides* are said to possess the power of arresting the formation of pus in threatened boils and abscesses, and are used internally for this purpose.

Calx sulphurata (*sulphurated lime*) consists of a mixture containing at least 60 per cent. of calcium monosulphide together with unchanged calcium sulphate and carbon, in varying proportions, and is given in doses of $\frac{1}{10}$ - $\frac{1}{2}$ (0.006-0.03), repeated every $\frac{1}{2}$ -1 hour. *Sulphurated potassa* will be found among the sulphur preparations (v. "*Cathartics*").

ACIDUM BORICUM—BORIC ACID.

SODII BORAS—BORAX.

What are the effects and uses of boric acid and its salts?

Locally boric acid is antiseptic and deodorant. *Internally* they are somewhat diuretic, and retard the decomposition of the urine. Two fatal cases from the injection of large quantities into the pleural sac, and the cavity of a large lumbar abscess, have been reported, death being due to collapse and cardiac failure.

Boric acid is used as an antiseptic in powder or solution in many external inflammations and wounds, and in inflammations and aphthous or diphtheritic ulcerations of the mouth and fauces, the acid or its salts are of undoubted value.

In solution it has also been used to wash out *abscess cavities* and as an injection in *gonorrhœa*. As a *disinfectant* and *deodorant* in ophthalmic and aural practice it is very useful. Boric acid is recommended internally in *cystitis* depending on ammoniacal decomposition of the urine, and as an injection into the bladder in the cystitis of spinal diseases. Dose of the *acid* gr. v-x (0.3-0.6), t. i. d.; of the salt gr. xx-3j (1.2-4.0).

Glyceritum boroglycerini (*glycerite of boroglycerin*—*glycerite of glyceryl borate*—*solution of boroglycerite*) is made by dissolving 31 per cent. of boric acid in glycerin by means of heat, and is used as an external application for the disinfection of unhealthy ulcers or suppurating surfaces generally.

It is a useful application to burns, and has been applied with advantage to inflamed mucous membranes, as the pharynx, nasal passages, or conjunctiva.

ACIDUM CARBOLICUM—CARBOLIC ACID.

What is carbolic acid?

Carbolic acid, *phenol* or *phenic acid* is an oily liquid, or crystalline solid, which readily deliquesces, and is derived from coal-tar by fractional distillation.

What are the tests for carbolic acid?

The addition of *ferric sulphate* produces a lilac color; *bromine-water* causes a yellowish precipitate; it coagulates albumen, and precipitates nitro-cellulose from collodion.

What are its physiological actions?

Carbolic acid is a local anæsthetic and caustic (producing a white eschar) when used in concentrated solutions; it is antiseptic and germicide. *Internally*, it is a powerful gastro-intestinal irritant when taken undiluted. In diluted solutions it is an astringent.

It first stimulates and then exhausts the vagi (causing slowing, followed by acceleration of the heart's action, which is finally paralyzed), reduces the arterial pressure by paralyzing the vasomotor centres of the cord, increases and then depresses the respiratory movements, from its influence on the vagi and respiratory centre, and in large doses lowers the temperature slightly. In poisonous doses it causes vertigo, tremors, contracted pupils, stupor, coma, impaired motion, sensation and reflex action, and in the lower animals convulsions of spinal origin. It is eliminated chiefly by the kidneys and lungs, imparting a smoky color to the urine when the dose is excessive.

This is a sign of impending danger.

If *poisoning* occur from external applications, remove the dressings immediately; if the acid has been swallowed, give a saturated

solution of calcium carbonate. In either case administer atropine hypodermically, to counteract its respiratory and cardiac effects, and diluents freely, to aid its elimination.

What are its medicinal uses?

Carbolic acid was used externally as an antiseptic dressing and lotion in the various surgical affections and operations to which the mercuric chloride is applicable. It has also been used as a local anæsthetic previous to minor surgical operations, as the opening of small abscesses and felons, and in dilute solution to allay pruritus. It has been injected into the cavities of chronic abscesses and of the tunica vaginalis, to cause obliteration of their sacs after the withdrawal of the contained fluid.

Internally carbolic acid has been much used as a *gastric sedative* to check vomiting; as an *astringent* in diarrhœa; as an *antifermmentative*; as a *germicide* in *sarcina ventriculi*; and as an *antizymotic* in *typhoid fever*, *smallpox*, *scarlatina*, and *diphtheria*. It is also used by inhalation in *nasal catarrh*, *hay fever*, *bronchorrhœa*, *pertussis*, and *pulmonary gangrene*.

What are its preparations and doses?

ACIDUM CARBOLICUM (*carbolic acid*), dose gr. j-ij (0.06-0.13).

GLYCERITUM ACIDI CARBOLICI (*glycerite of carbolic acid*) contains 20 per cent. of the acid dissolved in glycerin. Dose ℥ ij-v (0.12-0.3).

ACIDUM CARBOLICUM CRUDUM (*crude carbolic acid*) may be used externally as a disinfectant. The *ointment* (*unguentum acidi carbolic*) contains 5 per cent. of the acid.

THE SULPHOCARBOLATES.

What are the sulphocarbolates?

Sodii sulphocarbolat (*sodium sulphocarbolate* or *sodium para-phenol sulphonate*)—the only one which is official—is used *internally* to fulfill the indications of carbolic acid, and as an *antifermmentative* in flatulency from fermentation of the food. It is also used *locally* as a sedative application to inflamed mucous membranes. Dose gr. x-xv (0.6-1.0) t. i. d.

CREOSOTUM—CREOSOTE.

What is creosote?

Creosote is a substance closely resembling carbolic acid, derived from wood-tar by dry distillation. It may be distinguished from carbolic acid by not coagulating collodion. It resembles carbolic acid in its effects and uses, and a solution of the latter is generally sold for it in the shops. The dose of pure creosote is $\mathfrak{m}\text{j}$ – ij (0.06–0.18).

Aqua creosoti (*creosote water*) is a 1 per cent. solution of creosote.

ACIDUM BENZOICUM—BENZOIC ACID.

What is benzoic acid?

Benzoic acid is obtained from benzoin by sublimation, and is an ingredient of many of the substances (*the balsams*) previously considered. *Locally* it is an irritant, antiseptic and germicide. *Internally* it increases the secretions generally, and is eliminated chiefly by the urine increasing the acidity of that fluid, hence its value in the phosphatic diathesis. It has been used *locally* as an antiseptic in the treatment of wounds and ulcers, but is not much employed for these purposes at present. It is also used as a stimulant and antiseptic application in the treatment of *chapped hands*, *fissured nipples*, etc., but for this purpose the preparations of benzoin are preferred. Benzoic acid or its salts are of great value in the treatment of *cystitis* from alkaline urine, *phosphatic deposits* in the urine, etc., for by acidulating the urine it renders the phosphates more soluble. They are also used internally as *antiseptics* in *diphtheria*, *erysipelas*, etc. Benzoic acid is added to unguents, to prevent rancidity. The dose is gr. v–xx (0.3–1.2); but for internal administration the salts are generally preferred.

Sodii benzoas (*sodium benzoate*) has been used as an *antipyretic* in *rheumatism*, and as an *antiseptic* in *pertussis*, *diphtheria*, and the *exanthemata*. Dose $\mathfrak{z}\text{j}$ – ij (4.0–8.0).

Ammonii benzoas (*ammonium benzoate*) is the best remedy for *incontinence* due to vesical irritation from alkaline decomposition of the urine. Dose gr. v–xx (0.3–1.2).

NAPHTALINUM—NAPHTALIN.

What is naphthalin?

Naphtalin, also called *naphthalene* or *naphthalin*, is a hydrocarbon obtained from coal-tar, occurring in colorless, transparent plates, and having a strong, peculiar odor, and a burning, somewhat aromatic taste. It is soluble in alcohol, but insoluble in water.

Naphtalin is *antiseptic* and *antiparasitic*. *Locally* it is mildly stimulant. When given internally in large doses or for a length of time it causes gastric irritation. Very little is absorbed when taken internally, the greater part being eliminated by the fæces and acting as an *intestinal antiseptic*. Some little, however, passes into the system and is eliminated by the kidneys and by the lungs. It is said that its continued administration to animals has resulted in cataract.

It is chiefly used as an *intestinal antiseptic* in typhoid fever, diarrhœas, and dysentery. It is also used as an *anthelmintic*, given internally for the removal of lumbrici, or used as an injection (gr. x-xx (0.6-1.2)) in sweet oil (f3ij (90.0)) to destroy the oxyuris. It has also been given in *pyelitis*, *cystitis*, etc., for its antiseptic effect on the urine, and has been employed in the treatment of *bronchorrhœa* and *fœtid bronchitis*. As a *local antiseptic* it is not much used. It has been used as a dressing for *wounds*, *ulcers*, and *chronic abscesses*. Dose gr. ij-x (0.12-0.64), administered in capsules.

NAPHTOL.

What is naphtol?

Naphtol, also called *isonaphthol* or *betanaphthol*, is a hydrocarbon prepared from naphthalene. It occurs as a white or yellowish-white powder, or in crystalline plates, with a faint phenol-like odor and a pungent taste. It is freely soluble in alcohol, but only slightly so in water.

In its action it resembles closely naphthalene. In animals large doses cause death by respiratory paralysis. No toxic symptoms have been reported in the human subject from its use.

Its *uses* are similar to those of naphthalin. It is a good *intestinal antiseptic*, preventing fermentation and consequent flatulence, and is used in gastric or intestinal fermentation, in dilated stomach, and in diarrhœas and dysentery. Dose gr. ij-v (0.12-0.32) although much larger doses have been given without deleterious effect. It is usually administered in capsules, which may be keratin-coated when

used as an intestinal antiseptic. It is often advantageously combined with bismuth subnitrate and sodium bicarbonate.

FORMALDEHYD.

What is this substance?

Formaldehyd, or *formic aldehyd* (not official), is an aqueous solution containing 40 per cent. of *formalin* or *formal*, obtained by oxidation of methyl alcohol. It forms a colorless solution having a pungent odor and taste. It has been much used as a deodorant and disinfectant and is said to be poisonous in a very moderate degree. Concentrated solutions applied to the skin are said (R. H. Cunningham) to cause a peculiar necrosis, unaccompanied by signs of inflammation. The gas is liberated by heating the solution, and it is said that an atmosphere containing 2.5 per cent. of the gas is speedily fatal to all micro-organisms.

It has been used to disinfect the surgeon's hands and instruments, in the preparation of aseptic catgut ligatures, and in dilute solutions (1 : 2000) in the treatment of *gonorrhœal affections of the female*. In the male it has sometimes effected a cure in *acute gonorrhœa*, more frequently when the disease was situated in the deeper parts of the urethra. Occasionally the irritation caused by its use was so great that the irrigation had to be discontinued. It has also been used in solutions of the same strength in *acute conjunctivitis*. Recently it has been recommended as a spray (1 per cent. solution, in the treatment of *pertussis*, and, used in the same manner, is strongly recommended in the treatment of *hay asthma*.

THE ANTIPYRETICS AND ANTISEPTIC OILS.

Are these substances used as antiseptics?

The *antipyretics* (*q. v.*) have been sufficiently considered.

The antiseptic properties of the *volatile oils* classed with the "*Carminatives*" (*q. v.*) have been alluded to under that head. Those of importance not hitherto mentioned are :

THYMOL.

What is thymol?

Thymol (*cymylic phenol*) is a crystalline substance found in the oil of thyme (*oleum thymi*) (*q. v.*). Locally it is a powerful antiseptic and

local anæsthetic. Given internally it lowers febrile temperature to some extent, and causes diaphoresis. In excessive amount it causes tinnitus aurium, deafness, diarrhœa, and sometimes delirium and collapse. It is used as an *antiseptic gargle* and *mouth-wash* in aphthous and diphtheritic ulcers of the mouth and fauces, and internally as a *gastric sedative* to allay vomiting. As an *anti-fermentative*, in flatulency from fermentation of the food and in intestinal indigestion, it is very useful. Dose gr. ss-ij (0.03-0.13). As much as gr. xx-xxx (1.2-2.0) have been given in 24 hours. It is administered in solution in glycerin, alcohol, and water.

Class II.—Irritants.

What are irritants ?

Irritants are medicines which are used locally to produce counter-irritation or inflammation of the parts to which they are applied. They are divided, according to the violence of their action, into Rubefacients, Vesicants, Suppurants, and Escharotics or Caustics.

ORDER I.—RUBEFACIENTS.

What are rubefacients ?

Rubefacients are medicines which are employed to produce a powerful but temporary irritation and congestion of the surface. They are useful in arousing the system in shock ; in stimulating the circulation of a part in case of frost-bite, or to aid in bringing on reaction in the cold stage of a congestive chill ; in all cases of congestion and in the formative stages of inflammations. The application of rubefacients for the relief of neuralgic and muscular pains is often followed by improvement which may even be permanent.

Mention the principal rubefacients.

Mustard ; capsicum ; oil of turpentine ; ammonia ; Burgundy pitch, and Canada pitch.

SINAPIS ALBA—WHITE MUSTARD.

SINAPIS NIGRA—BLACK MUSTARD.

What is mustard ?

Mustard is the seed of *Brassica alba*, or white mustard, and *B. nigra*, or black mustard (*Nat. Ord. Cruciferae*).

What chemical principles do they contain ?

Mustard seed contains a *fixed oil*. Black mustard seed yield on distillation a volatile oil (allyl sulphocyanide) which does not preëxist in the seed, but is formed by the action of water on *sinigrin* or *potassium myronate*. The white seed yield an *acrid fixed principle* which is formed by the action of water on *sinalbin*. These changes are probably due to the presence of *myrosin*, an albuminous ferment.

What are the effects and uses of mustard?

Locally, it is an irritant rapidly causing redness and a sensation of heat; if the application is long continued, inflammation and even gangrene may result. Internally it is a stomachic in small doses, an emetic in larger amounts, and a severe gastro-intestinal irritant if an excessive quantity be taken.

Mustard is *used* as a *rubefacient* in the form of a mustard plaster (made of the powdered seed known as mustard flour), whenever a mild counter-irritant effect is desired. The mustard should not be allowed to come in contact with the skin, but should be separated from it by a piece of gauze or a fold of muslin or newspaper. When a prolonged but gentle effect is desirable, the mustard may be added to a flaxseed poultice or diluted with flour. *Internally*, mustard-flour is used as an *emetic*, and is especially valuable when torpor of the stomach is present.

What are the preparations of mustard, and their doses?

The dose of powdered mustard as an emetic is ʒj-iv (4.0-16.0).

The official preparations are *mustard-papers* (*charta sinapis*), a convenient form for external use. Each square inch of paper contains about gr. vj of black mustard deprived of its oil. They should be dipped in warm water before applying. The *volatile oil* (*oleum sinapis volatile*) is very irritant, and is used in preparing the *compound liniment* (*linimentum sinapis compositum*).

CAPSICUM. OIL OF TURPENTINE. AMMONIA.

Are these substances used as rubefacients?

The uses of these substances as rubefacients will be found in detail under the respective drugs.

Linimentum ammoniac (*liniment of ammonia*) contains 3.5 per cent. of ammonia water in cotton-seed oil and alcohol.

PIX BURGUNDICA—BURGUNDY PITCH.

What is Burgundy pitch?

Burgundy pitch is the prepared *resinous exudation* from *Abies excelsa*, or Norway spruce (*Nat. Ord. Coniferae*). It contains a *resin* and a *volatile oil*, and is a mild rubefacient, used in the form of plaster in *subacute* and *chronic bronchitis*, to the back in *lumbago*, and to the joints in *chronic rheumatism*.

Pitch plaster with cantharides (*emplastrum picis cum cantharidē*) is more stimulating than the *plaster* (*emplastrum picis Burgundicæ*). *Iron and opium plasters* contain Burgundy pitch.

ORDER II.—VESICANTS.

What are vesicants?

Vesicants are remedies which, when applied to the skin, cause sufficient inflammation to produce an effusion of serum under the cuticle, or, in other words, to raise a blister. They are also called *epispastics*, or *blisters*. They are used in inflammations of the serous membranes, especially when accompanied by an outpouring of serum; to deplete the vessels in inflammations of the brain and its membranes; in chronic joint affections; in neuritis, and for the relief of various neuralgic pains, and are sometimes used in inflammations of the parenchymatous viscera. They are contraindicated in the very young or very old, where there is great depression of the vital powers, or where there is very high arterial tension.

Name the principal vesicants.

Spanish fly, potato fly, and stronger water of ammonia.

CANTHARIS—CANTHARIDES.

What is cantharides?

Cantharis vesicatoria or Spanish fly (*Class*, Insecta; *Order*, Coleoptera) is an insect found in the temperate parts of Europe, but especially in Spain and Italy. When dried they contain a *volatile oil* and a neutral crystalline principle, *cantharidin*, the vesicating principle.

What are its effects and uses?

Locally, cantharides is an intense irritant to any surface with which it comes in contact; thus, applied to the skin it causes inflammation of sufficient intensity to result in a serous effusion under the cuticle, and if the application be too long continued, gangrene may result. Constitutional effect may be caused from the local application of cantharides. *Internally, in small doses*, it is diuretic; in *larger amounts* it is an irritant, causing pain on urination, priapism, strangury, and even bloody urine. Sexual desire does not usually occur. In *excessive doses* it is a gastro-enteric irritant, causing

vomiting and purging, and even death, often preceded by dilated pupils, delirium, convulsions, and collapse. On *post-mortem examination* severe inflammation of the alimentary canal, kidneys, and bladder is found. The *treatment* consists in removing the application, or evacuating the contents of the stomach; the free use of diluents; opium (especially by the rectum); and avoiding all oily substances, as these aid the absorption of the poison.

What are its medicinal uses?

The internal uses of cantharides have previously been noticed (v. "*Emmenagogues*" and "*Diuretics*"). Externally it is used as a blistering agent, almost to the exclusion of other remedies. It must be remembered that poisoning may result from its local application.

What are its preparations and their doses?

The *cerate* (*ceratum cantharidis*) is used locally as a blistering agent, and is often known as *blistering cerate*. *Cantharidal collodion* (*collodium cantharidatum*) is applied with a brush, and is a convenient way of applying a blister to an irregular surface. The *tincture* (*tinctura cantharidis*) is used internally; dose gtt. v-x (0.3-0.6).

CANTHARIS VITTATA—POTATO FLY.

What is this substance?

The potato fly (not official) is an indigenous insect which contains *cantharidin*, and may be used as a substitute for the Spanish fly.

AQUA AMMONIÆ FORTIOR—STRONGER WATER OF AMMONIA.

Is this ever used as a blistering agent?

Stronger ammonia-water is a rapid but painful vesicant.

It forms the basis of *Granville's vesicating lotion* and of *Gondret's vesicating ointment*. It is but little used for this purpose.

ORDER III.—SUPPURANTS.

What are suppurants?

Suppurants are remedies which cause such powerful irritation of the surface to which they are applied as to produce a crop of pus-

tules. They are seldom employed at present except as an application to the throat and chest in *chronic laryngeal* or *bronchial affections*, or to the surface of a joint when *chronic inflammation* is present. The only medicines belonging to this order are *Croton oil* and *antimonial ointment*. These remedies have been previously described. Antimonial ointment is the more painful and its effects are more lasting. It is no longer official.

ORDER IV.—ESCHAROTICS.

What are escharotics?

Escharotics are medicines which are applied locally to destroy the structure and vitality of the tissues. The dead tissue excites inflammation, and is thrown off by ulcerative action. It is called the *sloUGH*, or *eschar*. Escharotics are *used* to destroy *morbid growths*, as lupus, epithelium, condylomata, warts, fungous granulations, etc.; to *destroy the virus* of malignant pustule, hospital gangrene, the bites of rabid animals or poisonous reptiles, the virus of chancre, etc.; to substitute a healthy for diseased action, and to stimulate indolent ulcers, sinuses, etc.

Mention the principal medicines used for these purposes?

Caustic potash; caustic soda; arsenious acid; bromine; zinc chloride; acid nitrate of mercury; corrosive sublimate; potassium bichromate; the mineral acids; silver nitrate; copper sulphate; and burnt alum.

POTASSA.

What is potassa?

Potassa is formed by the rapid evaporation of the solution of potassa, and is moulded into thin cylindrical sticks.

What are its effects and uses?

It is an exceedingly powerful caustic, destroying the surrounding tissues and penetrating to some depth below the surface. When it is used, it is well to protect the surrounding tissues from its action by adhesive plaster with a hole cut in its centre, through which the caustic is applied. It is used when a powerful effect is desired.

Potassa cum Calce (*potassa with lime*) consists of equal parts of potassa and quicklime, and is a powerful caustic. It is sometimes called *Vienna paste*.

SODA.

What is soda?

Caustic soda is made by the rapid evaporation of the solution of soda, and is a very powerful caustic, resembling potassa in its effects and uses. *London paste* is made by rubbing up equal parts of soda and lime.

ACIDUM ARSENOSUM—ARSENOUS ACID.

Is arsenous acid used as a caustic?

The uses of this substance as a caustic will be found under the article "Arsenic" (v. "Alteratives").

BROMUM—BROMINE.

What is bromine?

Bromine is a halogen element, and is a very powerful caustic. It is a volatile liquid, and has very great affinity for hydrogen. *Locally* it is a powerful deodorant and disinfectant, and a severe and deeply penetrating caustic. When *inhaled*, it is an intense irritant, producing laryngitis, bronchitis and even pneumonia. *Internally*, if not sufficiently diluted, it is a corrosive poison. It is *used* as a *caustic* in *hospital gangrene* after the removal of the slough. As it is heavier than water it may be poured into a wide-mouthed bottle half filled with water, which will prevent it from giving off irritating fumes. From this it may be sucked up with a syringe and injected into the wound which it is desired to cauterize. The fumes are sometimes cautiously inhaled for the cure of *acute coryza*, *chronic nasal catarrh*, *ozæna*, *hay asthma*, etc. For purposes of inhalation a solution of 3ss (2.0) in 3jv (120.0) of alcohol may be used.

ZINCI CHLORIDUM—ZINC CHLORIDE.

LIQUOR HYDRARGYRI NITRATIS—SOLUTION OF MERCURIC NITRATE.

HYDRARGYRI CHLORIDUM CORROSIVUM—MERCURIC CHLORIDE.

Are these medicines escharotics?

The uses of these substances as caustics will be found under Zinc (v. "Astringents") and Mercury (v. "Alteratives").

POTASSII BICHROMAS—POTASSIUM BICHROMATE.

What is potassium bichromate?

It is an irritant *internally*, and in overdoses a corrosive poison. *Externally* it is a *caustic*, used to destroy syphilitic condylomata, warts, etc. *Internally* it is used as an *alterative* in doses of gr. $\frac{1}{5}$ (0.013) t. i. d.

THE MINERAL ACIDS.

Are the mineral acids used as caustics?

The mineral acids are powerful caustics, and are often employed to remove warts, or other excrescences, to cauterize chancres, indolent sinuses and ulcers. Nitric acid is preferred as an escharotic.

SILVER NITRATE, COPPER SULPHATE, and ALUM have all been previously spoken of as caustics under their appropriate heads.

Class III.—Demulcents.

What are demulcents?

Demulcents are bland remedies which, when applied to inflamed or irritated surfaces, exert a soothing influence upon them and are employed internally for this purpose. They are *used* in acute inflammations of the alimentary canal; in pharyngeal, laryngeal and bronchial irritation; in renal, cystic, and urinary affections generally; and to suspend medicinal substances which are insoluble in water. They are also used as light articles of diet.

Describe the demulcents.

AQUA (*water*) is used in medicine in its purest attainable state, *i. e.*, rain or melted snow. For pharmaceutical purposes it should only be used in the form of *distilled water* (*aqua destillata*). A certain amount of water is necessary for the proper performance of the functions of the body. It is an *active diuretic*, increasing not only the watery, but also the solid ingredients of the urine. It is *eliminated* by the kidneys, skin, lungs and bowels. It is *used* as a *diuretic* in acute renal diseases, and is also often serviceable in chronic Bright's disease. To dilute a highly concentrated urine in *ardor urinæ*, especially when there is a tendency to the formation of gravel or calculi, it is invaluable.

It is useful in fevers, as it aids the elimination of the waste products. *Externally*, it is used in the form of baths, either hot or cold, not only for purposes of cleanliness, but also as a therapeutic agent. Impregnated with carbonic acid gas under pressure, it makes an agreeable drink known as *soda-water*, which is often useful in allaying nausea and vomiting.

ACACIA, or *gum arabic*, is a *gummy exudation* from Acacia Senegal (*Nat. Ord.* Leguminosæ), a tree of Africa. It contains *arabic acid*, and is much *used* in solution as a demulcent in various forms of *sore throat*. It is also used in making emulsions and pills.

Its preparations are: a *mucilage* (*mucilago acaciæ*) and a *syrup* (*syrupus acaciæ*). *Emulsion of almonds* (*emulsum amygdalæ*) is an agreeable form for administration.

TRAGACANTHA (*tragacanth*) is a *gummy exudation* from *Astragalus gummifer* and other species of *Astragalus* (*Nat. Ord. Leguminosæ*), and contains *arabin* and *tragacanthin*. It is only used internally as an agent to suspend heavy powders or to give consistence to lozenges. A *mucilage* (*mucilago tragacanthæ*) is official.

ULMUS (*elm*) is the *inner bark* of the *Ulmus fulva*, or slippery elm (*Nat. Ord. Urticacæ*), and contains a large amount of mucilaginous matter, and though more often used *externally* as an emollient poultice, is sometimes employed as a *demulcent* in inflammations of the bowels, and in laryngitis. A *mucilage* (*mucilago ulmi*) is official.

ALTHÆA, the root of *A. officinalis*, or marsh-mallow (*Nat. Ord. Malvaceæ*), is a pleasant demulcent. A *syrup* (*syrupus althææ*) is official.

LINUM (*flaxseed*) is the *seed* of *Linum usitatissimum*, or flax (*Nat. Ord. Linææ*), and contains *fixed oil*, *mucilaginous matters*, *vegetable albumen*, etc. The *oil* (*oleum lini*) is a laxative, but is chiefly used externally in emollient liniments. The seed may be used internally as an infusion, flavored with lemon and sugar (*flaxseed tea*), in bronchial catarrhs, enteric inflammations, dysentery, and renal, vesical, or urethral irritations. The ground seed are much used to make emollient poultices.

SASSAFRAS MEDULLA (*sassafras pith*) is the pith of the stems of *S. variifolium* (*Nat. Ord. Laurinææ*). The *mucilage* (*mucilago sassafras medullæ*) may be used internally as a demulcent, but is chiefly employed locally in inflammations of the eye.

GLYCYRRHIZA, commonly called *liquorice-root*, is the root of *G. glabra* (*Nat. Ord. Leguminosæ*). It contains a *glucoside* (*glycyrrhizin*), *asparagin*, *resin*, etc. It is chiefly used as a demulcent in catarrhal affections of the respiratory organs and to disguise the taste of quinine, of the ammonium salts, and of other disagreeable medicines; occasionally, also, in kidney and bowel complaints. The preparations are an *extract* (*extractum glycyrrhizæ*), often called *liquorice*; *refined extract* (*extractum glycyrrhizæ purum*); *fluid extract* (*extractum glycyrrhizæ fluidum*), used as an addendum to

cough medicines; *ammoniated glycyrrhizin* (*glycyrrhizinum ammoniatum*) dose gr. v-x (0.3-0.6); *compound glycyrrhiza mixture*, *brown mixture* (*mistura glycyrrhizæ composita*) contains the refined extract, mucilage of acacia, syrup, paregoric (12 per cent.), anti-monial wine (6 per cent.), sweet spirits of nitre and water; dose f3ss (15.0). *Compound liquorice powders* (containing sulphur and senna, —used as a laxative), have been previously mentioned.

CETRARIA.—*Cetraria islandica*, or Iceland moss (*Nat. Ord. Lichenes*), is a lichen growing in the northern portions of both continents. It contains *lichenin* and a bitter principle, *cetrarin*, and is used as a demulcent. A *decoction* (*decoctum cetrariæ*) is official.

CHONDRUS.—*Chondrus crispus*, Irish Moss, or Carrageen (*Nat. Ord. Algæ*), is a marine plant found on the coast of Ireland and New England. It contains *carrageenin*, etc., and is used as a nutritive (in the form of a blance-mange), as a demulcent, and locally as an emollient poultice.

AMYLUM (*starch*).—This is the *fecula* of the *seed* of *Zea mays* (*Nat. Ord. Graminaceæ*). With iodine its solution forms a rich blue iodide. It is converted into glucose in the alimentary canal, and thus absorbed. It is used as an article of diet and locally as a dusting powder. Its solution is employed as a vehicle for laudanum enemata.

Glyceritum amyli (*glycerite of starch* or *plasmine*) is used as a vehicle for astringent applications to the eye, and as an unguent in scarlatina and smallpox.

LYCOPODIUM, the *sporules* of *L. clavatum*, or club-moss and other species of lycopodium (*Nat. Ord. Lycopodiaceæ*), is used as a dusting powder for intertrigo, and is very useful in children with excoriated nates.

ICHTHYOCOLLA (*isinglass*) is prepared from the swimming bladder of *Acipenser Huso* (the sturgeon) and other species. (*Class, Pisces; Ord. Sturiones*). It is the purest form of gelatine, and is chiefly used locally as a *plaster* (*emplastrum ichthyocollæ*—*court-plaster*). Gelatine is also used as an article of diet, and in the form of capsules for administering other medicines.

Class IV.—Emollients.

What are emollients?

Emollients are bland, fatty substances used externally to soften the skin and render it more pliable or to serve as a vehicle for other medicaments.

Describe the emollients.

ADEPS (*lard*), the prepared internal fat of the abdomen of *Sus Scrofa* (the hog), consists of olein and stearine, and is used in the preparation of unguents. *Ointment (unguentum)* consists of lard (80 per cent.) melted with yellow wax. *Cerate (ceratum)* contains 70 per cent. of lard and 30 per cent. of white wax. *Adeps (benzoinated lard)* contains 2 per cent. of benzoin to prevent rancidity.

ADEPS LANÆ HYDROSUS (*hydrous wool-fat—lanolin*) is the purified fat of the wool of sheep (*Ovis Aries*) mixed with not more than 30 per cent. of water.

SEVUM (*suet*), the internal fat of the abdomen of *Ovis Aries* (the sheep), consists, when purified, almost entirely of stearine.

CETACEUM (*spermaceti*), a peculiar, concrete, fatty substance obtained from *Physeter macrocephalus* or sperm-whale. A *cerate (ceratum cetacei)* is official, and it also enters into the preparation of *cold cream*.

CERA FLAVA (*yellow*) and CERA ALBA (*white wax*). The former is a peculiar, concrete substance, prepared by *Apis mellifica*, or honey bee. The latter is the former which has been bleached. They are used in making cerates, plasters, and ointments.

ACIDUM OLEICUM (*oleic acid*), used chiefly in preparing the oleates of mercury and veratrine.

GLYCERINUM (*glycerin*) is the hydrate of glyceril and exists in animal oils combined with fatty acids. Internally it is nutrient, demulcent, and, in large doses, laxative. Administered by the

bowel it is laxative even in small amounts, for which purpose the *suppositories* (*suppositoria glycerini*) are much used. Locally, it is much used as an emollient. *Glyceritum Vitelli* (*glycerite of yolk of egg*) is used in making emulsions. *Glycerite of starch* (v. "*Amylum*") is used externally.

The *glycerites of carbolic acid, of tannic acid, of boroglycerin, and of hydrastis* are also official.

OLEUM THEOBROMATIS (*oil of theobroma*) or cacao-butter, is the *fixed oil expressed from the seed* of *Theobroma Cacao* (*Nat. Ord. Sterculiaceæ*), and is used as an unguent and to give form and consistence to suppositories.

OLEUM GOSSYPH SEMINIS (*cottonseed oil*), a *fixed oil expressed from the seed* of *Gossypium herbaceum* (*q. v.*), is used as a substitute for almond or olive oil, and enters into the composition of numerous liniments.

OLEUM SESAMI (*oil of sesamum—oil of benne*) is a *fixed oil expressed from the seed* of *Sesamum indicum* (*Nat. Ord. Pedaliaceæ*). It is used for the same purposes as cottonseed oil.

PETROLATUM MOLLE (*soft petrolatum*) is a mixture of hydrocarbons, chiefly of the marsh-gas series, obtained by distilling off the lighter and more volatile portions from petroleum and purifying the residue.

It differs from *Petrolatum Liquidum* (*liquid petrolatum*) and from *Petrolatum Spissum* (*hard petrolatum*) chiefly in its consistence, in its melting-point, and in its specific gravity. The latter is of the consistence of cerate, and the former that of an oil, while soft petrolatum more nearly approaches that of an ointment. Under the names of vaseline and cosmoline, the heavier petroleum oils are much used externally as emollients and as a basis for ointments, and internally to allay irritation of the mucous membranes.

Many of the "*Demulcents*" also are employed in the form of poultices as emollients.

Class V.—Protectives.

What are protectives?

Protectives are substances used to exclude light or air from the surface to which they are applied, or otherwise to protect it from injurious influences. Many *surgical appliances* are thus “protectives,” but do not belong to the domain of “Materia Medica.” The *plasters* and *unguents*, when simply used to protect the surface, come under this class. The following agents have not been previously considered: Collodion, solution of gutta-percha and solution of sodium silicate.

What are these substances?

CALCI SULPHAS EXSICCATUS (*dried gypsum, dried calcium sulphate or plaster of Paris*) is a powder containing about 95 per cent., by weight, of calcium sulphate.

It is used as a permanent dressing in the treatment of fractures, caries of the vertebræ, etc., where it is necessary to give continual support to the parts for some time. For this purpose it is moistened and thoroughly rubbed into successive layers of previously moistened roller bandage. It *sets* very rapidly, forming with the bandage a stiff, board-like support.

COLLODIUM (*collodion*) is a solution of *pyroxylin* or soluble gun-cotton (*pyroxylinum*) (4 per cent.), in stronger ether and alcohol. When applied to the skin, the solvents rapidly evaporate, leaving a colorless, transparent, flexible and strongly contractile film, closely adherent, difficult to detach, and impervious alike to air and water. It has been used in surgery as an antiphlogistic in inflammation, on account of its contractile power, and also applied on gauze, as a substitute for adhesive plaster in the treatment of wounds. Its contractile power is an objection to its use as a simple protective, for which purpose it is better to employ *flexible collodion* (*collodium flexile*), which contains Canada turpentine (5 per cent.) and castor oil (3 per cent.), and is more pliable and elastic. *Styptic collodion* (*collodium stypticum*) contains, in addition to collodion, tannic acid (20 per cent.), ether (25 per cent.), and alcohol (5 per cent.).

ELASTICA (*India rubber or caoutchouc*) is the prepared *milk-juice* of various species of *Hevea* (*Nat. Ord.* Euphorbiaceæ), known in commerce as *Para Rubber*. It is used in making plasters, bougies, etc., and has been used dissolved in commercial chloroform as a protective to small abrasions, chaps, threatened bedsores, etc.

LIQUOR SODII SILICATIS (*solution of sodium silicate*), commonly called *solution of soluble glass*, is principally used in the treatment of fractures, applied on successive layers of roller bandage as a permanent dressing.

Class VI.—Coloring Agents.


What are coloring agents?

They are employed almost exclusively to give their particular color to tinctures, spirits and ointments, and are—

COCCUS (*cochineal*), the *female dried insect*. *Coccus cacti*, a native of Mexico and Central America, containing *carminic acid*, to which it owes its peculiar red color. It has been used also as an *antispasmodic*; dose gr. $\frac{1}{3}$ (0.02), t. i. d.

CROCUS (*saffron*), the stigmas of *Crocus sativus* (*Nat. Ord.* Irideæ), contains *safranin*. A *tincture* (*tinctura croci*) is official.

SANTALUM RUBRUM (*red sanders*), the *wood* of *Pterocarpus santalinus* (*Nat. Ord.* Leguminosæ), contains *sandal*, *santalic acid*, and *pterocarpin*, and is used to give a red color to pharmaceutical preparations.



INDEX.

- ABSINTHIUM, 40
 Acacia, 263
 mucilage of, 263
 syrup of, 263
 Aceta, 22
 Acetanilid, 80, 81
 Acetanilidum, 80, 81
 Acetic acid, 168
 diluted, 168
 glacial, 168
 Acetphenctidin, 81
 Acetum opii, 94
 scillæ, 208
 Acid nitrate of mercury, 63
 Acids, 168, 169
 Acidum aceticum, 168
 dilutum, 168
 glaciale, 168
 arsenosum, 69, 261
 benzoicum, 252
 boricum, 249, 250
 carbolicum, 250, 251
 citricum, 168, 169
 gallicum, 230
 hydrobromicum dilutum, 99
 hydrochloricum, 58
 hydrocyanicum dilutum, 139, 141
 lacticum, 59
 nitricum, 57
 nitrohydrochloricum, 58
 oleicum, 266
 phosphoricum, 58
 sulphuricum, 57
 sulphurosum, 248
 tannicum, 229
 tartaricum, 169
 Aconine, 162
 Aconite, 162-164, 218
 preparations and doses of, 163, 164
 Aconitine, 162-164
 Aconitum, 162, 164
 Adeps, 265
 benzoinatus, 226
 lanæ hydrosus, 266
 Administration of medicines, 32
 Adonidin, 156, 157
 Æther, 116-119, 149
 fortior, U. S. P. 1880, 119
 preparations of, 119
 Alcohol, 146-149
 absolutum, 148
 official preparations, 148, 149
 Alcoholic extract of belladonna
 leaves, 110
 Alkaloids, cinchona, 42, 46
 Allium, 225
 Allspice, 184
 Allyl, sulphocyanide, 256
 Almonds, preparations of, 142
 Aloe purificata, 192
 Aloes, 192
 pills of asafœtida and, 128
 Aloin, 192
 Aloinum, 192
 Alteratives, 59-77
 Althea, 264
 Alum, 181, 236, 262
 Alumen, 181, 236
 Alumin sulphas, 236
 Aluminum sulphate, 236
 Amidophenacetin, 81, 82
 Ammonia, 151, 242, 257
 gas, 149
 liniment, 257
 stronger water of, 259
 Ammoniac, 222, 226
 Ammoniacum, 226
 Ammoniated glycyrrhizin, 265
 mercury, 63
 tincture of guaiac, 75
 Ammonii benzoas, 252
 bromidum, 99
 carbonas, 151
 chloridum, 72, 73, 223
 iodidum, 66
 valerianas, 127
 Ammonium, 149, 151
 acetate, solution of, 219
 benzoate, 252
 carbonate, 151
 chloride, 72, 73, 222, 223
 iodide, 66
 muriate, 223
 valerianate, 127
 Amulsum amygdalæ, 142
 Amyl nitris, 138, 151
 nitrite, 138, 151
 Amylum, 265, 267
 Anæsthetics, 116, 123
 Anemonic acid, 167
 Anemonii, 167
 Anise, 187
 Anisum, 187
 Anthemis, 39, 41

- Antifebrin, 80, 81
 Antimonial ointment, 269
 powder, 161
 Antimonii et potassii tartras, 161, 181
 oxidum, 161
 sulphidum, 162
 sulphidum purificatum, 162
 sulphuratum, 162
 Antimonium, 159, 162
 and potassium tartrate, 161, 181
 Antimonious oxide, 161
 sulphide, 162
 Antimony, 159, 162
 Antipyretics, 77, 89
 Antipyrine, 78, 80, 116
 Antispasmodics, 126, 130
 Apiol, 177
 Apocynum, 210
 Apomorphine, 222
 hydrochlorate, 179, 180
 Apothecaries' measure, 24
 relation of, to metrical system, 25
 weight, 23
 relation of, to metrical system, 25
 Approximate measurements, 26
 Aqua, 262
 ammonia, 151
 fortior, 151, 259
 amygdalæ, 144
 anisi, 187
 aurantii florum fortior, 187
 camphoræ, 129
 chloroformi, 121
 cinnamomi, 184
 creosoti, 252
 destillata, 263
 feniculi, 187
 hydrogenii dioxidi, 244, 245
 menthæ piperitæ, 186
 viridis, 186
 pagliarii, 236
 rosæ, 232
 Aquæ, 21
 Arbutin, 212
 Argenti cyanidum, 141, 240
 iodidum, 240
 nitras, 240
 dilatatus, 240
 fusus, 240
 oxidum, 240
 Argentum, 239, 240
 Arnica, effects and uses of, 167
 flowers, tincture of, 167
 fluid extract of, 167
 plaster of, 167
 root, extract of, 167
 tincture of, 167
 Arnicin, 167
 Aromatic bitters, 39-41
 preparations and doses of, 41
 elixir, 188
 powder, 185
 spirit of ammonia, 151
 sulphuric acid, 57
 Arsenic, 67-70
 Arsenic iodide, 69
 preparations and doses of, 69
 symptoms and treatment of poison-
 ing by, 68, 69
 Arseni iodidum, 69
 Arsenous acid, 69, 261
 Arsenum, 69-70
 Asafetida, 127, 128
 emulsion of, 128
 pills of, 128
 pills of aloes, etc., 128
 preparations and doses of, 128
 tincture of, 128
 Asclepias, 222, 224
 fluid extract of, 224
 Aspidium, 200
 Aspidosperma, 224
 fluid extract of, 224
 Atropinæ, 152
 Atropine, 107-110, 152, 220
 sulphas, 110
 treatment of poisoning by, 108, 109
 Aurantii amari cortex, 187
 dulcis cortex, 187
 Auri et sodii chloridum, 64
 Aurum, 64

 BALSAM OF FIR, 214
 Peru, 227
 tolu, 222, 227
 Balsamum peruvianum, 227
 tolutanum, 222, 227
 Bark of cotton root, 173, 174
 Basham's mixture, 50
 Belladonna, 107, 110
 liniment of, 110
 ointment of, 110
 plaster of, 110
 root, fluid extract of, 110
 tincture of, 110
 treatment of poisoning by, 108, 109
 Benne, oil of, 267
 Benzoic acid, 226, 252
 Benzoin, 222, 226
 compound tincture of, 226
 tincture of, 226
 Benzoinated lard, 226, 266
 Benzoinum, 226
 Berberine, 37, 39, 46
 Beta-naphtol, 243
 Bismuth, 240, 244
 and ammonium citrate, 241
 subcarbonate, 241
 subnitrate, 241
 Bismuthi et ammonii citras, 241
 subcarbonas, 241
 subnitras, 241
 Bismuthum, 240, 241
 Bitter almond, water of, 141
 essence of, 142
 spirits of, 141, 142
 tonics, 37
 wine of iron, 51
 Bitters, 37-39
 aromatic, 39-41

- Bitters, peculiar, 42-47
 simple, 37-39
 Black draft, 193
 drop, 94
 Blaud's pills, 49
 Blue gum tree, 39
 mass, 61, 62, 193
 ointment, 62
 stone, 237
 vitriol, 237
 Borax, 249, 250
 Boric acid, 249, 250
 Brandy, 148
 Brayera, 201
 Bromide of ammonium, 99
 calcium, 99
 potassium, 99
 sodium, 99
 strontium, 99
 zinc, 99
 Bromides, 98, 99, 136
 Bromine, 246
 Bromoform, 123
 Bromum, 246
 Brown mixture, 265
 Brucine, 131
 Bryonia, 196
 Bryony, 196
 Buchu, 211
 fluid extract of, 211
 Burdock, 77
 Burgundy, 257
 Burnett's disinfecting fluid, 238
 Butyl chloral hydrate, 97

 CAFFEINA, 155, 156
 citrate, 156
 effervescens, 156
 Caffeine, 155, 156, 208
 administration of, 156
 citrate, 156
 effervescent, 156
 Cajuput, oil of, 185
 Calamus, 185
 fluid extract of, 185
 Calci bromidum, 99
 carbonas præcipitatus, 243
 chloridum, 72
 hypophosphis, 72
 phosphas præcipitatus, 72
 sulphas exsiccatus, 268
 Calcined magnesia, 194
 Calcium, 243
 chloride, 72
 hypophosphite, 72
 phosphate, 71, 72
 sulphate, dried, 268
 Calendula, 76
 tincture of, 76
 Calomel, 62, 139, 211
 Calumba, 37, 38, 39
 Calumbo, 38, 39
 Calx chlorata, 246
 sulphurata, 249
 Cambogia, 197

 Campeachy wood, 232
 Camphor, 128, 129
 cerate, 129
 liniment, 129
 monobromated, 129
 preparations and uses of, 129
 spirit of, 129
 water of, 129
 Camphora, 128, 129
 monobromata, 129
 Camphorated tincture of opium, 94
 Canada balsam, 214
 Canadensis, 46
 Cannabis Indica, 104, 105
 extract of, 105
 fluid extract of, 105
 tincture of, 105
 Cantharidal collodion, 239
 Cantharides, 177, 217, 258, 259
 tincture of, 217, 259
 Cantharis, 217, 258, 259
 vesicatoria, 258
 vittata, 259
 Caoutchouc, 269
 Capsicum, 183, 257
 fluid extract of, 183
 oleoresin of, 183
 plaster of, 183
 tincture of, 183
 Capsules, 23
 Caraway, 187
 Carbolic acid, 250, 251
 crude, 251
 glycerite of, 251
 ointment of, 251
 Cardamom, 185
 compound tincture of, 185
 tincture of, 185
 Cardamomum, 185
 Carica papaya, 35
 Carrageen, 265
 Carron oil, 243
 Carum, 187
 Caryophyllus, 184
 Cascara sagrada, 189
 Cascarilla, 39
 Cassia cinnamon, 183, 184
 fistula, 189
 Castanæ, 233
 fluid extract of, 233
 Castor oil, 189, 190, 191
 Catechu, 231
 compound tincture of, 231
 troches of, 231
 Caustic soda, 242, 261
 Celandine, 197
 Centimeter, 24
 Cera alba, 266
 Cera flava, 266
 Cerata, 21
 Cerates, 20, 21
 Ceratum camphoræ, 129
 cantharidis, 259
 cetacei, 266
 plumbi subacetatis, 235

- Cerii oxalas, 241
 Cerium oxalate, 241
 Cetaceum, 266
 Cetraria, 265
 islandica, 265
 Chalk, 243
 compound powder, 243
 mixture, 243
 prepared, 243
 troches of, 243
 Chamomile, 39
 German, 39
 Charta potassii nitratis, 205
 Charta sinapis, 257
 Chartæ, 20
 Chelidonium, 197
 Chenopodium, 200
 oil of, 200
 Chian turpentine, 214
 Chimaphila, 212
 fluid extract of, 212
 Chinoline, 89
 Chirata, 37, 39
 Chiratae, 39
 Chittam bark, 189
 Chloral, 95-97, 136
 treatment of poisoning by, 96
 Chloralamide, 97
 Chloralose, 97, 98
 Chlorinated lime, 246
 soda, solution of, 247
 Chlorine-water, 245, 246
 Chloroform, 119-121
 emulsion of, 121
 liniment of, 121
 preparations of, 121
 spirit of, 121
 water, 121
 Chloroformum, 119-121.
 Choline, 106
 Chondrus, 265
 Cimicifuga, 158, 159
 extract of, 159
 fluid extract of, 158
 tincture of, 159
 Cinchona, 42-46
 administration of, 46
 alkaloids, 42-46
 calisaya, 42
 chemical constituents of, 427
 condaminea, 42
 infusion of, 45
 micrantha, 42
 pallida, 42
 succirubra, 42
 Cinchonine, 43
 Cinchonidinæ sulphas, 45
 Cinchonidine, 42
 sulphate, 45
 Cinchonæ sulphas, 45
 Cinchonine, 42
 sulphate, 45
 Cinnamomum cassia, 183, 184
 saigoncum, 183
 zeylanicum, 184
 Cinnamon, 183, 184
 oil of, 184
 spirit of, 184
 tincture of, 184
 water of, 184
 Citric acid, 168, 169
 syrup of, 169
 Citrine ointment, 63
 Classification of medicines, 33
 Cloves, 184
 oil of, 184
 Coca, 113-115
 preparations and doses of, 115
 Cocainæ hydrochloras, 115
 Cocaine, 114, 124, 125
 hydrochlorate, 115
 treatment of poisoning by, 114, 115
 Coccus, 269
 Cochineal, 269
 Codeine, 95
 Cod-liver oil, 70, 71
 dose and administration, 71
 Colchicia, 73
 Colchicine, 73, 74
 Colchicum, 74
 fluid extract of, 74
 preparations and doses of, 74
 root, extract of, 74
 fluid extract of, 74
 Cold cream, 232
 Collodion, flexible, 268
 styptic, 268
 Collodium cantharidatum, 259
 flexile, 268
 stypticum, 268
 Colocynth, 197
 Colocynthis, 197
 Compound cathartic pills, 198
 decoction of sarsaparilla, 75
 effervescent powder, 195
 fluid extract of sarsaparilla, 75
 glycyrrhiza mixture, 265
 liquorice powders, 265
 pills of antimony, 162
 powder of morphine, 94
 solution of iodine, 66
 syrup of sarsaparilla, 75
 squill, 161
 Confectio rosæ, 232
 sennæ, 193
 Confectiones, 20
 Confections, 20
 Conium, 143-144
 extract of, 143
 fluid extract of, 144
 preparations and doses of, 143, 144
 Conserves, 20
 Convallamarin, 159
 Convallaria, 159
 fluid extract of, 159
 Convallarin, 159
 Conydrine, 143
 Copaiba, 216
 massa, 216
 oil of, 216

- Copper, 237
 sulphas, 181
 sulphate, 237, 262
 Copperas, 49
 Coriander, 187
 Coriandrum, 187
 Cornutin, 171
 Corrosive sublimate, 60, 61, 211
 treatment of poisoning by, 61
 Cotton bark, 174
 fluid extract of, 174
 root, 173, 174
 Cottonseed oil, 267
 Court-plaster, 265
 Cowlings rule for dose in children, 31
 Coxe's hive-syrup, 161, 208
 Cream of tartar, 195, 205
 Cremor tartari, 195
 Creosote, 252
 water, 252
 Creosotum, 252
 Cretæ præparata, 243
 Crocus, 269
 tincture of, 269
 Croton chloral hydrate, 97
 eluteria, 39
 oil, 198, 260
 Cubebæ, 216
 Cubeba, 216
 fluid extract of, 217
 oil of, 217
 oleoresin of, 217
 tincture of, 217
 troches of, 217
 Cubebin, 216
 acid, 216
 Cupri sulphas, 181, 237
 Cuprum, 237
 Curare, 144
 Curarine, 144
 Cusso, 201
 Cyanids, 141
 Cypripedium, 130
 fluid extract, 130

 DATURINE, 111
 Decimal system, 24, 25
 relation to apothecaries' weight and
 measure, 25, 26
 weights and measures, 24, 25
 Decocta, 22
 Decoctions, 20, 22
 Decoctum cetrariæ, 265
 sarsaparillæ compositum, 75
 Definitions, 18-23
 Depresso-motors, 134, 145
 Diachylon ointment, 235
 Digallic acid, 229
 Digestants, 35, 37
 Digitalein, 152
 Digitalin, 152
 Digitalis, 152-154, 208
 extract of, 154
 fluid extract of, 154
 infusion of, 154
 Digitalis, preparations of, 154
 tincture of, 154
 Digitalosmin, 152
 Digitin, 152
 Digitonin, 152
 Digitoxin, 152
 Diluted hydrobromic acid, 99
 hydrocyanic acid, 139-141
 Dional, 100
 Dispensatory, 19
 Donovan's solution, 69
 Dosage, 29-32
 Dover's powder, 94, 218
 Dried gypsum, 268
 Duboisia, 113
 Duboisine, 113
 Dulcamara, 77
 fluid extract of, 77

 EFFERVESCENT MAGNESIUM CITRATE, 194
 Effervescing draught, 204
 Elastica, 269
 Elaterin, 198
 Elaterinum, 198
 Elixir aromaticum, 188
 phosphorii, 55
 phosphorus, 55
 vitriol, 57
 Ellis' magnesia, 194
 Elm, 264
 Emetine, 178
 Emplastra, 21
 Emplastrum, ammoniaci cum hydrar-
 gyri, 62, 226
 arnicæ, 167
 belladonnæ, 110
 capsici, 183
 ferri, 49
 hydrargyri, 62
 ichthyocollæ, 265
 opii, 94
 picis cum cantharidè, 258
 burgundicæ, 258
 plumbi, 235
 saponis, 235
 Emulsa, 23
 Emulsion of almonds, 263
 Emulsions, 23
 Emulsum, ammoniaci, 226
 amygdalæ, 263
 asafoetida, 128
 chloroformi, 121
 Enemata, 23, 199
 Epsom salt, 194
 Ergot, 170, 173
 extract of, 172
 fluid extract of, 173
 preparations and doses of, 172, 173
 wine of, 173
 Ergota, 170, 173
 Eriodictyon, 222, 225
 fluid extract of, 225
 Erythema, 44
 Erythroxylin, 113
 fluid extract of, 115

- Ether, 116-119, 130
 compound spirit of, 130
 preparations of, 119
 spirit of, 119
 Ethereal oil, 130
 Ethyl bromide, 122, 123
 chloride, 124
 hydrate, 146
 nitrite, 219
 oxide, 116-119
 Eucaine, 125
 Eucalyptol, 41
 Eucalyptus, 39, 41, 222, 228
 fluid extract of, 41
 Euonymus, 193
 Eupatorium, 39
 fluid extract of, 41
 Excito-motors, 130-134
 Extracta, 20
 fluida, 22
 Extracts, 20
 fluid, 20-22
 Extractum aconiti, 163
 fluidum, 163
 aloës, 192
 apocyni fluidum, 210
 arnicæ radicis, 167
 fluidum, 167
 aromaticum fluidum, 185
 asclepiadis fluidum, 224
 aspidospermatis fluidum, 224
 aurantii amari fluidum, 188
 dulcis fluidum, 188
 belladonnæ foliorum alcoholicum,
 110
 radicis fluidum, 110
 buchu fluidum, 211
 calami fluidum, 185
 cannabis indicæ, 105
 fluidum, 105
 capsici fluidum, 183
 castanæ fluidum, 233
 chimaphilæ fluidum, 212
 cimicifugæ, 159
 fluidum, 158
 cinchonæ, 45
 fluidum, 45
 cocæ fluidum, 115
 colchici radicis, 74
 fluidum, 74
 seminis fluidum, 74
 colocynthis, 197
 compositum, 197
 conii, 143
 fluidum, 144
 convallariæ fluidum, 159
 cubebæ fluidum, 217
 cusso fluidum, 201
 cypripedii fluidum, 130
 digitalis, 154
 fluidum, 154
 dulcamaræ fluidum, 77
 ergotæ, 173
 fluidum, 174
 eriodicyti fluidum, 225
 Extractum eucalypti fluidum, 41
 euonymi, 193
 eupatorii fluidum, 41
 frangulæ fluidum, 189
 gelsemii fluidum, 139
 geranii fluidum, 232
 glycyrrhizæ, 264
 fluidum, 264
 purum, 264
 gossypii radicis fluidum, 174
 grindeliæ fluidum, 222
 guaranæ fluidum, 156
 hæmatoxyli, 232
 hamamelidis fluidum, 232
 hydrastis fluidum, 47
 hyoscyami, 112
 fluidum, 112
 ipecacuanhæ fluidum, 179
 iris, 197
 fluidum, 197
 jalapæ, 196
 juglandis, 192
 krameriæ, 231
 fluidum, 231
 lappæ fluidum, 77
 leptandræ, 193
 fluidum, 193
 lobeliæ fluidum, 142
 lupulini fluidum, 106
 matico, 217
 menispermii fluidum, 76
 mezerei fluidum, 76
 nucis vomicæ, 133
 fluidum, 133
 opii, 93
 pareiræ fluidum, 212
 physostigmatis, 136
 pilocarpi fluidum, 221
 podophylli, 196
 fluidum, 196
 pruni virginianæ fluidum, 41
 rhamni purshiana fluidum, 189
 rhei, 191
 fluidum, 191
 rhois glabræ fluidum, 233
 rosæ fluidum, 232
 sabinæ fluidum, 176
 sanguinariæ fluidum, 225
 sarsaparillæ fluidum, 75
 compositum, 75
 scillæ fluidum, 208
 scoparii fluidum, 210
 scutellariæ fluidum, 130
 senegæ fluidum, 224
 sennæ fluidum, 193
 serpentariæ fluidum, 41
 spigeliæ fluidum, 200
 stillingiæ fluidum, 77
 stramonii seminis, 111
 fluidum, 111
 taraxaci, 210
 fluidum, 210
 tritici fluidum, 212
 uvæ ursi, 212
 fluidum, 212

- Extractum valerianæ fluidum, 127
 veratri viridis fluidum, 165
 viburnum fluidum, 175
 zææ fluidum, 214
 zingiberis fluidum, 185
 FENNEL, 186
 Ferri carbonas saccharatus, 49
 chloridum, 50
 citras, 50
 et ammonii citras 51
 sulphas, 51
 tartras, 51
 potassii tartras, 51
 quininæ citras, 51
 solubilis, 51
 strychninæ citras, 51
 hypophosphis, 50
 iodidum saccharatum, 50
 lactas, 51
 oxidum hydratum, 49
 cum magnesia, 49
 phosphas solubilis, 50
 pyrophosphas, 50
 sulphas, 49
 exsiccatus, 49
 granulatus, 49
 valerianas, 51
 Ferric ammonium sulphate, 51
 chloride, 50
 citrate, 50
 hypophosphite, 50
 valerianate, 51
 Ferrous carbonate, mass of, 49
 iodide, pills of, 50
 lactate, 51
 sulphate, 49
 dried, 49
 granulated, 49
 Ferrum, 47-52
 reductum, 49
 Flaxseed, 264
 tea, 264
 Flower of sulphur, 190
 Fluid extracts, 20, 22
 Fœniculum, 187
 Fomentations, 23
 Formaldehyd, 254
 Formalin, 254
 Formic aldehyd, 254
 Formyl terchloride, 119
 Fowler's solution, 69
 Frangula, 189
 GALLA, 231
 ointment of, 231
 tincture of, 231
 Gallic acid, 230
 Gallo-tannic acid, 229
 Gamboge, 197
 Gargles, 23
 Garlic, 222, 225
 Gaultheria, 186
 oil of, 186
 spirit of, 186
 Gelatine, 265
 Gelsemine, 138
 Gelseminic acid, 138
 Gelseminine, 138
 Gelsemium, 138, 139, 166
 fluid extract of, 139
 preparations and doses of, 139
 tincture of, 139
 Gentian, 37, 38, 39
 Gentiana, 37
 Gentianæ, 38
 Gentisic acid, 37
 Gentisin, 37
 Geranium, 232
 fluid extract of, 232
 Ginger, 184, 185
 fluid extract of, 185
 oleoresin of, 185
 syrup of, 185
 tincture of, 185
 troches of, 185
 Glauber's salt, 194, 195
 Glycerin, 266, 267
 Glycerinum, 266, 267
 Glycerita, 22
 Glycerite of boroglycerin, 250, 267
 of carbolic acid, 267
 glyceryl borate, 250
 hydrastis, 267
 starch, 267
 or plasmine, 265
 tannic acid, 267
 yolk of egg, 267
 Glycerites, 20, 22
 Glyceritum acidi carbolici, 251
 tannici, 230
 amyli, 265
 boroglycerini, 250
 hydrastis, 47
 vitelli, 267
 Glycyrrhiza, 264
 extract of, 264
 fluid extract of, 264, 265
 refined extract of, 264
 troches of opium and, 94
 Glycyrrhizinum ammoniatum, 265
 Gold, 64
 and sodium chloride, 64
 preparations and doses of, 64
 salts, 64
 Golden seal, 46
 Gossypii radiceis cortex, 173, 174
 Goulard's extract, 235
 cerate, 235
 Gramme, 24
 Granatum, 200
 Granules, 23
 Granville's vesicating lotion, 259
 Gray powder, 62
 Green vitriol, 49
 Griffith's anti-hectic mixture, 49
 Grindelia, 222
 fluid extract of, 222
 Grondret's vesicating ointment, 259
 Guaiac, 75, 177

- Guaiac acid, 75
 beta-resin, 75
 preparations and doses of, 75
 resina, 75
 wood, 75
 yellow, 75
 Guaiaci lignum, 75
 Guaiacin, 75
 Guaiacol, 86, 88, 126
 carbonate, 86-88
 Guaiaconic acid, 75
 Guarana, 156
 fluid extract of, 156
 Gum arabic, 263
 Gypsum, dried, 268
- HÆMATOXYLON**, 232
 extract of, 232
 Hamamelis, 232
 fluid extract of, 232
 Hedeoma, 186
 Henry's magnesia, 194
 Hoffmann's anodyne, 130
 Homatropine hydrobromate, 110
 Honey, 20, 22
 Hope's camphor mixture, 129
 Horehound, 186
 Humulus, 106
 tincture of, 106
 Husband's magnesia, 194
 Hydrargyri chloridum corrosivum,
 62, 63, 246, 261
 mite, 62, 193, 201
 cyanidum, 63
 iodidum flavum, 63
 rubrum, 63, 246
 oxidum flavum, 62
 rubrum, 62
 subsulphas flavus, 63, 181
 Hydrargyrum, 60-64
 ammoniatum, 63
 cum cretæ, 62
 preparations and doses of, 61-63
 Hydrastine, 46
 Hydrastinina hydrochloras, 47, 173
 Hydrastinine hydrochlorate, 47, 173
 Hydrastis, 42, 46, 45, 173
 fluid extract of, 47
 glycerite of, 47
 preparations and doses of, 47
 tincture of, 47
 Hydrate, 222
 Hydrated ferric oxide, 49
 with magnesia, 49
 Hydrochloric acid, 56, 58
 diluted, 58
 ether, 124
 Hydrocyanic acid, 139, 141
 Hydrogen dioxide, solution of, 244, 245
 peroxide, 244, 245
 Hydroquinone, 88, 89
 Hydrous wool-fat, 266
 Hyoscine hydrobromas, 100
 Hyoscine hydrobromate, 100
 Hyoscyamina hydrobromas, 113
 Hyoscyamina sulphas, 112, 113
 Hyoscyamine hydrobromate, 113
 sulphate, 112, 113
 Hyoscyamus, 111, 113
 extract of, 112
 fluid extract of, 112
 preparations and doses of, 112, 113
 tincture of, 112
 Hypnotics, 90-106
 Hypophosphites, 71, 72
 preparations and doses of, 72
- ICELAND MOSS**, 265
 Ichthyocollo, 265
 Igasuric acid, 131
 Igasurine, 131
 Illicium, 187
 India rubber, 269
 Indian cannabis, 104, 105
 preparations and doses, 105
 Infusa, 22
 Infusions, 20, 22
 Infusum cinchonæ, 45
 digitalis, 154
 pruni virginianæ, 41
 sennæ compositum, 193
 Ingluvin, 35, 36
 Inhalations, 23
 Injections, 23
 Iodides, 65-67
 preparations and doses of, 66, 67
 Iodine, 65-67, 246
 ointment, 66
 preparations and doses of, 66, 67
 treatment of poisoning by, 65
 Iodoform, 247, 248
 ointment, 247
 Iodoformum, 247, 248
 Iodol, 248
 Iodum, 65-67, 246
 Ipecac, 178, 218, 222
 fluid extract of, 179
 powder of opium and, 90, 179
 preparations and doses of, 179
 syrup of, 179
 tincture of opium and, 90, 179
 troches of, 179
 Ipecacuanha, 178, 179
 preparations and doses of, 179
 Ipecacuanhic acid, 179
 Iris, 197
 extract of, 197
 fluid extract of, 197
 Irish moss, 265
 Iron, 47-52
 and ammonium tartrate, 51
 citrate, 51
 potassium tartrate, 51
 quinine citrate, 51
 strychnine citrate, 51
 carbonate, pills of, 49
 compound mixture of, 49
 plaster of, 49
 preparations and doses of, 49-51
 reduced, 49

- Iron sulphate, 39
 therapeutic uses of, 48, 49
 troches of, 49
 Isaconitine, 162
 Isinglass, 265
 Isonaphthol, 253

 JABORANDINE, 220
 Jalap, 196
 compound powder of, 196
 extract of, 196
 resin of, 196
 Jalapa, 196
 James' powder, 161
 Jervine, 164
 Juglans, 191
 Juniper, 213
 compound spirit of, 213
 spirit of, 213
 Juniperus, 213

 KAIRINE, 89
 Kamala, 201
 Kermes mineral, 162
 Kino, 231
 powder, 231
 tincture of, 231
 tannic acid, 229
 Kouso, 201
 Krameria, 231
 extract of, 231
 fluid extract of, 231
 syrup of, 231
 tincture of, 231
 troches of, 231

 LABARRAQUE'S SOLUTION, 246
 Lac asafetidae, 128
 Lactic acid, 59
 preparations and doses of, 59
 Lactucarium, 106
 syrup of, 106
 tincture of, 106
 Lady Webster's dinner pills, 192
 Lanolin, 266
 Lappa, 77
 fluid extract of, 77
 Lard, 266
 Laudanum, 94
 Lauder Brunton's rule for dose in chil-
 dren, 31
 Lavender, compound tincture of, 186
 flowers, oil of, 186
 spirit of, 186
 Lead, 233, 236
 carbonate, 235, 236
 iodide, 235
 nitrate, 235
 ointment, 235
 oxide, 235
 preparations of, 235
 subacetate, cerate of, 235
 solution of, 235
 diluted, 235
 sugar of, 235

 Ledoyen's disinfecting fluid, 235
 Lemon, essence of, 169
 juice of, 169
 oil of, 169
 spirit of, 169
 Leptandra, 193
 Lime liniment, 243
 solution, 243
 syrup of, 243
 water, 243
 Limonis succus, 169
 Linimenta, 21
 Liniments, 20, 21
 Linimentum ammoniæ, 257
 belladonnæ, 110
 calcis, 243
 camphoræ, 129
 chloroformi, 121
 saponis, 129
 sinapis compositum, 257
 terebinthinæ, 216
 Linum, 264
 Liquids, 19, 20
 Liquor acidi arsenosi, 69
 ammonii acetatis, 219
 arseni et hydrargyri iodidi, 69
 ferri acetatis, 50
 chloridum, 50
 citratiss, 50
 et ammonii acetatis, 50
 nitratiss, 50
 subsulphatiss, 50
 tersulphatiss, 50
 hydrargyri nitratiss, 63, 261
 iodi compositus, 66
 magnesi citratiss, 194
 plumbi subacetatiss, 235
 dilutus, 235
 potassæ, 204
 potassii arsenitiss, 69
 citratiss, 204
 sodæ, 242
 chloratæ, 246
 sodii arsenatiss, 69
 silicatis, 269
 zinci chloridi, 239
 Liqueurice, 193, 264
 compound powder of, 193
 root, 264
 Liquores, 21
 Litharge, 235
 Lithii benzoas, 206
 carbonas, 206
 citras, 206
 effervescens, 206
 Lithium, 206
 benzoate, 206
 bromide, 206
 carbonate, 206
 citrate, effervescent, 206
 salicylate, 206
 Lobelia, 142, 222
 fluid extract of, 142
 preparations and doses of, 142
 tincture of, 142

- Lobelic acid, 142
 Lobeline, 142
 Local anæsthetics, 123-126
 Logwood, 232
 London paste, 261
 Lotions, 23
 Lozenges, 19, 20
 Lugol's solution, 66, 67
 Lupulin, 106
 fluid extract of, 106
 oleoresin of, 106
 Lupulinum, 106
 Lycopodium, 165

 MACE, 184
 Macis, 184
 Magendie's solution, 94
 Magnesia, 194
 alba, 194
 ponderosa, 194
 Magnesii carbonas, 194
 citras effervescens, 194
 sulphas, 194
 Magnesium carbonate, 194
 citrate, effervescent, 194
 solution of, 194
 sulphate, 194
 Manganese, 52, 53
 dioxide, 53
 preparations and doses of, 53
 sulphate, 53, 195
 Mangani dioxidum, 53
 sulphas, 53, 195
 Manganum, preparations and doses
 of, 52, 53
 Manna, 189
 Marrubium, 186
 Massa copaibæ, 216
 hydrargyri, 61, 62, 193
 Materia medica, 17
 proper, 17
 Matico, 217
 fluid extract of, 217
 tincture of, 217
 Matricaria, 39, 41
 Measure, wine, 24
 Measurements, approximate, 26
 Measures, 23, 26
 Mellita, 22
 Mel rosæ, 232
 Menispermum, 76
 fluid extract of, 76
 Mentha piperita, 186
 viridis, 186
 Menthol, 125, 126
 Mercurial cathartics, 193
 ointment, 62
 oleate, 62
 plaster, 62
 Mercuric chloride, 62, 63, 246, 261
 cyanide, 63
 iodide, 63, 246
 Mercurous chloride, 62, 193
 iodide, 63
 Mercury, 60-64

 Mercury, mass of, 61, 62
 metallic, 60
 preparations and doses of, 61-63
 with chalk, 62
 Methenyl chloride, 119
 Methyl-conine, 143
 Methylene bichloride, 122, 123
 Methylpyrocatechin, 86, 88
 Metrical weights, 24, 25
 Mezereon, 75, 76
 fluid extract of, 76
 Mezereum, 75, 76
 Mineral acids, 55, 59
 names of, 56
 therapeutic uses of, 56
 treatment of poisoning by, 56
 toulies, 47, 59
 Mistura cretæ, 243
 ferri compositæ, 49
 glycyrrhizæ compositæ, 265
 rhei et sodæ, 191
 Misturæ, 21
 Mixtures, 20, 21
 Morphine acetat, 94
 hydrochloras, 94
 sulphas, 94
 Morphine acetate, 94
 hydrochlorate, 94
 sulphate, 94
 troches of ipecac and, 94
 Moschus, 126, 127
 Mucilago acaciæ, 263
 sassafras medullæ, 264
 tragacanthæ, 264
 ulmi, 264
 Muriatic acid, 58
 diluted, 58
 Musk, 126, 127
 preparations of, 127
 tincture of, 127
 Mustard, 180, 181
 black, 256, 257
 liniment, 257
 papers, 257
 volatile oil of, 257
 white, 256, 257
 Mydriatic anodynes, 107, 116
 Myristica, 184

 NAPHTALIN, 253
 Naphtalinum, 253
 Naphthalin, 253
 Naphtol, 253
 Neutral mixture, 204
 Nicotianin, 144
 Nicotine, 144
 Nitre, 205
 Nitric acid, 56, 57
 diluted, 57
 Nitrites, 136-138
 preparations and doses of, 138
 Nitrogen monoxide, 122
 Nitroglycerin, 151
 triturations of, 138
 Nitrohydrochloric acid, 56-58

- Nitrohydrochloric acid, aqua regia, 58
 diluted, 58
 Nitromuriatic acid, aqua regia, 58
 diluted, 58
 Nitrous oxide, 122
 Nutmeg, 184
 oil of, 184
 spirit of, 184
 Nux vomica, 131-133
 extract of, 133
 fluid extract of, 133
 preparations and doses of, 133
 tincture of, 133

 OFFICIAL PREPARATIONS, 19
 Oil of bitter almond, 141
 cade, 227
 erigeron, 213
 gaultheria, 84
 neroli, 187
 orange flowers, 187
 santal, 213
 sweet birch, 84
 thyme, 186
 turpentine, 188, 201, 215, 257
 rectified, 216
 vitriol, 57
 wintergreen, 84
 Ointment, 266
 of ammoniated mercury, 63
 of mercurial nitrate, 63
 of potassium iodide, 66
 of red mercuric oxide, 62
 of yellow mercuric oxide, 62
 Ointments, 20, 21
 Oleata, 21
 Oleate of zinc, 238
 Oleates, 20, 21
 Oleatum hydrargyri, 62
 veratrinæ, 166
 zinci, 238
 Oleic acid, 266
 Oleoresina, aspidii, 200
 cubebæ, 217
 lupulini, 106
 piperis, 183
 zingiberis, 185
 Oleoresinæ, 23
 Oleoresins, 20, 23
 Oleum æthereum, 130
 amygdalæ amaræ, 141
 anisi, 187
 aurantii corticis, 187
 florum, 187
 betulæ volatile, 84
 cadinum, 227
 cajuputi, 185
 cari, 187
 caryophylli, 184
 chenopodii, 200
 cinnamomi, 184
 copaibæ, 216
 coriandri, 187
 erigerontis, 213
 feniculi, 187
 gaultheriæ, 84, 186
 gossypii seminis, 267
 hedeoma, 186
 juniperi, 213
 lavendulæ florum, 186
 limonis, 169
 lini, 264
 menthæ piperita, 186
 viridis, 186
 morrhuae, 70, 71
 myristicæ, 184
 phosphoratum, 55
 picis liquidæ, 227
 ricini, 189
 rosmarini, 186
 sabinæ, 176
 santali, 213
 sassafras, 76
 sesami, 267
 sinapis volatile, 257
 terebinthinæ, 201, 214, 215
 rectificatum, 216
 thymi, 186, 254, 255
 tiglii, 198
 Opii emplastrum, 94
 extractum, 93
 pulvis, 93
 Opium, 90, 95, 179
 chemical ingredients, 91
 deodoratum, 93
 deodorized, 93
 extract of, 93
 pills, 93
 plaster, 94
 powdered, 93
 powder of ipecac and, 90, 179
 preparations and doses of, 93, 94
 tincture of ipecac and, 90, 179
 treatment of poisoning by, 92
 Orange, bitter, 187
 compound spirit of, 187
 fluid extract of bitter, 187
 oil of, 187
 spirit of, 187
 syrup of, 187
 Orange flower, stronger water of, 187
 syrup of, 187
 triple water of, 187
 Orange peel, sweet, 187
 tincture of bitter and sweet, 188
 Ovis aries, 266

 PALE ROSE, 232
 ointment of, 232
 stronger water of, 232
 Pancreatin, 35, 36, 37
 Pancreatinum, 35, 36
 Papain, 35, 36, 37
 Papers, 19, 20
 Paraldehydum, 100
 Pareira, 211
 fluid extract of, 212
 Parillin, 74
 Parvules, 23
 Peculiar bitters, 42-47

- Pelletierine tannate, 200
 Pennyroyal, 186
 Pepo, 20
 Pepper, 183
 oleoresin of, 183
 Peppermint, 186
 troches of, 186
 Pepsin, 35, 36
 Pepsinum, 35, 36
 saccharatum, 36
 Peruvian balsam, 222
 Petrolatum, liquid, 267
 Petrolatum liquidum, 267
 molle, 267
 spissum, 267
 Pharmacology, 17
 Pharmacopeia, 19
 Pharmacy, 17
 Phenacetin, 81
 doses of, 81
 Phenic acid, 250
 Phenocoll, 81, 82
 Phenol, 250
 cymylic, 254, 255
 Phosphates, 71, 72
 preparations and doses of, 72
 Phosphide, 72
 Phosphite, 71
 Phosphorated oil, 55
 Phosphoric acid, 56, 58
 diluted, 58
 Phosphorus, 53, 55
 pills of, 55
 preparations and doses of, 55
 tincture of, 55
 toxic effects of, 54
 treatment of poisoning by, 54
 uses of, 54
 Physostigma, 135, 136
 extract of, 136
 preparations and doses of, 136
 tincture of, 136
 Physostigminæ salicylas, 136
 sulphas, 136
 Physostigmine salicylate, 136
 sulphate, 136
 Picroconitine, 162
 Pierotoxin, 134
 Pierotoxinum, 134
 Pills, 19, 20
 Pilocarpidine, 220
 Pilocarpinæ hydrochloras, 221
 Pilocarpine, 222
 hydrochlorate, 221
 Pilocarpos, 220, 221
 fluid extract of, 221
 preparations and doses of, 221
 Pilulæ, 20
 aloes, 192
 et asafetidæ, 128, 192
 ferri, 192
 mistaches, 192
 myrrhæ, 192
 antimonii compositæ, 162
 asafetidæ, 128
 Pilulæ catharticæ compositæ, 198
 vegetables, 198
 ferri carbonatis, 49
 iodidi, 50
 opii, 93
 phosphori, 55
 rhei, 191
 compositæ, 197
 Pimenta, 184
 oil of, 184
 Piperin, 183
 Piperinum, 183
 Pitch plaster with cantharides, 258
 Pix burgundica, 257
 liquida, 227
 Plaster of ammoniac with mercury, 62
 of Paris, 268
 Plasters, 20, 21
 Pleurisy, 224
 Plumbi acetas, 235
 carbonas, 235, 236
 iodidum, 235
 nitras, 235
 oxidum, 235
 Plumbum, 233, 236
 Plummer's pills, 162
 Podophyllotoxin, 196
 Podophyllum, 196
 Pomegranate, 200
 Potassa, 204, 260
 cum calce, 260
 solution of, 204, 242
 sulphurata, 190, 191
 with lime, 260
 Potassii acetas, 204
 bicarbonas, 204
 bichromas, 262
 bitartras, 195, 205
 bromidum, 99
 chloras, 205
 citras, 204
 effervescens, 204
 cyanidum, 141
 et sodii tartras, 195
 hypophosphis, 72
 iodidum, 66
 nitras, 205
 permanganas, 53, 245
 sulphas, 195
 Potassio-tartrate of iron, 51
 Potassium acetate, 204
 and sodium tartrate, 195
 bicarbonate, 204
 bichromate, 262
 bitartrate, 195, 205
 bromide, 99
 carbonate, 204
 chlorate, 205
 citrate, 204, 219
 effervescent, 204
 solution of, 204
 cyanide, 141
 hypophosphite, 72
 iodide, 66
 myronate, 256

Potassium nitrate, 205
 nitrite, 138
 permanganate, 52, 53, 245
 sulphate, 194
 Potato fly, 259
 Poultices, 23
 Powder of ipecac and opium, 94
 Powders, 19, 20
 Precipitated calcium carbonate, 243
 phosphate, 72
 sulphur, 190
 Preparations, official, 19
 Prepared chalk, 243
 Prescription-writing, 26, 29
Prunus virginiana, 39, 41, 228
 fluid extract of, 41
 infusion of, 41
 syrup of, 41
 Prussic acid, 139, 141
 treatment of poisoning by, 140
 Pseudoconhydrine, 143
 Pulsatilla, 167
 Pulveris, 20
Pulvis antimonialis, 161
 aromaticus, 185
 cretæ compositus, 243
 effervescens compositus, 195
 glycyrrhizæ compositus, 193
 ipecacuanhæ et opii, 94
 jalapæ compositus, 196
 morphinæ compositus, 94
 rhei compositus, 191
 Pumpkin seed, 201
 Pure bitters, 37
 Purified antimonious sulphide, 161
 Pyrocatechin, 88, 89

QUASSIA, 37, 39
Quassia, 38
Quebracho, 222
Quercus alba, 232
 Quevenne's iron, 49
 Quicksilver, 60
Quillaja, 222, 224
 tincture of, 224
Quinine, 43
Quinidine sulphas, 45
 sulphate, 42
 sulphate, 45
Quinine, 174
 bisulphas, 45
 hydrobromas, 45
 hydrochloras, 45
 sulphas, 45
 valerianas, 45
Quinine, 42, 45, 82, 174
 bisulphate, 45
 hydrobromate, 45
 hydrochlorate, 45
 sulphate, 45
 valerianate, 45

RED BARK, 42
 mercuric oxide, 62
 pepper, 183

Red precipitate ointment, 62
 rose, 232
 confection, 232
 fluid extract of, 232
 honey, 232
 syrup, 232
 saunders, 269
 wine, 148, 149
 Remedies, 18, 19
 dietetic, 18
 hygienic, 18, 19
 mechanical, 18
 pharmaceutical, 18, 19
 physical, 18
 Resin, 214, 215
 Resina, 214, 215
 jalapæ, 196
 scammonii, 197
 Resorcin, 88, 89
 Resorcinum, 88, 89
Rhamnus purshiana, 180
 Rheum, 191
 Rhigolene, 124
 Rhubarb, 191
 aromatic tincture of, 191
 syrup of, 191
 compound pills of, 191
 extract of, 191
 fluid extract of, 191
 mixture of soda and, 191
 pills, 191
 sweet, tincture of, 191
 syrup of, 191
 tincture of, 191
Rhus glabra, 232, 233
 toxicodendron, 133
 Rochelle salt, 195
Rosa centifolia, 232
 gallica, 232
 Rosemary, oil of, 186
 Rose water, 232
 Rufus pill, 192

SABINA, 176
 Saccharated ferrous iodide, 50
 pepsin, 36
 Saffron, 269
 Sage, 39
 Saigon cinnamon, 183
 Sal ammoniac, 223
 Salicin, 82-86
 preparations and doses of, 83, 84
 Salicylas, 84
 lithii, 84
 methyl, 84
 sodii, 84
 Salicylate, 84
 lithium, 84
 methyl, 84
 phenyl, 84
 physostigmine, 84
 sodium, 84
 Salicylates, 82-86
 preparations and doses of, 83, 84
 Salicylic acid, 82, 84, 219

- Salicylic acid, preparations and doses of, 83, 84
 Salicylicum, 83
 acidum, 83, 84
 Salol, 85
 doses of, 85
 Salophen, 85, 86
 Saltpetre, 205
 Salts of gold, 64
 lead, 39
 silver, 39
 Salvia, 186
 Sanguinaria, 222, 225
 fluid extract of, 225
 tincture of, 225
 Sanguinarine, 225
 Santalum rubrum, 269
 Santonica, 200
 Santonin, 200
 troches of, 200
 Santoninum, 200
 Saponin, 224
 Sarsaparilla, 74, 75
 compound fluid extract of, 75
 fluid extract of, 75
 preparations and doses of, 75
 Sassafras, 76
 medulla, 264
 oil of, 76
 pith, 264
 Savine, 176
 fluid extract of, 176
 oil of, 176
 Scammonium, 197
 Scammony, 197
 Scilla, 207, 208
 Scoparius, 210
 Scutellaria, 130
 fluid extract of, 130
 Seidlitz powder, 195
 Semi-solids, 19, 20
 Senega, 222, 223
 fluid extract of, 224
 syrup of, 224
 Senna, 193
 compound infusion of, 193
 powder of glycyrrhiza and, 193
 confection of, 193
 fluid extract of, 193
 syrup of, 193
 Serpentaria, 39, 41, 222, 228
 fluid extract of, 41
 tincture of, 41
 Sesamum, oil of, 267
 Sevum, 266
 Silver, 239, 240
 cyanide, 141, 240
 iodide, 240
 nitrate, 240, 262
 diluted, 240
 moulded, 240
 oxide, 240
 Simple bitters, 37-39
 preparations and doses of, 38, 39
 syrup, 22
 Sinalbin, 256
 Sinapis, 180, 181
 alba, 256, 257
 nigra, 256, 257
 Sinigrin, 256
 Sinilacin, 74
 Soap liniment, 129
 plaster, 235
 Socotrine aloes, 192
 Soda, 261
 caustic, 242, 261
 solution of, 242
 Sodii acetas, 207, 208
 arsenas, 69
 bicarbonas, 242
 benzoas, 252
 bisulphis, 249
 boras, 249, 250
 bromidum, 99
 carbonas, 242
 exsiccatus, 242
 hypophosphis, 72
 hyposulphis, 249
 iodidum, 66
 nitris, 138
 phosphas, 195
 pyrophosphas, 195
 sulphas, 194, 195
 sulphis, 249
 sulphocarbolas, 251
 Sodio-theobromine salicylate, 208, 209
 Sodium acetate, 207
 arsenate, 69
 benzoate, 252
 bicarbonate, 242
 troches of, 242
 bisulphate, 249
 carbonate, 242
 hypophosphite, 72, 249
 nitrite, 138
 paraphenol sulphonate, 251
 phosphate, 196
 preparations of, 242
 pyrophosphas, 195
 silicate, solution of, 269
 sulphate, 194, 195
 sulphocarbonate, 251
 Soft petrolatum, 267
 Solidified copabia, 216
 Solids, 19
 Soluble ferric phosphate, 50
 iron and quinine citrate, 51
 pyrophosphate, 50
 Solution of ammonium acetate, 219
 of arsenic and mercuric iodide, 69
 of arsenous acid, 69
 of boroglycerite, 250
 of ferric acetate, 50
 chloride, 50
 citrate, 50
 nitrate, 50
 subsulphate, 50
 sulphate, 50
 of iron and ammonium acetate, 50
 of mercuric nitrate, 63, 261

- Solution of potassium arsenite, 69
 of sodium arsenate, 69
 silicate, 269
 of zinc chloride, 239
 Solutions, 20, 21
 Spanish fly, 258
 Sparteinæ sulphas, 157, 210
 Sparteine sulphate, 157
 Spearmint, 186
 Spermaceti, 266
 Sphacelinic acid, 171
 Spigelia, 199
 fluid extract of, 200
 Spirit of glonoin, 138
 of mindererus, 219
 of nitroglycerin, 138
 of nitrous ether, 210, 211, 219
 of phosphorus, 55
 Spirits, 20, 22
 Spiritus, 22
 ætheris, 119, 130
 compositus, 130
 nitrosi, 210, 211, 219
 ammonia, 151
 aromaticus, 151
 amygdalæ amaræ, 141, 142
 anisi, 187
 aurantii, 187
 compositus, 187
 camphoræ, 129
 chloroformi, 121
 cinnamomi, 184
 frumenti, 148
 gaultheriæ, 186
 glonoini, 138
 juniperi, 213
 compositus, 213
 lavendulæ, 186
 limonis, 169
 myristicæ, 184
 phosphori, 55
 vini gallici, 148
 Squill, 207, 222, 228
 compound syrup of, 208
 fluid extract of, 208
 syrup of, 208
 tincture of, 208
 Star anise, 187
 Starch, 265
 Stillingia, 76, 77
 fluid extract of, 77
 Stramonium, 111
 fluid extract of, 111
 ointment of, 111
 preparations of, 111
 seed, extract of, 111
 tincture of, 111
 Strontii bromidum, 99
 iodidum, 67
 lactas, 207
 Strontium iodide, 67
 lactate, 207
 Strophanthin, 157
 Strophanthus, 157
 tincture of, 157
 Strychninæ, 133
 sulphas, 133
 Strychnine, 131-133, 166
 preparations and doses of, 133
 sulphate, 133
 treatment of poisoning by, 132
 Sublimated sulphur, 190
 Suet, 266
 Sugar of lead, 235
 Sulphides, 248, 249
 Sulphites, 248, 249
 Sulphonal, 100, 102
 Sulphur, flowers of, 190
 iodide, 166
 lotum, 190
 ointment, 190
 sublimatum, 190
 washed, 190
 Sulphurated antimony, 162
 lime, 249
 potassa, 190, 191, 249
 Sulphuric acid, 56, 57
 diluted, 58
 Sulphuris iodidum, 66
 Sulphurous acid, 248
 Sumach, 232, 233
 fluid extract, 233
 Suppositoria, 21
 glycerini, 267
 Suppositories, 20, 21
 Sweet spirit of nitre, 219
 Swertia chirata, 37
 Sydenham's laudanum, 94
 Syrupi, 22
 Syrup of calcium lactophosphate, 72
 of ferrous iodide, 50
 of hypophosphites, 72
 with iron, 72
 of iron, quinine, and strychnine
 phosphate, 51
 simple, 22
 Syrups, 20, 22
 Syrupus acaciæ, 263
 acidi citrici, 169
 alli, 225
 althææ, 264
 amygdalæ, 142
 aurantii, 187
 florum, 187
 calcii lactophosphatis, 72
 calcis, 243
 ferri iodidi, 50
 quininæ, et strychninæ phos-
 phatum, 50
 hypophosphitum, 72
 cum ferro, 72
 ipecacuanhæ, 179
 krameriæ, 231
 lactucarii, 106
 picis liquidæ, 227
 prunus virginianæ, 41
 rhei, 191
 aromaticus, 191
 rosæ, 232
 sarsaparillæ compositus, 161, 208

- Syrupus senegæ, 224
 sennæ, 193
 tolutanus, 227
 zingiberis, 185
 Systemic medicines, 35, 244
- TABACUM, 144, 145
 Tamarind, 189
 Tamarindus, 189
 Tanacetin, 176
 Tanacetum, 176
 Tannic acid, 229
 glycerite of, 230
 ointment of, 230
 troches of, 230
 Tansy, 176
 Tar, 222, 227
 oil of, 227
 ointment of, 227
 water, 227
 Taraxacum, 210
 extract of, 210
 fluid extract of, 210
 Tartarated antimony, 161
 Tartar emetic, 161, 181, 222
 Tartaric acid, 169
 Terebene, 222, 228
 Terebenum, 228
 Terebinthina, 214, 216
 Canadensis, 214
 Terpin, 222
 hydrate, 228
 Terpini hydras, 227
 Test for cinchona alkaloids, 43
 Tetronal, 103, 104
 Tetterswort, 197
 Thalline, 87
 Theine, 155
 Therapeutics, 17, 18
 empirical, 18
 rational, 18
 Thoroughwort, 39
 Thymol, 186, 254, 255
 Tinctura aconiti, 163
 alocs, 192
 et myrrhæ, 192
 arnicæ radicis, 167
 asafetida, 128
 belladonnæ, 110
 benzoini, 226
 composita, 226
 byroniæ, 196
 calendulæ, 76
 cannabis indicæ, 105
 cantharidis, 217, 259
 cardamomi, 185
 composita, 185
 catechu composita, 231
 cimicifugæ, 159
 cinchonæ, 45
 composita, 45
 cinnamomi, 184
 colchici, 74
 croci, 269
 cubcbæ, 217
- Tinctura capsici, 183
 digitalis, 154
 ferri chloridi, 50
 gallæ, 231
 gelsemii, 139
 guaiaci, 75
 humuli, 106
 hyoscyami, 112
 iodi, 66
 ipæcacuanhæ, 94
 kino, 231
 krameriæ, 231
 lactucarii, 106
 lavendulæ composita, 186
 lobeliæ, 142
 matico, 217
 moschi, 127
 nucis vomicæ, 133
 opii, 94
 camphorata, 94
 deodorata, 94
 physostigmatis, 136
 quillajæ, 224
 rhei, 191
 aromaticus, 192
 dulcis, 192
 sanguinarie, 225
 scillæ, 208
 serpentariæ, 41
 stramonii seminis, 111
 strophanthi, 157
 tolutani, 226
 valerianæ, 127
 ammoniata, 127
 vanillæ, 188
 veratri viridis, 165
 zingiberis, 185
 Tincturæ, 22
 Tincture of aconite, 126
 of camphorated opium, 94
 of colchicum, 74
 of deodorized opium, 94
 of elaterine, 198
 of ferric chloride, 50
 of guaiac, 75
 of iodine, 66
 of ipecac and opium, 94
 of opium, 94
 Tinctures, 20, 22
 Tobacco, 144, 145
 camphor, 144
 Tolu, balsam of, 227
 syrup of, 227
 tincture of, 227
 Tonics, mineral, 47, 59
 vegetable, 37-47
 subdivision of, 37
 Toxicology, 17
 Tragacanth, 264
 Tragacantha, 264
 Trional, 102
 Triticum, 212
 fluid extract of, 212
 Trituratio elaterine, 198
 Triturations, 20

- Triturations, 19, 20
 Troches, 20
 Trochisci, 20
 acidi tannici, 230
 ammonii chloridi, 233
 catechu, 231
 crete, 243
 cubebæ, 217
 ferri, 49
 glycyrrhizæ et opii, 94
 ipêcacuanhæ, 179
 krameriæ, 231
 menthæ piperitæ, 186
 morphinæ et ipêcacuanhæ, 94
 potassii chloratis, 206
 santonini, 200
 sodii bicarbonatis, 242
 zingiberis, 185
 Tully's powder, 94
 Turpentine, 214, 216, 222, 228
 Bordeaux, 214
 Canada, 214
 Chian, 214
 Venice, 214
 Turpeth mineral, 63, 181
 Tutty, 238

 ULMUS, 264
 Unguenta, 21
 Unguents, 21
 Unguentum, 266
 acidi carbolici, 251
 tannici, 230
 aquæ rosæ, 232
 belladonnæ, 110
 diachylon, 235
 gallæ, 231
 hydrargyri, 62
 ammoniati, 63
 nitratis, 63
 oxidi flavi, 62
 rubri, 62
 iodi, 66
 iodoformi, 248
 pici liquidæ, 227
 plumbi carbonatis, 236
 iodidi, 235
 potassii iodidi, 66
 stramonii, 111
 sulphuris, 190
 veratrinæ, 166
 zinci oxidi, 238
 Unslaked lime, 243
 Urethan, 104
 Uva ursi, 212

 VALERIAN, 127
 ammoniated tincture of, 127
 fluid extract of, 127
 preparations and doses of, 127
 tincture of, 127
 Valeriana, 127
 Vanilla, 188
 Vanillin, 188
 Vegetable cathartic pills, 198

 Vegetable tonics, 37-47
 Veratrine, 165, 166
 ointment of, 166
 oil of, 166
 uses and preparations of, 166
 Veratroidine, 164
 Veratrum viride, 164, 165, 218
 fluid extract of, 165
 preparations and doses of, 165
 tincture of, 165
 Viburnum, fluid extract of, 175
 prunifolium, 175
 Vienna paste, 260
 Vina, 22
 Vinegar of opium, 94
 Vinegars, 20, 22
 Vinum album, 148
 antimonii, 161
 colehici radidis, 74
 ergotæ, 173
 ferri amarum, 51
 citratiss, 51
 ipêcacuanhæ, 179
 opii, 94
 rubrum, 148, 149
 seminis, 74
 Viola tricolor, 189
 Virginia snakeroot, 39
 Veratrina, 165, 166
 uses and preparations of, 166
 Volatile oil of betula, 84

 WAHOO, 193
 Washed sulphur, 190
 Washes, 23
 Waters, 20, 21
 Wax, white, 266
 yellow, 266
 Weight, apothecaries', 23
 relation of, to metrical system, 25
 Weights and measures, 23-26.
 decimal, 24
 metrical, 24
 Whiskey, 148
 White arsenic, 69
 lead, 232
 oak, 232
 precipitate, 63
 vitriol, 238
 wine, 148
 Wild cherry, 39
 Wine measure, 24
 relation of, to metrical system, 28
 of antimony, 161
 of colchicum root, 74
 seed, 74
 of iron citrate, 51
 of opium, 94
 Wines, 20, 22
 Wintergreen, 186
 Wisters' cough drops, 94
 Witch hazel, 232
 Woorali, 144
 Woorara, 144
 Wormwood, 40

- YELLOW BARK, 42
 mercurial subsulphate, 61, 181
 mercuric oxide, 62
 root, 46
Young's rule for dose in children, 31
- ZEAL, 214
 fluid extract of, 214
- Zinc acetate, 238
 bromide, 99, 239
 carbonate, precipitated, 238
 chloride, 238, 261
 iodide, 239
 ointment, 238
- Zinc oxide, 238
 phosphide, 55, 239
 sulphate, 181, 238
 valerianate, 239
- Zinci acetate, 238
 bromidum, 99, 239
 carbonas præcipitatus, 238
 chloridum, 238, 261
 iodidum, 239
 phosphidum, 55, 239
 sulphas, 181, 238
 valerianas, 239
- Zincum, 238, 239
- Zingiber, 184, 185

STANDARD

Medical and Surgical Works

PUBLISHED BY

W. B. SAUNDERS, 925 Walnut Street, Philadelphia, Pa.

	PAGE		PAGE
*American Text-Book of Applied Therapeutics	4	Lockwood's Practice of Medicine	26
*American Text-Book of Dis. of Children	9	Long's Syllabus of Gynecology	20
*An American Text-Book of Diseases of the Eye, Ear, Nose, and Throat	31	Macdonald's Surgical Diagnosis and Treatment	31
*An American Text-Book of Genito-Urinary and Skin Diseases	31	McFarland's Pathogenic Bacteria	16
*American Text-Book of Gynecology	8	Mallory and Wright's Pathological Technique	31
*American Text-Book of Obstetrics	5	Martin's Surgery	28
*American Text-Book of Physiology	3	Martin's Minor Surgery, Bandaging, and Venereal Diseases	28
*American Text-Book of Practice	6	Meigs' Feeding in Early Infancy	16
*American Text-Book of Surgery	7	Moore's Orthopedic Surgery	31
Anders' Theory and Practice of Medicine	31	Morris' Materia Medica and Therapeutics	28
Ashton's Obstetrics	28	Morris' Practice of Medicine	28
Atlas of Skin Diseases	12	Morten's Nurses' Dictionary	24
Ball's Bacteriology	28	Nancrede's Anatomy and Dissection	17
Bastin's Laboratory Exercises in Botany	22	Nancrede's Anatomy	28
Beck's Surgical Asepsis	26	Norris' Syllabus of Obstetrical Lectures	21
Boislaniere's Obstetric Accidents	23	Penrose's Gynecology	31
Brockway's Physics	28	Powell's Diseases of Children	28
Burr's Nervous Diseases	26	Pye's Bandaging and Surgical Dressing	29
Butler's Materia Medica and Therapeutics	29	Raymond's Physiology	26
Cerna's Notes on the Newer Remedies	18	Rowland's Clinical Skiagraphy	14
Chapin's Compendium of Insanity	31	Saundby's Renal and Urinary Diseases	29
Chapman's Medical Jurisprudence	26	*Saunders' American Year-Book of Medicine and Surgery	32
Church and Peterson's Nervous and Mental Diseases	31	Saunders' Medical Hand-Atlases	33, 34
Clarkson's Histology	14	Saunders' Pocket Medical Formulary	19
Cohen and Eshner's Diagnosis	28	Saunders' Pocket Medical Lexicon	32
Corwin's Diagnosis of the Thorax	29	Saunders' New Series of Manuals	25, 26
Cragin's Gynecology	28	Saunders' Series of Question Compend	27, 28
Crookshank's Text-Book of Bacteriology	13	Sayre's Practice of Pharmacy	28
DaCosta's Manual of Surgery	26	Seiple's Pathology and Morbid Anatomy	28
De Schweinitz's Diseases of the Eye	15	Seiple's Legal Medicine, Toxicology, and Hygiene	28
Dorland's Obstetrics	26	Senn's Genito-Urinary Tuberculosis	31
Frothingham's Bacteriological Guide	16	Senn's Tumors	11
Garrigues' Diseases of Women	20	Senn's Syllabus of Lectures on Surgery	21
Gleason's Diseases of the Ear	28	Shaw's Nervous Diseases and Insanity	28
*Gould and Pyle's Anomalies and Curiosities of Medicine	30	Starr's Diet-Lists for Children	24
Griffin's Materia Medica and Therapeutics	26	Stelwagon's Diseases of the Skin	28
Griffith's Care of the Baby	24	Stengel's Manual of Pathology	26
Gross's Autobiography	10	Stevens' Materia Medica and Therapeutics	18
Hampton's Nursing	23	Stevens' Practice of Medicine	17
Hare's Physiology	28	Stewart's Manual of Physiology	21
Hart's Diet in Sickness and in Health	22	Stewart and Lawrance's Medical Electricity	28
Haynes' Manual of Anatomy	26	Stoney's Practical Points in Nursing	13
Heisler's Embryology	31	Sutton and Giles' Diseases of Women	26
Hirst's Obstetrics	31	Thomas's Diet-List and Sick-Room Dietary	24
Hyde's Syphilis and Venereal Diseases	26	Thornton's Dose-Book and Manual of Prescription-Writing	26
Jackson and Gleason's Diseases of the Eye, Nose, and Throat	28	Van Valzah and Nisbet's Diseases of the Stomach	31
Keating's Pronouncing Dictionary	10	Vierordt and Stuart's Medical Diagnosis	12
Keating's Life Insurance	23	Warren's Surgical Pathology	11
Keen's Operation Blanks	22	Wolff's Chemistry	28
Keen's Surgery of Typhoid Fever	31	Wolff's Examination of Urine	28
Kyle's Diseases of Nose and Throat	26		
Laine's Temperature Charts	18		

For list of the latest publications, see page 31.

The works indicated thus (*) are sold by SUBSCRIPTION (not by booksellers), usually through travelling solicitors, but they can be obtained *direct* from the office of publication (charges of shipment prepaid) by remitting the quoted prices. Full *descriptive circulars* of such works will be sent to any address upon application.

All the other books advertised in this catalogue are commonly for sale by *booksellers* in all parts of the United States; but any book will be sent by the publisher to any address (post-paid) on receipt of the price herein given.

GENERAL INFORMATION.

One Price. One price absolutely without deviation. No discounts allowed, regardless of the number of books purchased at one time. Prices on all works have been fixed extremely low, with the view to selling them strictly net and for cash.

Orders. An order accompanied by remittance will receive prompt attention, books being sent to any address in the United States, by mail or express, all charges prepaid. We prefer to send books **by express** when possible, and if sent C. O. D. **we pay all charges** for returning the money. Small orders of three dollars or less **must invariably be accompanied by remittance.**

How to Send Money by Mail. There are four ways by which money can be sent at our risk, namely: a post-office money order, an express money order, a bank-check (draft), and in a registered letter. Money sent in any other way is at the sender's risk. Silver should not be sent through the mail.

Shipments. All books, being packed in patent metal-edged boxes, necessarily reach our patrons by mail or express in excellent condition.

Subscription Books. Books in our catalogue marked "For sale by subscription only" may be secured by ordering them through any of our authorized travelling salesmen, or direct from the Philadelphia office; they are **not** for sale by booksellers. All other books in our catalogue can be procured of any bookseller at the advertised prices, or directly from us. *We handle only our own publications,* and cannot supply second-hand books nor the publications of other houses.

Latest Editions. In every instance the latest revised edition is sent.

Bindings. In ordering, be careful to state the style of binding desired—Cloth, Sheep, or Half-Morocco.

Descriptive Circulars. A complete descriptive circular, giving table of contents, etc. of any book sold by subscription only, will be sent free on application.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF PHYSIOLOGY. Edited by WILLIAM H. HOWELL, PH. D., M. D., Professor of Physiology in the Johns Hopkins University, Baltimore, Md. One handsome octavo volume of 1052 pages, fully illustrated. Prices: Cloth, \$6.00 net; Sheep or Half-Morocco, \$7.00 net.

This work is the most notable attempt yet made in America to combine in one volume the entire subject of Human Physiology by well-known teachers who have given especial study to that part of the subject upon which they write. The completed work represents the present status of the science of Physiology, particularly from the standpoint of the student of medicine and of the medical practitioner.

The collaboration of several teachers in the preparation of an elementary text-book of physiology is unusual, the almost invariable rule heretofore having been for a single author to write the entire book. One of the advantages to be derived from this collaboration method is that the more limited literature necessary for consultation by each author has enabled him to base his elementary account upon a comprehensive knowledge of the subject assigned to him; another, and perhaps the most important, advantage is that the student gains the point of view of a number of teachers. In a measure he reaps the same benefit as would be obtained by following courses of instruction under different teachers. The different standpoints assumed, and the differences in emphasis laid upon the various lines of procedure, chemical, physical, and anatomical, should give the student a better insight into the methods of the science as it exists to-day. The work will also be found useful to many medical practitioners who may wish to keep in touch with the development of modern physiology.

The main divisions of the subject-matter are as follows: General Physiology of Muscle and Nerve—Secretion—Chemistry of Digestion and Nutrition—Movements of the Alimentary Canal, Bladder, and Ureter—Blood and Lymph—Circulation—Respiration—Animal Heat—Central Nervous System—Special Senses—Special Muscular Mechanisms—Reproduction—Chemistry of the Animal Body.

CONTRIBUTORS:

HENRY P. BOWDITCH, M. D.,
Professor of Physiology, Harvard Medical School.

JOHN G. CURTIS, M. D.,
Professor of Physiology, Columbia University, N. Y. (College of Physicians and Surgeons).

HENRY H. DONALDSON, Ph. D.,
Head-Professor of Neurology, University of Chicago.

W. H. HOWELL, Ph. D., M. D.,
Professor of Physiology, Johns Hopkins University.

FREDERIC S. LEE, Ph. D.,
Adjunct Professor of Physiology, Columbia University, N. Y. (College of Physicians and Surgeons).

WARREN P. LOMBARD, M. D.,
Professor of Physiology, University of Michigan.

GRAHAM LUSK, Ph. D.,
Professor of Physiology, Yale Medical School.

W. T. PORTER, M. D.,
Assistant Professor of Physiology, Harvard Medical School.

EDWARD T. REICHERT, M. D.,
Professor of Physiology, University of Pennsylvania.

HENRY SEWALL, Ph. D., M. D.,
Professor of Physiology, Medical Department, University of Denver.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF APPLIED THERAPEUTICS. For the Use of Practitioners and Students. Edited by JAMES C. WILSON, M. D., Professor of the Practice of Medicine and of Clinical Medicine in the Jefferson Medical College. One handsome octavo volume of 1326 pages. Illustrated. Prices: Cloth, \$7.00 net; Sheep or Half-Morocco, \$8.00 net.

The arrangement of this volume has been based, so far as possible, upon modern pathologic doctrines, beginning with the intoxications, and following with infections, diseases due to internal parasites, diseases of undetermined origin, and finally the disorders of the several bodily systems—digestive, respiratory, circulatory, renal, nervous, and cutaneous. It was thought proper to include also a consideration of the disorders of pregnancy.

The list of contributors comprises the names of many who have acquired distinction as practitioners and teachers of practice, of clinical medicine, and of the specialties.

CONTRIBUTORS:

Dr. I. E. Atkinson, Baltimore, Md.
Sanger Brown, Chicago, Ill.
John B. Chapin, Philadelphia, Pa.
William C. Dabney, Charlottesville, Va.
John Chalmers DaCosta, Philada., Pa.
I. N. Danforth, Chicago, Ill.
John L. Dawson, Jr., Charleston, S. C.
F. X. Dercum, Philadelphia, Pa.
George Dock, Ann Arbor, Mich.
Robert T. Edes, Jamaica Plain, Mass.
Augustus A. Eshner, Philadelphia, Pa.
J. T. Eskridge, Denver, Col.
F. Forchheimer, Cincinnati, O.
Carl Frese, Philadelphia, Pa.
Edwin E. Graham, Philadelphia, Pa.
John Guit ras, Philadelphia, Pa.
Frederick P. Henry, Philadelphia, Pa.
Guy Hinsdale, Philadelphia, Pa.
Orville Horwitz, Philadelphia, Pa.
W. W. Johnston, Washington, D. C.
Ernest Laplace, Philadelphia, Pa.
A. Laveran, Paris, France.

Dr. James Hendrie Lloyd, Philadelphia, Pa.
John Noland Mackenzie, Baltimore, Md.
J. W. McLaughlin, Austin, Texas.
A. Lawrence Mason, Boston, Mass.
Charles K. Mills, Philadelphia, Pa.
John K. Mitchell, Philadelphia, Pa.
W. P. Northrup, New York City.
William Osler, Baltimore, Md.
Frederick A. Packard, Philadelphia, Pa.
Theophilus Parvin, Philadelphia, Pa.
Beaven Rake, London, England.
E. O. Shakespeare, Philadelphia, Pa.
Wharton Sinkler, Philadelphia, Pa.
Louis Starr, Philadelphia, Pa.
Henry W. Stelwagon, Philadelphia, Pa.
James Stewart, Montreal, Canada.
Charles G. Stockton, Buffalo, N. Y.
James Tyson, Philadelphia, Pa.
Victor C. Vaughan, Ann Arbor, Mich.
James T. Whitaker, Cincinnati, O.
J. C. Wilson, Philadelphia, Pa.

The articles, with two exceptions, are the contributions of American writers. Written from the standpoint of the practitioner, the aim of the work is to facilitate the application of knowledge to the prevention, the cure, and the alleviation of disease. The endeavor throughout has been to conform to the title of the book—Applied Therapeutics—to indicate the course of treatment to be pursued at the bedside, rather than to name a list of drugs that have been used at one time or another.

While the scientific superiority and the practical desirability of the metric system of weights and measures is admitted, it has not been deemed best to discard entirely the older system of figures, so that both sets have been given where occasion demanded.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF OBSTETRICS. Edited by RICHARD C. NORRIS, M. D.; Art Editor, ROBERT L. DICKINSON, M. D. One handsome octavo volume of over 1000 pages, with nearly 900 colored and half-tone illustrations. Prices: Cloth, \$7.00; Sheep or Half-Morocco, \$8.00.

The advent of each successive volume of the *series* of the AMERICAN TEXT-BOOKS has been signalized by the most flattering comment from both the Press and the Profession. The high consideration received by these text-books, and their attainment to an authoritative position in current medical literature, have been matters of deep *international* interest, which finds its fullest expression in the demand for these publications from all parts of the civilized world.

In the preparation of the "AMERICAN TEXT-BOOK OF OBSTETRICS" the editor has called to his aid proficient collaborators whose professional prominence entitles them to recognition, and whose disquisitions exemplify **Practical Obstetrics**. While these writers were each assigned special themes for discussion, the correlation of the subject-matter is, nevertheless, such as ensures logical connection in treatment, the deductions of which thoroughly represent the latest advances in the science, and which elucidate *the best modern methods of procedure*.

The more conspicuous feature of the treatise is its wealth of illustrative matter. The production of the illustrations had been in progress for several years, under the personal supervision of Robert L. Dickinson, M. D., to whose artistic judgment and professional experience is due the **most sumptuously illustrated work of the period**. By means of the photographic art, combined with the skill of the artist and draughtsman, conventional illustration is superseded by rational methods of delineation.

Furthermore, the volume is a revelation as to the possibilities that may be reached in mechanical execution, through the unsparing hand of its publisher.

CONTRIBUTORS:

Dr. James C. Cameron.
Edward P. Davis.
Robert L. Dickinson.
Charles Warrington Earle.
James H. Etheridge.
Henry J. Garrigues.
Barton Cooke Hirst.
Charles Jewett.

Dr. Howard A. Kelly.
Richard C. Norris.
Chauncey D. Palmer.
Theophilus Parvin.
George A. Piersol.
Edward Reynolds.
Henry Schwarz.

"At first glance we are overwhelmed by the magnitude of this work in several respects, viz.: First, by the size of the volume, then by the array of eminent teachers in this department who have taken part in its production, then by the profuseness and character of the illustrations, and last, but not least, the conciseness and clearness with which the text is rendered. This is an entirely new composition, embodying the highest knowledge of the art as it stands to-day by authors who occupy the front rank in their specialty, and there are many of them. We cannot turn over these pages without being struck by the superb illustrations which adorn so many of them. We are confident that this most practical work will find instant appreciation by practitioners as well as students."—*New York Medical Times*.

Permit me to say that your American Text-Book of Obstetrics is the most magnificent medical work that I have ever seen. I congratulate you and thank you for this superb work, which alone is sufficient to place you first in the ranks of medical publishers.

With profound respect I am sincerely yours,

ALEX. J. C. SKENE.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK ON THE THEORY AND PRACTICE OF MEDICINE. By American Teachers. Edited by WILLIAM PEPPER, M. D., LL.D., Provost and Professor of the Theory and Practice of Medicine and of Clinical Medicine in the University of Pennsylvania. Complete in two handsome royal-octavo volumes of about 1000 pages each, with illustrations to elucidate the text wherever necessary. Price per Volume: Cloth, \$5.00 net; Sheep or Half-Morocco, \$6.00 net.

VOLUME I. CONTAINS:

Hygiene.—Fever (Ephemeral, Simple Continued, Typhus, Typhoid, Epidemic Cerebro-spinal Meningitis, and Relapsing).—Scarlatina, Measles, Röteln, Variola, Varioloid, Vaccinia, Varicella, Mumps, Whooping-cough, Anthrax, Hydrophobia, Trichinosis, Actino-

mycosis, Glanders, and Tetanus.—Tuberculosis, Scrofula, Syphilis, Diphtheria, Erysipelas, Malaria, Cholera, and Yellow Fever.—Nervous, Muscular, and Mental Diseases etc.

VOLUME II. CONTAINS:

Urine (Chemistry and Microscopy).—Kidney and Lungs.—Air-passages (Larynx and Bronchi) and Pleura.—Pharynx, Oesophagus, Stomach and Intestines (including Intestinal Parasites), Heart, Aorta, Arteries and Veins.

—Peritoneum, Liver, and Pancreas.—Diathetic Diseases (Rheumatism, Rheumatoid Arthritis, Gout, Lithæmia, and Diabetes).—Blood and Spleen.—Inflammation, Embolism, Thrombosis, Fever, and Bacteriology.

The articles are not written as though addressed to students in lectures, but are exhaustive descriptions of diseases, with the newest facts as regards Causation, Symptomatology, Diagnosis, Prognosis, and Treatment, including a large number of approved formulæ. The recent advances made in the study of the bacterial origin of various diseases are fully described, as well as the bearing of the knowledge so gained upon prevention and cure. The subjects of Bacteriology as a whole and of Immunity are fully considered in a separate section.

Methods of diagnosis are given the most minute and careful attention, thus enabling the reader to learn the very latest methods of investigation without consulting works specially devoted to the subject.

CONTRIBUTORS:

Dr. J. S. Billings, Philadelphia.
Francis Delafield, New York.
Reginald H. Fitz, Boston.
James W. Holland, Philadelphia.
Henry M. Lyman, Chicago.
William Osler, Baltimore.

Dr. William Pepper, Philadelphia.
W. Gilman Thompson, New York.
W. H. Welch, Baltimore.
James T. Whittaker, Cincinnati.
James C. Wilson, Philadelphia.
Horatio C. Wood, Philadelphia.

"We reviewed the first volume of this work, and said: 'It is undoubtedly one of the best text-books on the practice of medicine which we possess.' A consideration of the second and last volume leads us to modify that verdict and to say that the completed work is, in our opinion, THE BEST of its kind it has ever been our fortune to see. It is complete, thorough, accurate, and clear. It is well written, well arranged, well printed, well illustrated, and well bound. It is a model of what the modern text-book should be."—*New York Medical Journal*.

"A library upon modern medical art. The work must promote the wider diffusion of sound knowledge."—*American Lancet*.

"A trusty counsellor for the practitioner or senior student, on which he may implicitly rely."—*Edinburgh Medical Journal*.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF SURGERY. Edited by WILLIAM W. KEEN, M. D., LL.D., and J. WILLIAM WHITE, M. D., PH. D. Forming one handsome royal-octavo volume of 1250 pages (10 x 7 inches), with 500 wood-cuts in text, and 37 colored and half-tone plates, many of them engraved from original photographs and drawings furnished by the authors. Prices: Cloth, \$7.00 net; Sheep or Half-Morocco, \$8.00 net.

SECOND EDITION, REVISED AND ENLARGED,

With a Section devoted to "The Use of the Röntgen Rays in Surgery."

The want of a text-book which could be used by the practitioner and at the same time be recommended to the medical student has been deeply felt, especially by teachers of surgery; hence, when it was suggested to a number of these that it would be well to unite in preparing a text-book of this description, great unanimity of opinion was found to exist, and the gentlemen below named gladly consented to join in its production. While there is no distinctive American Surgery, yet America has contributed very largely to the progress of modern surgery, and among the foremost of those who have aided in developing this art and science will be found the authors of the present volume. All of them are teachers of surgery in leading medical schools and hospitals in the United States and Canada.

Especial prominence has been given to Surgical Bacteriology, a feature which is believed to be unique in a surgical text-book in the English language. Asepsis and Antisepsis have received particular attention. The text is brought well up to date in such important branches as cerebral, spinal, intestinal, and pelvic surgery, the most important and newest operations in these departments being described and illustrated.

The text of the entire book has been submitted to all the authors for their mutual criticism and revision—an idea in book-making that is entirely new and original. The book as a whole, therefore, expresses on all the important surgical topics of the day the consensus of opinion of the eminent surgeons who have joined in its preparation.

One of the most attractive features of the book is its illustrations. Very many of them are original and faithful reproductions of photographs taken directly from patients or from specimens.

CONTRIBUTORS:

Dr. Charles H. Burnett, Philadelphia.
Phineas S. Conner, Cincinnati.
Frederic S. Dennis, New York.
William W. Keen, Philadelphia.
Charles B. Nancrede, Ann Arbor, Mich.
Roswell Park, Buffalo, N. Y.
Lewis S. Pilcher, New York.

Dr. Nicholas Senn, Chicago.
Francis J. Shepherd, Montreal, Canada.
Lewis A. Stimson, New York.
William Thomson, Philadelphia.
J. Collins Warren, Boston.
J. William White, Philadelphia.

"If this text-book is a fair reflex of the present position of American surgery, we must admit it is of a very high order of merit, and that English surgeons will have to look very carefully to their laurels if they are to preserve a position in the van of surgical practice."—*London Lancet*.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF GYNECOLOGY, MEDICAL AND SURGICAL, for the use of Students and Practitioners.

Edited by J. M. BALDY, M. D. Forming a handsome royal-octavo volume, with 341 illustrations in text and 38 colored and half-tone plates. Prices: Cloth, \$6.00 net; Sheep or Half-Morocco, \$7.00 net.

SECOND EDITION, THOROUGHLY REVISED.

In this volume all anatomical descriptions, excepting those essential to a clear understanding of the text, have been omitted, the illustrations being largely depended upon to elucidate the anatomy of the parts. This work, which is thoroughly practical in its teachings, is intended, as its title implies, to be a working text-book for physicians and students. A clear line of treatment has been laid down in every case, and although no attempt has been made to discuss mooted points, still the most important of these have been noted and explained. The operations recommended are fully illustrated, so that the reader, having a picture of the procedure described in the text under his eye, cannot fail to grasp the idea. All extraneous matter and discussions have been carefully excluded, the attempt being made to allow no unnecessary details to cumber the text. The subject-matter is brought up to date at every point, and the work is as nearly as possible the combined opinions of the ten specialists who figure as the authors.

CONTRIBUTORS:

Dr. Henry T. Byford.
John M. Baldy.
Edwin Cragin.
J. H. Etheridge.
William Goodell.

Dr. Howard A. Kelly.
Florian Krug.
E. E. Montgomery.
William R. Pryor.
George M. Tuttle.

"The most notable contribution to gynecological literature since 1887, . . . and the most complete exponent of gynecology which we have. No subject seems to have been neglected, . . . and the gynecologist and surgeon, and the general practitioner who has any desire to practise diseases of women, will find it of practical value. In the matter of illustrations and plates the book surpasses anything we have seen."—*Boston Medical and Surgical Journal*.

"A valuable addition to the literature of Gynecology. The writers are progressive, aggressive, and earnest in their convictions."—*Medical News*, Philadelphia.

"A thoroughly modern text-book, and gives reliable and well-tempered advice and instruction."—*Edinburgh Medical Journal*.

"The harmony of its conclusions and the homogeneity of its style give it an individuality which suggests a single rather than a multiple authorship."—*Annals of Surgery*.

"It must command attention and respect as a worthy representation of our advanced clinical teaching."—*American Journal of Medical Sciences*.

For Sale by Subscription.

AN AMERICAN TEXT-BOOK OF THE DISEASES OF CHILDREN. By American Teachers. Edited by LOUIS STARR, M. D., assisted by THOMPSON S. WESTCOTT, M. D. In one handsome royal-8vo volume of 1250 pages, profusely illustrated with wood-cuts, half-tone and colored plates. Net Prices: Cloth, \$7.00; Sheep or Half-Morocco, \$8.00.

SECOND EDITION, REVISED AND ENLARGED.

The plan of this work embraces a series of original articles written by some sixty well-known pædiatrists, representing collectively the teachings of the most prominent medical schools and colleges of America. The work is intended to be a PRACTICAL book, suitable for constant and handy reference by the practitioner and the advanced student.

Especial attention has been given to the latest accepted teachings upon the etiology, symptoms, pathology, diagnosis, and treatment of the disorders of children, with the introduction of many special formulæ and therapeutic procedures.

Special chapters embrace at unusual length the Diseases of the Eye, Ear, Nose and Throat, and the Skin; while the introductory chapters cover fully the important subjects of Diet, Hygiene, Exercise, Bathing, and the Chemistry of Food. Tracheotomy, Intubation, Circumcision, and such minor surgical procedures coming within the province of the medical practitioner are carefully considered.

CONTRIBUTORS:

Dr. S. S. Adams, Washington.
 John Ashhurst, Jr., Philadelphia.
 A. D. Blackader, Montreal, Canada.
 David Bovaird, New York.
 Dillon Brown, New York.
 Edward M. Buckingham, Boston.
 Charles W. Burr, Philadelphia.
 W. E. Casselberry, Chicago.
 Henry Dwight Chapin, New York.
 W. S. Christopher, Chicago.
 Archibald Church, Chicago.
 Floyd M. Crandall, New York.
 Andrew F. Currier, New York.
 Roland G. Curtin, Philadelphia.
 J. M. DaCosta, Philadelphia.
 I. N. Danforth, Chicago.
 Edward P. Davis, Philadelphia.
 John B. Deaver, Philadelphia.
 G. E. de Schweinitz, Philadelphia.
 John Dorning, New York.
 Charles Warrington Earle, Chicago.
 Wm. A. Edwards, San Diego, Cal.
 F. Forchheimer, Cincinnati.
 J. Henry Fruitnight, New York.
 J. P. Crozer Griffith, Philadelphia.
 W. A. Hardaway, St. Louis.
 M. P. Hatfield, Chicago.
 Barton Cooke Hirst, Philadelphia.
 H. Illoway, Cincinnati.
 Henry Jackson, Boston.
 Charles G. Jennings, Detroit.
 Henry Koplik, New York.

Dr. Thomas S. Latimer, Baltimore.
 Albert R. Leeds, Hoboken, N. J.
 J. Hendrie Lloyd, Philadelphia.
 George Roe Lockwood, New York.
 Henry M. Lyman, Chicago.
 Francis T. Miles, Baltimore.
 Charles K. Mills, Philadelphia.
 James E. Moore, Minneapolis.
 F. Gordon Morrill, Boston.
 John H. Musser, Philadelphia.
 Thomas R. Neilson, Philadelphia.
 W. P. Northrup, New York.
 William Osler, Baltimore.
 Frederick A. Packard, Philadelphia.
 William Pepper, Philadelphia.
 Frederick Peterson, New York.
 W. T. Plant, Syracuse, New York.
 William M. Powell, Atlantic City.
 B. Alexander Randall, Philadelphia.
 Edward O. Shakespeare, Philadelphia.
 F. C. Shattuck, Boston.
 J. Lewis Smith, New York.
 Louis Starr, Philadelphia.
 M. Allen Starr, New York.
 Charles W. Townsend, Boston.
 James Tyson, Philadelphia.
 W. S. Thayer, Baltimore.
 Victor C. Vaughan, Ann Arbor, Mich.
 Thompson S. Westcott, Philadelphia.
 Henry R. Wharton, Philadelphia.
 J. William White, Philadelphia.
 J. C. Wilson, Philadelphia.

A NEW PRONOUNCING DICTIONARY OF MEDICINE, with Phonetic Pronunciation, Accentuation, Etymology, etc. By JOHN M. KEATING, M. D., LL.D., Fellow of the College of Physicians of Philadelphia; Vice-President of the American Pædiatric Society; Ex-President of the Association of Life Insurance Medical Directors; Editor "Cyclopædia of the Diseases of Children," etc.; and HENRY HAMILTON, author of "A New Translation of Virgil's Æneid into English Rhyme;" co-author of "Saunders' Medical Lexicon," etc.; with the Collaboration of J. CHALMERS DACOSTA, M. D., and FREDERICK A. PACKARD, M. D. With an Appendix containing important Tables of Bacilli, Micrococci, Leucomaïnes, Ptomaïnes, Drugs and Materials used in Antiseptic Surgery, Poisons and their Antidotes, Weights and Measures, Thermometric Scales, New Official and Unofficial Drugs, etc. One very attractive volume of over 800 pages. Second Revised Edition. Prices: Cloth, \$5.00 net; Sheep or Half-Morocco, \$6.00 net; with Denison's Patent Ready-Reference Index; without patent index, Cloth, \$4.00 net; Sheep or Half-Morocco, \$5.00 net.

PROFESSIONAL OPINIONS.

"I am much pleased with Keating's Dictionary, and shall take pleasure in recommending it to my classes."

HENRY M. LYMAN, M. D.,

Professor of Principles and Practice of Medicine, Rush Medical College, Chicago, Ill.

"I am convinced that it will be a very valuable adjunct to my study-table, convenient in size and sufficiently full for ordinary use."

C. A. LINDSLEY, M. D.,

*Professor of Theory and Practice of Medicine, Medical Dept. Yale University;
Secretary Connecticut State Board of Health, New Haven, Conn.*

AUTOBIOGRAPHY OF SAMUEL D. GROSS, M. D., Emeritus Professor of Surgery in the Jefferson Medical College of Philadelphia, with Reminiscences of His Times and Contemporaries. Edited by his sons, SAMUEL W. GROSS, M. D., LL.D., late Professor of Principles of Surgery and of Clinical Surgery in the Jefferson Medical College, and A. HALLER GROSS, A. M., of the Philadelphia Bar. Preceded by a Memoir of Dr. Gross, by the late Austin Flint, M. D., LL.D. In two handsome volumes, each containing over 400 pages, demy 8vo, extra cloth, gilt tops, with fine Frontispiece engraved on steel. Price per Volume, \$2.50 net.

This autobiography, which was continued by the late eminent surgeon until within three months of his death, contains a full and accurate history of his early struggles, trials, and subsequent successes, told in a singularly interesting and charming manner, and embraces short and graphic pen-portraits of many of the most distinguished men—surgeons, physicians, divines, lawyers, statesmen, scientists, etc.—with whom he was brought in contact in America and in Europe; the whole forming a retrospect of more than three-quarters of a century.

SURGICAL PATHOLOGY AND THERAPEUTICS. By JOHN COLLINS WARREN, M. D., LL.D., Professor of Surgery, Medical Department Harvard University; Surgeon to the Massachusetts General Hospital, etc. A handsome octavo volume of 832 pages, with 136 relief and lithographic illustrations, 33 of which are printed in colors, and all of which were drawn by William J. Kaula from original specimens. Prices: Cloth, \$6.00 net; Half-Morocco, \$7.00 net.

"The volume is for the bedside, the amphitheatre, and the ward. It deals with things not as we see them through the microscope alone, but as the practitioner sees their effect in his patients; not only as they appear in and affect culture-media, but also as they influence the human body; and, following up the demonstrations of the nature of diseases, the author points out their logical treatment." (*New York Medical Journal*). "It is the handsomest specimen of book-making * * * that has ever been issued from the American medical press" (*American Journal of the Medical Sciences*, Philadelphia).

Without Exception, the Illustrations are the Best ever Seen in a Work of this Kind.

"A most striking and very excellent feature of this book is its illustrations. Without exception, from the point of accuracy and artistic merit, they are the best ever seen in a work of this kind. * * * Many of those representing microscopic pictures are so perfect in their coloring and detail as almost to give the beholder the impression that he is looking down the barrel of a microscope at a well-mounted section."—*Annals of Surgery*, Philadelphia.

PATHOLOGY AND SURGICAL TREATMENT OF TUMORS.

By N. SENN, M. D., Ph. D., LL. D., Professor of Practice of Surgery and of Clinical Surgery, Rush Medical College; Professor of Surgery, Chicago Polyclinic; Attending Surgeon to Presbyterian Hospital; Surgeon-in-Chief, St. Joseph's Hospital, Chicago. One volume of 710 pages, with 515 engravings, including full-page colored plates. Prices: Cloth, \$6.00 net; Half-Morocco, \$7.00 net.

Books specially devoted to this subject are few, and in our text-books and systems of surgery this part of surgical pathology is usually condensed to a degree incompatible with its scientific and clinical importance. The author spent many years in collecting the material for this work, and has taken great pains to present it in a manner that should prove useful as a text-book for the student, a work of reference for the busy practitioner, and a reliable, safe guide for the surgeon. The more difficult operations are fully described and illustrated. More than *one hundred* of the illustrations are original, while the remainder were selected from books and medical journals not readily accessible.

"The most exhaustive of any recent book in English on this subject. It is well illustrated, and will doubtless remain as the principal monograph on the subject in our language for some years. The book is handsomely illustrated and printed, . . . and the author has given a notable and lasting contribution to surgery."—*Journal of American Medical Association*, Chicago.

MEDICAL DIAGNOSIS. By Dr. OSWALD VIERORDT, Professor of Medicine at the University of Heidelberg. Translated, with additions, from the Fifth Enlarged German Edition, with the author's permission, by FRANCIS H. STUART, A. M., M. D. In one handsome royal-octavo volume of 600 pages. 194 fine wood-cuts in the text, many of them in colors. Prices: Cloth, \$4.00 net; Sheep or Half-Morocco, \$5.00 net.

FOURTH AMERICAN EDITION, FROM THE FIFTH REVISED AND ENLARGED GERMAN EDITION.

In this work, as in no other hitherto published, are given full and accurate explanations of the phenomena observed at the bedside. It is distinctly a clinical work by a master teacher, characterized by thoroughness, fulness, and accuracy. It is a mine of information upon the points that are so often passed over without explanation. Especial attention has been given to the germ-theory as a factor in the origin of disease.

This valuable work is now published in German, English, Russian, and Italian. The issue of a third American edition within two years indicates the favor with which it has been received by the profession.

THE PICTORIAL ATLAS OF SKIN DISEASES AND SYPHILITIC AFFECTIONS. (American Edition.) Translation from the French. Edited by J. J. PRINGLE, M. B., F. R. C. P., Assistant Physician to, and Physician to the department for Diseases of the Skin at, the Middlesex Hospital, London. Photo-lithochromes from the famous models of dermatological and syphilitic cases in the Museum of the Saint-Louis Hospital, Paris, with explanatory wood-cuts and letter-press. In 12 Parts, at \$3.00 per Part. Parts 1 to 8 now ready.

"The plates are beautifully executed."—JONATHAN HUTCHINSON, M. D. (London Hospital).

"The plates in this Atlas are remarkably accurate and artistic reproductions of *typical* examples of skin disease. The work will be of great value to the practitioner and student."—WILLIAM ANDERSON, M. D. (St. Thomas Hospital).

"If the succeeding parts of this Atlas are to be similar to Part 1, now before us, we have no hesitation in cordially recommending it to the favorable notice of our readers as one of the finest dermatological atlases with which we are acquainted."—*Glasgow Medical Journal*, Aug., 1895.

"Of all the atlases of skin diseases which have been published in recent years, the present one promises to be of greatest interest and value, especially from the standpoint of the general practitioner."—*American Medico-Surgical Bulletin*, Feb. 22, 1896.

"The introduction of explanatory wood-cuts in the text is a novel and most important feature which greatly furthers the easier understanding of the excellent plates, than which nothing, we venture to say, has been seen better in point of correctness, beauty, and general merit."—*New York Medical Journal*, Feb. 15, 1896.

"An interesting feature of the Atlas is the descriptive text, which is written for each picture by the physician who treated the case or at whose instigation the models have been made. We predict for this truly beautiful work a large circulation in all parts of the medical world where the names *St. Louis* and *Baretta* have preceded it."—*Medical Record*, N. Y., Feb. 1, 1896.

PRACTICAL POINTS IN NURSING. For Nurses in Private Practice. By EMILY A. M. STONEY, Graduate of the Training-School for Nurses, Lawrence, Mass.; Superintendent of the Training-School for Nurses, Carney Hospital, South Boston, Mass. 456 pages, handsomely illustrated with 73 engravings in the text, and 9 colored and half-tone plates. Cloth. Price, \$1.75 net.

SECOND EDITION, THOROUGHLY REVISED.

In this volume the author explains, in popular language and in the shortest possible form, the entire range of *private* nursing as distinguished from *hospital* nursing, and the nurse is instructed how best to meet the various emergencies of medical and surgical cases when distant from medical or surgical aid or when thrown on her own resources.

An especially valuable feature of the work will be found in the directions to the nurse how to *improvise* everything ordinarily needed in the sick-room, where the embarrassment of the nurse, owing to the want of proper appliances, is frequently extreme.

The work has been logically divided into the following sections :

- I. The Nurse : her responsibilities, qualifications, equipment, etc.
- II. The Sick-Room : its selection, preparation, and management.
- III. The Patient : duties of the nurse in medical, surgical, obstetric, and gynecologic cases.
- IV. Nursing in Accidents and Emergencies.
- V. Nursing in Special Medical Cases.
- VI. Nursing of the New-born and Sick Children.
- VII. Physiology and Descriptive Anatomy.

The APPENDIX contains much information in compact form that will be found of great value to the nurse, including Rules for Feeding the Sick ; Recipes for Invalid Foods and Beverages ; Tables of Weights and Measures ; Table for Computing the Date of Labor ; List of Abbreviations ; Dose-List ; and a full and complete Glossary of Medical Terms and Nursing Treatment.

"This is a well-written, eminently practical volume, which covers the entire range of private nursing as distinguished from hospital nursing, and instructs the nurse how best to meet the various emergencies which may arise and how to prepare everything ordinarily needed in the illness of her patient."—*American Journal of Obstetrics and Diseases of Women and Children*, Aug., 1896.

A TEXT-BOOK OF BACTERIOLOGY, including the Etiology and Prevention of Infective Diseases and an account of Yeasts and Moulds, Hæmatozoa, and Psorospersms. By EDGAR M. CROOKSHANK, M. B., Professor of Comparative Pathology and Bacteriology, King's College, London. A handsome octavo volume of 700 pages, with 273 engravings in the text, and 22 original and colored plates. Price, \$6.50 net.

This book, though nominally a Fourth Edition of Professor Crookshank's "MANUAL OF BACTERIOLOGY," is practically a new work, the old one having been reconstructed, greatly enlarged, revised throughout, and largely rewritten, forming a text-book for the Bacteriological Laboratory, for Medical Officers of Health, and for Veterinary Inspectors.

A TEXT-BOOK OF HISTOLOGY, DESCRIPTIVE AND PRACTICAL. For the Use of Students. By ARTHUR CLARKSON, M. B., C. M., Edin., formerly Demonstrator of Physiology in the Owen's College, Manchester; late Demonstrator of Physiology in the Yorkshire College, Leeds. Large 8vo, 554 pages, with 22 engravings in the text, and 174 beautifully colored original illustrations. Price, strongly bound in Cloth, \$6.00 net.

The purpose of the writer in this work has been to furnish the student of Histology, in one volume, with both the descriptive and the practical part of the science. The first two chapters are devoted to the consideration of the general methods of Histology; subsequently, in each chapter, the structure of the tissue or organ is first systematically described, the student is then taken tutorially over the specimens illustrating it, and, finally, an appendix affords a short note of the methods of preparation.

"We would most cordially recommend it to all students of histology."—*Dublin Medical Journal*.

"It is pleasant to give unqualified praise to the colored illustrations; . . . the standard is high, and many of them are not only extremely beautiful, but very clear and demonstrative. . . . The plan of the book is excellent."—*Liverpool Medical Journal*.

ARCHIVES OF CLINICAL SKIAGRAPHY. By SYDNEY ROWLAND, B. A., Camb. A series of collotype illustrations, with descriptive text, illustrating the applications of the New Photography to Medicine and Surgery. Price, per Part, \$1.00. Parts I. to V. now ready.

The object of this publication is to put on record in permanent form some of the most striking applications of the new photography to the needs of Medicine and Surgery.

The progress of this new art has been so rapid that, although Prof. Röntgen's discovery is only a thing of yesterday, it has already taken its place among the approved and accepted aids to diagnosis.

WATER AND WATER SUPPLIES. By JOHN C. THRESH, D. Sc., M. B., D. P. H., Lecturer on Public Health, King's College, London; Editor of the "Journal of State Medicine," etc. 12mo, 438 pages, illustrated. Handsomely bound in Cloth, with gold side and back stamps. Price, \$2.25 net.

This work will furnish any one interested in public health the information requisite for forming an opinion as to whether any supply or proposed supply is sufficiently wholesome and abundant, and whether the cost can be considered reasonable.

The work does not pretend to be a treatise on Engineering, yet it contains sufficient detail to enable any one who has studied it to consider intelligently any scheme which may be submitted for supplying a community with water.

DISEASES OF THE EYE. A Hand-Book of Ophthalmic Practice. By G. E. DE SCHWEINITZ, M. D., Professor of Ophthalmology in the Jefferson Medical College, Philadelphia, etc. A handsome royal-octavo volume of 679 pages, with 256 fine illustrations, many of which are original, and 2 chromo-lithographic plates. Prices: Cloth, \$4.00 net; Sheep or Half-Morocco, \$5.00 net.

The object of this work is to present to the student, and to the practitioner who is beginning work in the fields of ophthalmology, a plain description of the optical defects and diseases of the eye. To this end special attention has been paid to the clinical side of the question; and the method of examination, the symptomatology leading to a diagnosis, and the treatment of the various ocular defects have been brought into prominence.

THIRD EDITION, THOROUGHLY REVISED.

The entire book has been thoroughly revised. In addition to this general revision, special paragraphs on the following new matter have been introduced: Filamentous Keratitis, Blood-staining of the Cornea, Essential Phthisis Bulbi, Foreign Bodies in the Lens, Circinate Retinitis, Symmetrical Changes at the Macula Lutea in Infancy, Hyaline Bodies in the Papilla, Monocular Diplopia, Subconjunctival Injections of Germicides, Infiltration-Anæsthesia, and Sterilization of Collyria. Brief mention of Ophthalmia Nodosa, Electric Ophthalmia, and Angioid Streaks in the Retina also finds place. An Appendix has been added, containing a full description of the method of determining the corneal astigmatism with the ophthalmometer of Javal and Schiötz, and the rotations of the eyes with the tropometer of Stevens. The chapter on Operations has been enlarged and rewritten.

"A clearly written, comprehensive manual. . . . One which we can commend to students as a reliable text-book, written with an evident knowledge of the wants of those entering upon the study of this special branch of medical science."—*British Medical Journal*.

"The work is characterized by a lucidity of expression which leaves the reader in no doubt as to the meaning of the language employed. . . . We know of no work in which these diseases are dealt with more satisfactorily, and indications for treatment more clearly given, and in harmony with the practice of the most advanced ophthalmologists."—*Maritime Medical News*.

"It is hardly too much to say that for the student and practitioner beginning the study of Ophthalmology, it is the best single volume at present published."—*Medical News*.

"The latest and one of the best books on Ophthalmology. The book is thoroughly up to date, and is certainly a work which not only commends itself to the student, but is a ready reference for the busy practitioner."—*International Medical Review*.

PROFESSIONAL OPINIONS.

"A work that will meet the requirements not only of the specialist, but of the general practitioner in a rare degree. I am satisfied that unusual success awaits it."

WILLIAM PEPPER, M. D.

Provost and Professor of Theory and Practice of Medicine and Clinical Medicine in the University of Pennsylvania.

"Contains in concise and reliable form the accepted views of Ophthalmic Science."

WILLIAM THOMSON, M. D.,

Professor of Ophthalmology, Jefferson Medical College, Philadelphia, Pa.

TEXT-BOOK UPON THE PATHOGENIC BACTERIA. Specially written for Students of Medicine. By JOSEPH MCFARLAND, M. D., Professor of Pathology and Bacteriology in the Medico-Chirurgical College of Philadelphia, etc. 497 pages, finely illustrated. Price, Cloth, \$2.50 net.

SECOND EDITION, REVISED AND GREATLY ENLARGED.

The work is intended to be a text-book for the medical student and for the practitioner who has had no recent laboratory training in this department of medical science. The instructions given as to needed apparatus, cultures, stainings, microscopic examinations, etc. are ample for the student's needs, and will afford to the physician much information that will interest and profit him relative to a subject which modern science shows to go far in explaining the etiology of many diseased conditions.

In this second edition the work has been brought up to date in all departments of the subject, and numerous additions have been made to the technique in the endeavor to make the book fulfil the double purpose of a systematic work upon bacteria and a laboratory guide.

"It is excellently adapted for the medical students and practitioners for whom it is avowedly written. . . . The descriptions given are accurate and readable, and the book should prove useful to those for whom it is written.—*London Lancet*, Aug. 29, 1896.

"The author has succeeded admirably in presenting the essential details of bacteriological technics, together with a judiciously chosen summary of our present knowledge of pathogenic bacteria. . . . The work, we think, should have a wide circulation among English-speaking students of medicine."—*N. Y. Medical Journal*, April 4, 1896.

"The book will be found of considerable use by medical men who have not had a special bacteriological training, and who desire to understand this important branch of medical science."—*Edinburgh Medical Journal*, July, 1896.

LABORATORY GUIDE FOR THE BACTERIOLOGIST. By LANGDON FROTHINGHAM, M. D. V., Assistant in Bacteriology and Veterinary Science, Sheffield Scientific School, Yale University. Illustrated. Price, Cloth, 75 cents.

The technical methods involved in bacteria-culture, methods of staining, and microscopical study are fully described and arranged as simply and concisely as possible. The book is especially intended for use in laboratory work

"It is a convenient and useful little work, and will more than repay the outlay necessary for its purchase in the saving of time which would otherwise be consumed in looking up the various points of technique so clearly and concisely laid down in its pages."—*American Med.-Surg. Bulletin*.

FEEDING IN EARLY INFANCY. By ARTHUR V. MEIGS, M. D. Bound in limp cloth, flush edges. Price, 25 cents net.

SYNOPSIS: Analyses of Milk—Importance of the Subject of Feeding in Early Infancy—Proportion of Casein and Sugar in Human Milk—Time to Begin Artificial Feeding of Infants—Amount of Food to be Administered at Each Feeding—Intervals between Feedings—Increase in Amount of Food at Different Periods of Infant Development—Unsuitableness of Condensed Milk as a Substitute for Mother's Milk—Objections to Sterilization or "Pasteurization" of Milk—Advances made in the Method of Artificial Feeding of Infants.

ESSENTIALS OF ANATOMY AND MANUAL OF PRACTICAL DISSECTION, containing "Hints on Dissection." By CHARLES B. NANCREDE, M. D., Professor of Surgery and Clinical Surgery in the University of Michigan, Ann Arbor; Corresponding Member of the Royal Academy of Medicine, Rome, Italy; late Surgeon Jefferson Medical College, etc. Fourth and revised edition. Post 8vo, over 500 pages, with handsome full-page lithographic plates in colors, and over 200 illustrations. Price: Extra Cloth or Oilcloth for the dissection-room, \$2.00 net.

Neither pains nor expense has been spared to make this work the most exhaustive yet concise Student's Manual of Anatomy and Dissection ever published, either in America or in Europe.

The colored plates are designed to aid the student in dissecting the muscles, arteries, veins, and nerves. The wood-cuts have all been specially drawn and engraved, and an Appendix added containing 60 illustrations representing the structure of the entire human skeleton, the whole being based on the eleventh edition of Gray's *Anatomy*.

"The plates are of more than ordinary excellence, and are of especial value to students in their work in the dissecting-room."—*Journal of American Medical Association*.

"Should be in the hands of every medical student."—*Cleveland Medical Gazette*.

"A concise and judicious work."—*Buffalo Medical and Surgical Journal*.

A MANUAL OF PRACTICE OF MEDICINE. By A. A. STEVENS, A. M., M. D., Instructor of Physical Diagnosis in the University of Pennsylvania, and Demonstrator of Pathology in the Woman's Medical College of Philadelphia. Specially intended for students preparing for graduation and hospital examinations, and includes the following sections: General Diseases, Diseases of the Digestive Organs, Diseases of the Respiratory System, Diseases of the Circulatory System, Diseases of the Nervous System, Diseases of the Blood, Diseases of the Kidneys, and Diseases of the Skin. Each section is prefaced by a chapter on General Symptomatology. Post 8vo, 520 pages. Numerous illustrations and selected formulæ. Price, \$2.50, bound in flexible leather.

FIFTH EDITION, REVISED AND ENLARGED.

Contributions to the science of medicine have poured in so rapidly during the last quarter of a century that it is well-nigh impossible for the student, with the limited time at his disposal, to master elaborate treatises or to cull from them that knowledge which is absolutely essential. From an extended experience in teaching, the author has been enabled, by classification, to group allied symptoms, and by the judicious elimination of theories and redundant explanations to bring within a comparatively small compass a complete outline of the practice of medicine.

MANUAL OF MATERIA MEDICA AND THERAPEUTICS.

By A. A. STEVENS, A. M., M. D., Instructor of Physical Diagnosis in the University of Pennsylvania, and Demonstrator of Pathology in the Woman's Medical College of Philadelphia. 445 pages. Price, Cloth, \$2.25.

SECOND EDITION, REVISED.

This wholly new volume, which is based on the last edition of the *Pharmacopœia*, comprehends the following sections: Physiological Action of Drugs; Drugs; Remedial Measures other than Drugs; Applied Therapeutics; Incompatibility in Prescriptions; Table of Doses; Index of Drugs; and Index of Diseases; the treatment being elucidated by more than two hundred formulæ.

"The author is to be congratulated upon having presented the medical student with as accurate a manual of therapeutics as it is possible to prepare."—*Therapeutic Gazette*.

"Far superior to most of its class; in fact, it is very good. Moreover, the book is reliable and accurate."—*New York Medical Journal*.

"The author has faithfully presented modern therapeutics in a comprehensive work, . . . and it will be found a reliable guide."—*University Medical Magazine*.

NOTES ON THE NEWER REMEDIES: their Therapeutic Applications and Modes of Administration. By DAVID CERNA, M. D., PH. D., Demonstrator of and Lecturer on Experimental Therapeutics in the University of Pennsylvania. Post-octavo, 253 pages. Price, \$1.25.

SECOND EDITION, RE-WRITTEN AND GREATLY ENLARGED.

The work takes up in alphabetical order all the newer remedies, giving their physical properties, solubility, therapeutic applications, administration, and chemical formula.

It thus forms a very valuable addition to the various works on therapeutics now in existence.

Chemists are so multiplying compounds, that, if each compound is to be thoroughly studied, investigations must be carried far enough to determine the practical importance of the new agents.

"Especially valuable because of its completeness, its accuracy, its systematic consideration of the properties and therapy of many remedies of which doctors generally know but little, expressed in a brief yet terse manner."—*Chicago Clinical Review*.

TEMPERATURE CHART. Prepared by D. T. LAINÉ, M. D. Size 8 × 13½ inches. Price, per pad of 25 charts, 50 cents.

A conveniently arranged chart for recording Temperature, with columns for daily amounts of Urinary and Fecal Excretions, Food, Remarks, etc. On the back of each chart is given in full the method of Brand in the treatment of Typhoid Fever.

SAUNDERS' POCKET MEDICAL LEXICON; or, Dictionary of Terms and Words used in Medicine and Surgery. By JOHN M. KEATING, M. D., editor of "Cyclopædia of Diseases of Children," etc.; author of the "New Pronouncing Dictionary of Medicine;" and HENRY HAMILTON, author of "A New Translation of Virgil's *Æneid* into English Verse;" co-author of a "New Pronouncing Dictionary of Medicine." A new and revised edition. 32mo, 282 pages. Prices: Cloth, 75 cents; Leather Tucks, \$1.00.

This new and comprehensive work of reference is the outcome of a demand for a more modern handbook of its class than those at present on the market, which, dating as they do from 1855 to 1884, are of but trifling use to the student by their not containing the hundreds of new words now used in current literature, especially those relating to Electricity and Bacteriology.

"Remarkably accurate in terminology, accentuation, and definition."—*Journal of American Medical Association*.

"Brief, yet complete . . . it contains the very latest nomenclature in even the newest departments of medicine."—*New York Medical Record*.

SAUNDERS' POCKET MEDICAL FORMULARY. By WILLIAM M. POWELL, M. D., Attending Physician to the Mercer House for Invalid Women at Atlantic City. Containing 1800 Formulæ, selected from several hundred of the best-known authorities. Forming a handsome and convenient pocket companion of nearly 300 printed pages, with blank leaves for Additions; with an Appendix containing Posological Table, Formulæ and Doses for Hypodermatic Medication, Poisons and their Antidotes, Diameters of the Female Pelvis and Fœtal Head, Obstetrical Table, Diet List for Various Diseases, Materials and Drugs used in Antiseptic Surgery, Treatment of Asphyxia from Drowning, Surgical Remembrancer, Tables of Incompatibles, Eruptive Fevers, Weights and Measures, etc. Fourth edition, revised and greatly enlarged. Handsomely bound in morocco, with side index, wallet, and flap. Price, \$1.75 net.

A concise, clear, and correct record of the many hundreds of famous formulæ which are found scattered through the works of the *most eminent physicians and surgeons* of the world. The work is helpful to the student and practitioner alike, as through it they become acquainted with numerous formulæ which are not found in text-books, but have been collected from among *the rising generation of the profession, college professors, and hospital physicians and surgeons*.

"This little book, that can be conveniently carried in the pocket, contains an immense amount of material. It is very useful, and as the name of the author of each prescription is given is unusually reliable."—*New York Medical Record*.

"Designed to be of immense help to the general practitioner in the exercise of his daily calling."—*Boston Medical and Surgical Journal*.

DISEASES OF WOMEN. By HENRY J. GARRIGUES, A. M., M. D., Professor of Gynecology and Obstetrics in the New York School of Clinical Medicine; Gynecologist to St. Mark's Hospital and to the German Dispensary, New York City. In one handsome octavo volume of 728 pages, illustrated by 335 engravings and colored plates. Prices: Cloth, \$4.00 net; Sheep or Half Morocco, \$5.00 net.

A PRACTICAL work on gynecology for the use of students and practitioners, written in a terse and concise manner. The importance of a thorough knowledge of the anatomy of the female pelvic organs has been fully recognized by the author, and considerable space has been devoted to the subject. The chapters on Operations and on Treatment are thoroughly modern, and are based upon the large hospital and private practice of the author. The text is elucidated by a large number of illustrations and colored plates, many of them being original, and forming a complete atlas for studying *embryology* and the *anatomy* of the *female genitalia*, besides exemplifying, whenever needed, morbid conditions, instruments, apparatus, and operations.

Second Edition, Thoroughly Revised.

The first edition of this work met with a most appreciative reception by the medical press and profession both in this country and abroad, and was adopted as a text-book or recommended as a book of reference by nearly *one hundred* colleges in the United States and Canada. The author has availed himself of the opportunity afforded by this revision to embody the latest approved advances in the treatment employed in this important branch of Medicine. He has also more extensively expressed his own opinion on the comparative value of the different methods of treatment employed.

"One of the best text-books for students and practitioners which has been published in the English language; it is condensed, clear, and comprehensive. The profound learning and great clinical experience of the distinguished author find expression in this book in a most attractive and instructive form. Young practitioners, to whom experienced consultants may not be available, will find in this book invaluable counsel and help."

THAD. A. REAMY, M. D., LL.D.,

Professor of Clinical Gynecology, Medical College of Ohio; Gynecologist to the Good Samaritan and Cincinnati Hospitals.

A SYLLABUS OF GYNECOLOGY, arranged in conformity with "An American Text-Book of Gynecology." By J. W. LONG, M. D., Professor of Diseases of Women and Children, Medical College of Virginia, etc. Price, Cloth (interleaved), \$1.00 net.

Based upon the teaching and methods laid down in the larger work, this will not only be useful as a supplementary volume, but to those who do not already possess the text-book it will also have an independent value as an aid to the practitioner in gynecological work, and to the student as a guide in the lecture-room, as the subject is presented in a manner at once systematic, clear, succinct, and practical.

A MANUAL OF PHYSIOLOGY, with Practical Exercises. For Students and Practitioners. By G. N. STEWART, M. A., M. D., D. Sc., lately Examiner in Physiology, University of Aberdeen, and of the New Museums, Cambridge University; Professor of Physiology in the Western Reserve University, Cleveland, Ohio. Handsome octavo volume of 800 pages, with 278 illustrations in the text, and 5 colored plates. Price, Cloth, \$3.50 net.

"It will make its way by sheer force of merit, and *amply deserves to do so. It is one of the very best English text-books on the subject.*"—*London Lancet.*

"Of the many text-books of physiology published, we do not know of one that so nearly comes up to the ideal as does Professor Stewart's volume."—*British Medical Journal.*

ESSENTIALS OF PHYSICAL DIAGNOSIS OF THE THORAX.

By ARTHUR M. CORWIN, A. M., M. D., Demonstrator of Physical Diagnosis in the Rush Medical College, Chicago; Attending Physician to the Central Free Dispensary, Department of Rhinology, Laryngology, and Diseases of the Chest. 200 pages. Illustrated. Cloth, flexible covers. Price, \$1.25 net.

SYLLABUS OF OBSTETRICAL LECTURES in the Medical Department, University of Pennsylvania. By RICHARD C. NORRIS, A. M., M. D., Lecturer on Clinical and Operative Obstetrics, University of Pennsylvania. Third edition, thoroughly revised and enlarged. Crown 8vo. Price, Cloth, interleaved for notes, \$2.00 net.

"This work is so far superior to others on the same subject that we take pleasure in calling attention briefly to its excellent features. It covers the subject thoroughly, and will prove invaluable both to the student and the practitioner. The author has introduced a number of valuable hints which would only occur to one who was himself an experienced teacher of obstetrics. The subject-matter is clear, forcible, and modern. We are especially pleased with the portion devoted to the practical duties of the accoucheur, care of the child, etc. The paragraphs on antiseptics are admirable; there is no doubtful tone in the directions given. No details are regarded as unimportant; no minor matters omitted. We venture to say that even the old practitioner will find useful hints in this direction which he cannot afford to despise."—*New York Medical Record.*

A SYLLABUS OF LECTURES ON THE PRACTICE OF SURGERY, arranged in conformity with "An American Text-Book of Surgery." By N. SENN, M. D., PH. D., Professor of Surgery in Rush Medical College, Chicago, and in the Chicago Polyclinic. Price, \$2.00.

This, the latest work of its eminent author, himself one of the contributors to "An American Text-Book of Surgery," will prove of exceptional value to the advanced student who has adopted that work as his text-book. It is not only the syllabus of an unrivalled course of surgical practice, but it is also an epitome of or supplement to the larger work.

"The author has evidently spared no pains in making his Syllabus thoroughly comprehensive, and has added new matter and alluded to the most recent authors and operations. Full references are also given to all requisite details of surgical anatomy and pathology."—*British Medical Journal, London.*

AN OPERATION BLANK, with Lists of Instruments, etc. required in Various Operations. Prepared by W. W. KEEN, M. D., LL.D., Professor of Principles of Surgery in the Jefferson Medical College, Philadelphia. Price per Pad, containing Blanks for fifty operations, 50 cents net.

SECOND EDITION, REVISED FORM.

A convenient blank, suitable for all operations, giving complete instructions regarding necessary preparation of patient, etc., with a full list of dressings and medicines to be employed.

On the back of each blank is a list of instruments used—viz. general instruments, etc., required for all operations; and special instruments for surgery of the brain and spine, mouth and throat, abdomen, rectum, male and female genito-urinary organs, the bones, etc.

The whole forming a neat pad, arranged for hanging on the wall of a surgeon's office or in the hospital operating-room.

"Will serve a useful purpose for the surgeon in reminding him of the details of preparation for the patient and the room as well as for the instruments, dressings, and antiseptics needed."—*New York Medical Record*

"Covers about all that can be needed in any operation."—*American Lancet*.

"The plan is a capital one."—*Boston Medical and Surgical Journal*.

LABORATORY EXERCISES IN BOTANY. By EDSON S. BASTIN, M. A., Professor of Materia Medica and Botany in the Philadelphia College of Pharmacy. Octavo volume of 536 pages, 87 full-page plates. Price, Cloth, \$2.50.

This work is intended for the beginner and the advanced student, and it fully covers the structure of flowering plants, roots, ordinary stems, rhizomes, tubers, bulbs, leaves, flowers, fruits, and seeds. Particular attention is given to the gross and microscopical structure of plants, and to those used in medicine. Illustrations have freely been used to elucidate the text, and a complete index to facilitate reference has been added.

"There is no work like it in the pharmaceutical or botanical literature of this country, and we predict for it a wide circulation."—*American Journal of Pharmacy*.

DIET IN SICKNESS AND IN HEALTH. By MRS. ERNEST HART, formerly Student of the Faculty of Medicine of Paris and of the London School of Medicine for Women; with an INTRODUCTION by Sir Henry Thompson, F. R. C. S., M. D., London. 220 pages; illustrated. Price, Cloth, \$1.50.

Useful to those who have to nurse, feed, and prescribe for the sick. In each case the accepted causation of the disease and the reasons for the special diet prescribed are briefly described. Medical men will find the dietaries and recipes practically useful, and likely to save them trouble in directing the dietetic treatment of patients.

HOW TO EXAMINE FOR LIFE INSURANCE. By JOHN M. KEATING, M. D., Fellow of the College of Physicians and Surgeons of Philadelphia; Vice-President of the American Pædiatric Society; Ex-President of the Association of Life Insurance Medical Directors. Royal 8vo, 211 pages, with two large half-tone illustrations, and a plate prepared by Dr. McClellan from special dissections; also, numerous cuts to elucidate the text. Second edition. Price, Cloth, \$2.00 net.

"This is by far the most useful book which has yet appeared on insurance examination, a subject of growing interest and importance. Not the least valuable portion of the volume is Part II., which consists of instructions issued to their examining physicians by twenty-four representative companies of this country. As the proofs of these instructions were corrected by the directors of the companies, they form the latest instructions obtainable. If for these alone, the book should be at the right hand of every physician interested in this special branch of medical science."—*The Medical News*, Philadelphia.

NURSING: ITS PRINCIPLES AND PRACTICE. By ISABEL ADAMS HAMPTON, Graduate of the New York Training School for Nurses attached to Bellevue Hospital; Superintendent of Nurses and Principal of the Training School for Nurses, Johns Hopkins Hospital, Baltimore, Md.; late Superintendent of Nurses, Illinois Training School for Nurses, Chicago, Ill. In one very handsome 12mo volume of 484 pages, profusely illustrated. Price, Cloth, \$2.00 net.

This original work on the important subject of nursing is at once comprehensive and systematic. It is written in a clear, accurate, and readable style, suitable alike to the student and the lay reader. Such a work has long been a desideratum with those entrusted with the management of hospitals and the instruction of nurses in training-schools. It is also of especial value to the graduated nurse who desires to acquire a practical working knowledge of the care of the sick and the hygiene of the sick-room.

OBSTETRIC ACCIDENTS, EMERGENCIES, AND OPERATIONS. By L. CH. BOISLINIERE, M. D., late Emeritus Professor of Obstetrics in the St. Louis Medical College. 381 pages, handsomely illustrated. Price, \$2.00 net.

"For the use of the practitioner who, when away from home, has not the opportunity of consulting a library or of calling a friend in consultation. He then, being thrown upon his own resources, will find this book of benefit in guiding and assisting him in emergencies."

INFANT'S WEIGHT CHART. Designed by J. P. CROZER GRIFFITH, M. D., Clinical Professor of Diseases of Children in the University of Pennsylvania. 25 charts in each pad. Price per pad, 50 cents net.

A convenient blank for keeping a record of the child's weight during the first two years of life. Printed on each chart is a curve representing the average weight of a healthy infant, so that any deviation from the normal can readily be detected.

THE CARE OF THE BABY. By J. P. CROZER GRIFFITH, M. D., Clinical Professor of Diseases of Children, University of Pennsylvania; Physician to the Children's Hospital, Philadelphia, etc. 404 pages, with 67 illustrations in the text, and 5 plates. 12mo. Price, \$1.50.

SECOND EDITION, REVISED.

A reliable guide not only for mothers, but also for medical students and practitioners whose opportunities for observing children have been limited.

"The whole book is characterized by rare good sense, and is evidently written by a master hand. It can be read with benefit not only by mothers, but by medical students and by any practitioners who have not had large opportunities for observing children."—*American Journal of Obstetrics*.

THE NURSE'S DICTIONARY of Medical Terms and Nursing Treatment, containing Definitions of the Principal Medical and Nursing Terms, Abbreviations, and Physiological Names, and Descriptions of the Instruments, Drugs, Diseases, Accidents, Treatments, Operations, Foods, Appliances, etc. encountered in the ward or the sick-room. By HONNOR MORTEN, author of "How to Become a Nurse," "Sketches of Hospital Life," etc. 16mo, 140 pages. Price, Cloth, \$1.00.

This little volume is intended for use merely as a small reference-book which can be consulted at the bedside or in the ward. It gives sufficient explanation to the nurse to enable her to comprehend a case until she has leisure to look up larger and fuller works on the subject.

DIET LISTS AND SICK-ROOM DIETARY. By JEROME B. THOMAS, M. D., Visiting Physician to the Home for Friendless Women and Children and to the Newsboys' Home; Assistant Visiting Physician to the Kings County Hospital; Assistant Bacteriologist, Brooklyn Health Department. Price, Cloth, \$1.50 (Send for specimen List.)

One hundred and sixty detachable (perforated) diet lists for Albuminuria, Anæmia and Debility, Constipation, Diabetes, Diarrhoea, Dyspepsia, Fevers, Gout or Uric-Acid Diathesis, Obesity, and Tuberculosis. Also forty detachable sheets of Sick-Room Dietary, containing full instructions for preparation of easily-digested foods necessary for invalids. Each list is *numbered only*, the disease for which it is to be used in no case being mentioned, an index key being reserved for the physician's private use.

DIETS FOR INFANTS AND CHILDREN IN HEALTH AND IN DISEASE. By LOUIS STARR, M. D., Editor of "An American Text-Book of the Diseases of Children." 230 blanks (pocket-book size), perforated and neatly bound in flexible morocco. Price, \$1.25 net.

The first series of blanks are prepared for the first seven months of infant life; each blank indicates the ingredients, but not the *quantities*, of the food, the latter directions being left for the physician. After the seventh month, modifications being less necessary, the diet lists are printed in full. *Formula* for the preparation of diluents and foods are appended.



SAUNDERS' for Students NEW SERIES and OF MANUALS Practitioners.

THAT there exists a need for thoroughly reliable hand-books on the leading branches of Medicine and Surgery is a fact amply demonstrated by the favor with which the SAUNDERS NEW SERIES OF MANUALS have been received by medical students and practitioners and by the Medical Press. These manuals are not merely condensations from present literature, but are ably written by well-known authors and practitioners, most of them being teachers in representative American colleges. Each volume is concisely and authoritatively written and exhaustive in detail, without being encumbered with the introduction of "cases," which so largely expand the ordinary text-book. These manuals will therefore form an admirable collection of advanced lectures, useful alike to the medical student and the practitioner: to the latter, too busy to search through page after page of elaborate treatises for what he wants to know, they will prove of inestimable value; to the former they will afford safe guides to the essential points of study.

The SAUNDERS NEW SERIES OF MANUALS are conceded to be superior to any similar books now on the market. No other manuals afford so much information in such a concise and available form. A liberal expenditure has enabled the publisher to render the mechanical portion of the work worthy of the high literary standard attained by these books.

Any of these Manuals will be mailed on receipt of price (see next page for List).

W. B. SAUNDERS, Publisher,
925 Walnut Street, Philadelphia.

SAUNDERS' NEW SERIES OF MANUALS.

VOLUMES PUBLISHED.

PHYSIOLOGY. By JOSEPH HOWARD RAYMOND, A. M., M. D., Professor of Physiology and Hygiene and Lecturer on Gynecology in the Long Island College Hospital, etc. Price, \$1.25 net.

SURGERY, General and Operative. By JOHN CHALMERS DACOSTA, M. D., Demonstrator of Surgery, Jefferson Medical College, Philadelphia, etc. Octavo, 911 pages, 386 illustrations. Cloth, \$4.00 net; Half-Morocco, \$5.00 net.

DOSE-BOOK AND MANUAL OF PRESCRIPTION-WRITING. By E. Q. THORNTON, M. D., Demonstrator of Therapeutics, Jefferson Medical College, Philadelphia. Price, \$1.25 net.

MEDICAL JURISPRUDENCE. By HENRY C. CHAPMAN, M. D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia, etc. Price, \$1.50 net.

SURGICAL ASEPSIS. By CARL BECK, M. D., Surgeon to St. Mark's Hospital and to the German Poliklinik; Instructor in Surgery, New York Post-Graduate Medical School, etc. Price, \$1.25 net.

MANUAL OF ANATOMY. By IRVING S. HAYNES, M. D., Adjunct Professor of Anatomy and Demonstrator of Anatomy, Medical Department of the New York University, etc. (Double number.) Price, \$2.50 net.

SYPHILIS AND THE VENEREAL DISEASES. By JAMES NEVINS HYDE, M. D., Professor of Skin and Venereal Diseases, and FRANK H. MONTGOMERY, M. D., Lecturer on Dermatology and Genito-urinary Diseases, in Rush Medical College, Chicago. (Double number.) Price, \$2.50 net.

PRACTICE OF MEDICINE. By GEORGE ROE LOCKWOOD, M. D., Professor of Practice in the Woman's Medical College of the New York Infirmary, etc. (Double number.) Price, \$2.50 net.

OBSTETRICS. By W. A. NEWMAN DORLAND, M. D., Asst. Demonstrator of Obstetrics, University of Pennsylvania; Chief of Gynecological Dispensary, Pennsylvania Hospital. (Double number.) Price, \$2.50 net.

DISEASES OF WOMEN. By J. BLAND SUTTON, F. R. C. S., Assistant Surgeon to the Middlesex Hospital, and Surgeon to the Chelsea Hospital for Women, London; and ARTHUR E. GILES, M. D., B. Sc. Lond., F. R. C. S. Edin., Assistant Surgeon to the Chelsea Hospital for Women, London. 436 pages, handsomely illustrated. (Double number.) Price, \$2.50 net.

VOLUMES IN PREPARATION.

NERVOUS DISEASES. By CHARLES W. BURR, M. D., Clinical Professor of Nervous Diseases, Medico-Chirurgical College, Philadelphia, etc.

NOSE AND THROAT. By D. BRADEN KYLE, M. D., Clinical Professor of Laryngology and Rhinology, Jefferson Medical College, Philadelphia.

. There will be published in the same series, at short intervals, carefully prepared works on various subjects, by prominent specialists.

SAUNDERS' QUESTION COMPENDS.

Arranged in Question and Answer Form.

THE LATEST, MOST COMPLETE, and BEST ILLUSTRATED
SERIES OF COMPENDS EVER ISSUED.

Now the Standard Authorities in Medical Literature

WITH

Students and Practitioners in every City of the United
States and Canada.

THE REASON WHY.

They are the advance guard of "Student's Helps"—that DO HELP; they are the leaders in their special line, *well and authoritatively written by able men, who, as teachers in the large colleges, know exactly what is wanted by a student preparing for his examinations.* The judgment exercised in the selection of authors is fully demonstrated by their professional elevation. Chosen from the ranks of Demonstrators, Quiz-masters, and Assistants, most of them have become Professors and Lecturers in their respective colleges.

Each book is of convenient size (5×7 inches), containing on an average 250 pages, profusely illustrated, and elegantly printed in clear, readable type, on fine paper.

The entire series, numbering twenty-four subjects, has been kept thoroughly revised and enlarged when necessary, many of them being in their fourth and fifth editions.

TO SUM UP.

Although there are numerous other Quizzes, Manuals, Aids, etc. in the market, none of them approach the "Blue Series of Question Compends;" and the claim is made for the following points of excellence:

1. Professional distinction and reputation of authors.
2. Conciseness, clearness, and soundness of treatment.
3. Size of type and quality of paper and binding.

..* Any of these Compends will be mailed on receipt of price (see over for List).

SAUNDERS' QUESTION-COMPEND SERIES.

Price, Cloth, \$1.00 per copy, except when otherwise noted.

1. **ESSENTIALS OF PHYSIOLOGY.** 3d edition. Illustrated. Revised and enlarged by H. A. HARE, M. D. (Price, \$1.00 net.)
2. **ESSENTIALS OF SURGERY.** 6th edition, with an Appendix on Antiseptic Surgery. 90 illustrations. By EDWARD MARTIN, M. D.
3. **ESSENTIALS OF ANATOMY.** 5th edition, with an Appendix. 180 illustrations. By CHARLES B. NANCREDÉ, M. D.
4. **ESSENTIALS OF MEDICAL CHEMISTRY, ORGANIC AND INORGANIC.** 4th edition, revised, with an Appendix. By LAWRENCE WOLFF, M. D.
5. **ESSENTIALS OF OBSTETRICS.** 4th edition, revised and enlarged. 75 illustrations. By W. EASTERLY ASHTON, M. D.
6. **ESSENTIALS OF PATHOLOGY AND MORBID ANATOMY.** 7th thousand. 46 illustrations. By C. E. ARMAND SEMPLE, M. D.
7. **ESSENTIALS OF MATERIA MEDICA, THERAPEUTICS, AND PRESCRIPTION-WRITING.** 4th edition. By HENRY MORRIS, M. D.
- 8, 9. **ESSENTIALS OF PRACTICE OF MEDICINE.** By HENRY MORRIS, M. D. An Appendix on URINE EXAMINATION. Illustrated. By LAWRENCE WOLFF, M. D. 3d edition, enlarged by some 300 Essential Formulæ, selected from eminent authorities, by WM. M. POWELL, M. D. (Double number, price \$2.00.)
10. **ESSENTIALS OF GYNÆCOLOGY.** 4th edition, revised. With 62 illustrations. By EDWIN B. CRAGIN, M. D.
11. **ESSENTIALS OF DISEASES OF THE SKIN.** 3d edition, revised and enlarged. 71 letter-press cuts and 15 half-tone illustrations. By HENRY W. STELWAGON, M. D. (Price, \$1.00 net.)
12. **ESSENTIALS OF MINOR SURGERY, BANDAGING, AND VENEREAL DISEASES.** 2d edition, revised and enlarged. 78 illustrations. By EDWARD MARTIN, M. D.
13. **ESSENTIALS OF LEGAL MEDICINE, TOXICOLOGY, AND HYGIENE.** 130 illustrations. By C. E. ARMAND SEMPLE, M. D.
14. **ESSENTIALS OF DISEASES OF THE EYE, NOSE, AND THROAT.** 124 illustrations. 2d edition, revised. By EDWARD JACKSON, M. D., and E. BALDWIN GLEASON, M. D.
15. **ESSENTIALS OF DISEASES OF CHILDREN.** Second edition. By WILLIAM H. POWELL, M. D.
5. **ESSENTIALS OF EXAMINATION OF URINE.** Colored "VOGEL SCALE," and numerous illustrations. By LAWRENCE WOLFF, M. D. (Price, 75 cents.)
17. **ESSENTIALS OF DIAGNOSIS.** By S. SOLIS-COHEN, M. D., and A. A. ESHNER, M. D. 55 illustrations, some in colors. (Price, \$1.50 net.)
18. **ESSENTIALS OF PRACTICE OF PHARMACY.** By L. E. SAYRE. 2d edition, revised.
20. **ESSENTIALS OF BACTERIOLOGY.** 3d edition. 82 illustrations. By M. V. BALL, M. D.
21. **ESSENTIALS OF NERVOUS DISEASES AND INSANITY.** 48 illustrations. 3d edition, revised. By JOHN C. SHAW, M. D.
22. **ESSENTIALS OF MEDICAL PHYSICS.** 155 illustrations. 2d edition, revised. By FRED J. BROCKWAY, M. D. (Price, \$1.00 net.)
23. **ESSENTIALS OF MEDICAL ELECTRICITY.** 65 illustrations. By DAVID D. STEWART, M. D., and EDWARD S. LAWRENCE, M. D.
24. **ESSENTIALS OF DISEASES OF THE EAR.** By E. B. GLEA-

RECENT PUBLICATIONS

PENROSE'S DISEASES OF WOMEN

A Text-Book of Diseases of Women. By CHARLES B. PENROSE, M.D., Ph.D., Professor of Gynecology in the University of Pennsylvania; Surgeon to the Gynecean Hospital, Philadelphia. Octavo volume of 529 pages, handsomely illustrated. Cloth, \$3.50 net.

"I shall value very highly the copy of Penrose's "Diseases of Women" received. I have already recommended it to my class as THE BEST book."—HOWARD A. KELLY, *Professor of Gynecology and Obstetrics, Johns Hopkins University, Baltimore, Md.*

SENN'S GENITO-URINARY TUBERCULOSIS

Tuberculosis of the Genito-Urinary Organs, Male and Female. By NICHOLAS SENN, M.D., Ph.D., LL.D., Professor of the Practice of Surgery and of Clinical Surgery, Rush Medical College, Chicago. Handsome octavo volume of 320 pages, illustrated. Cloth, \$3.00 net.

SUTTON AND GILES' DISEASES OF WOMEN

Diseases of Women. By J. BLAND SUTTON, F.R.C.S., Assistant Surgeon to Middlesex Hospital, and Surgeon to Chelsea Hospital, London; and ARTHUR E. GILES, M.D., B.Sc. Lond., F.R.C.S. Edin., Assistant Surgeon to Chelsea Hospital, London. 436 pages, handsomely illustrated. Cloth, \$2.50 net.

BUTLER'S MATERIA MEDICA, THERAPEUTICS, AND PHARMACOLOGY. Second Edition, Revised

A Text-Book of Materia Medica, Therapeutics, and Pharmacology. By GEORGE F. BUTLER, Ph.G., M.D., Professor of Materia Medica and of Clinical Medicine in the College of Physicians and Surgeons, Chicago; Professor of Materia Medica and Therapeutics, Northwestern University, Woman's Medical School, etc. Octavo, 858 pages, illustrated. Cloth, \$4.00 net; Sheep, \$5.00 net.

SAUNDBY'S RENAL AND URINARY DISEASES

Lectures on Renal and Urinary Diseases. By ROBERT SAUNDBY, M.D. Edin., Fellow of the Royal College of Physicians, London, and of the Royal Medico-Chirurgical Society; Physician to the General Hospital; Consulting Physician to the Eye Hospital and to the Hospital for Diseases of Women; Professor of Medicine in Mason College, Birmingham, etc. Octavo volume of 434 pages, with numerous illustrations and 4 colored plates. Cloth, \$2.50 net.

PYE'S BANDAGING

Elementary Bandaging and Surgical Dressing, with Directions Concerning the Immediate Treatment of Cases of Emergency. For the Use of Dressers and Nurses. By WALTER PYE, F.R.C.S., Late Surgeon to St. Mary's Hospital, London. Small 12mo, with over 80 illustrations. Cloth, flexible covers. Price, 75 cents net.

MALLORY AND WRIGHT'S PATHOLOGICAL TECHNIQUE

Pathological Technique. By FRANK B. MALLORY, A.M., M.D., Assistant Professor of Pathology, Harvard University Medical School; and JAMES H. WRIGHT, A.M., M.D., Instructor in Pathology, Harvard University Medical School. Octavo volume of 396 pages, handsomely illustrated. Cloth, \$2.50 net.

"I have been looking forward to the publication of this book, and I am glad to say that I find it to be a most useful laboratory and post-mortem guide, full of practical information, and well up to date."—WILLIAM H. WELCH, *Professor of Pathology, Johns Hopkins University, Baltimore, Md.*

ANDERS' PRACTICE OF MEDICINE. Second Edition

A Text-Book of the Practice of Medicine. By JAMES M. ANDERS, M.D., Ph.D., LL.D., Professor of the Practice of Medicine and of Clinical Medicine, Medico-Chirurgical College, Philadelphia. In one handsome octavo volume of 1287 pages, fully illustrated. Cloth, \$5.50 net; Sheep or Half Morocco, \$6.50 net.

ANOMALIES AND CURIOSITIES OF MEDICINE.

BY

GEORGE M. GOULD, M. D., AND WALTER L. PYLE, M. D.

Several years of exhaustive research have been spent by the authors in the great medical libraries of the United States and Europe in collecting the material for this work. **Medical literature of all ages and all languages** has been carefully searched, as a glance at the Bibliographic Index will show. The facts, which will be of **extreme value to the author and lecturer**, have been arranged and annotated, and full reference footnotes given, indicating whence they have been obtained.

In view of the persistent and dominant interest in the anomalous and curious, a **thorough and systematic collection** of this kind (the first of which the authors have knowledge) must have its own peculiar sphere of usefulness.

As a complete and authoritative **Book of Reference** it will be of value not only to members of the medical profession, but to all persons interested in general scientific, sociologic, and medico-legal topics; in fact, the general interest of the subject and the dearth of any complete work upon it make this volume **one of the most important literary innovations of the day.**

An especially valuable feature of the book consists of the **Indexing.** Besides a complete and comprehensive **General Index**, containing numerous cross-references to the subjects discussed, and the names of the authors of the more important reports, there is a convenient **Bibliographic Index** and a **Table of Contents.**

The plan has been adopted of printing the **topical headings in bold-face type**, the reader being thereby enabled to tell at a glance the subject-matter of any particular paragraph or page.

Illustrations have been freely employed throughout the work, there being 165 relief cuts and 130 half-tones in the text, and 12 colored and half-tone full-page plates—a total of over 320 separate figures.

The careful rendering of the text and references, the wealth of illustrations, the mechanical skill represented in the typography, the printing, and the binding, combine to make this book one of the most attractive medical publications ever issued.

Handsome Imperial Octavo Volume of 968 Pages.

PRICES: Cloth, \$6.00 net; Half Morocco, \$7.00 net.

JUST ISSUED

AN AMERICAN TEXT-BOOK OF GENITO-URINARY AND SKIN DISEASES

Edited by L. BOLTON BANGS, M.D., Late Professor of Genito-Urinary and Venereal Diseases, New York Post-Graduate Medical School and Hospital; and WILLIAM A. HARDAWAY, M.D., Professor of Diseases of the Skin, Missouri Medical College. Octavo volume of over 1200 pages, with 300 illustrations in the text, and 20 full-page colored plates. Prices: Cloth, \$7.00 net; Sheep or Half Morocco, \$8.00 net.

MOORE'S ORTHOPEDIC SURGERY

A Manual of Orthopedic Surgery. By JAMES E. MOORE, M.D., Professor of Orthopedics and Adjunct Professor of Clinical Surgery, University of Minnesota, College of Medicine and Surgery. 8vo, 356 pages, handsomely illustrated. Cloth, \$2.50 net.

MACDONALD'S SURGICAL DIAGNOSIS AND TREATMENT

Surgical Diagnosis and Treatment. By J. W. MACDONALD, M.D. Edin., L.R.C.S. Edin., Professor of the Practice of Surgery and of Clinical Surgery in Hamline University; Visiting Surgeon to St. Barnabas' Hospital, Minneapolis, etc. Octavo volume of 800 pages, handsomely illustrated. Cloth, \$5.00 net; Half Morocco, \$6.00 net.

CHAPIN ON INSANITY

A Compendium of Insanity. By JOHN B. CHAPIN, M.D., LL.D., Physician-in-Chief, Pennsylvania Hospital for the Insane; late Physician-Superintendent of the Willard State Hospital, New York, etc. 12mo., 234 pages, illustrated. Cloth, \$1.25 net.

KEEN ON THE SURGERY OF TYPHOID FEVER

The Surgical Complications and Sequels of Typhoid Fever. By WM. W. KEEN, M.D., LL.D., Professor of the Principles of Surgery and of Clinical Surgery, Jefferson Medical College, Philada. Octavo volume of 400 pages. Cloth, \$3.00 net.

VAN VALZAH AND NISBET'S DISEASES OF THE STOMACH

Diseases of the Stomach. By WILLIAM W. VAN VALZAH, M.D., Professor of General Medicine and Diseases of the Digestive System and the Blood, New York Polyclinic; and J. DOUGLAS NISBET, M.D., Adjunct Professor of General Medicine and Diseases of the Digestive System and the Blood, New York Polyclinic. Octavo volume of 674 pages, illustrated. Cloth, \$3.50 net.

IN PREPARATION

AN AMERICAN TEXT-BOOK OF DISEASES OF THE EYE, EAR, NOSE, AND THROAT

Edited by G. E. DE SCHWEINITZ, M.D., Professor of Ophthalmology in the Jefferson Medical College; and B. ALEXANDER RANDALL, M.D., Professor of Diseases of the Ear in the University of Pennsylvania and in the Philadelphia Polyclinic.

CHURCH AND PETERSON'S NERVOUS AND MENTAL DISEASES

Nervous and Mental Diseases. By ARCHIBALD CHURCH, M.D., Professor of Mental Diseases and Medical Jurisprudence, Northwestern University Medical School, Chicago; and FREDERICK PETERSON, M.D., Clinical Professor of Mental Diseases, Woman's Medical College, New York, etc.

KYLE ON THE NOSE AND THROAT

Diseases of the Nose and Throat. By D. BRADEN KYLE, M.D., Clinical Professor of Laryngology and Rhinology, Jefferson Medical College, Philadelphia; Consulting Laryngologist, Rhinologist, and Otologist, St. Agnes' Hospital, etc.

STENGEL'S PATHOLOGY

A Manual of Pathology. By ALFRED STENGEL, M.D., Physician to the Philadelphia Hospital; Professor of Clinical Medicine in the Woman's Medical College; Physician to the Children's Hospital, etc.

HIRST'S OBSTETRICS

A Text-Book of Obstetrics. By BARTON COOKE HIRST, M.D., Professor of Obstetrics, University of Pennsylvania.

HEISLER'S EMBRYOLOGY

A Text-Book of Embryology. By JOHN C. HEISLER, M.D., Professor of Anatomy, Medico-Chirurgical College, Philadelphia.

NOW READY, VOLUMES FOR 1896, 1897, 1898.

SAUNDERS'
AMERICAN YEAR-BOOK OF MEDICINE and SURGERY.

Edited by GEORGE M. GOULD, A. M., M. D.

Assisted by Eminent American Specialists and Teachers.

NOTWITHSTANDING the rapid multiplication of medical and surgical works, still these publications fail to meet fully the requirements of the *general physician*, inasmuch as he feels the need of something more than mere text-books of well-known principles of medical science. Mr. Saunders has long been impressed with this fact, which is confirmed by the unanimity of expression from the profession at large, as indicated by advices from his large corps of canvassers.

This deficiency would best be met by current journalistic literature, but most practitioners have scant access to this almost unlimited source of information, and the busy practiser has but little time to search out in periodicals the many interesting cases whose study would doubtless be of inestimable value in his practice. Therefore, a work which places before the physician in convenient form *an epitomization of this literature by persons competent to pronounce upon*

The Value of a Discovery or of a Method of Treatment

cannot but command his highest appreciation. It is this critical and judicial function that will be assumed by the Editorial staff of the "American Year-Book of Medicine and Surgery."

It is the special purpose of the Editor, whose experience peculiarly qualifies him for the preparation of this work, not only to review the contributions to American journals, but also the methods and discoveries reported in the leading medical journals of Europe, thus enlarging the survey and making the work characteristically **international**. These reviews will not simply be a series of undigested abstracts indiscriminately run together, nor will they be retrospective of "news" *one or two years old*, but the treatment presented will be *synthetic* and *dogmatic*, and will include **only what is new**. Moreover, through expert condensation by experienced writers these discussions will be

Comprised in a Single Volume of about 1200 Pages.

The work will be replete with **original** and **selected** illustrations skilfully reproduced, for the most part in Mr. Saunders' own studios established for the purpose, thus ensuring accuracy in delineation, affording efficient aids to a right comprehension of the text, and adding to the attractiveness of the volume. Prices: Cloth, \$6.50 net; Half Morocco, \$7.50 net.

W. B. SAUNDERS, Publisher,
925 Walnut Street, Philadelphia.

The knowledge gained is equal to a post-graduate course.

Uniform with the "American Text-Book" Series.





LIBRARY OF CONGRESS



00026101256

